Front propagation in anomalous diffusive media governed by time-fractional diffusion

Andrea Mentrelli^{a,b}, Gianni Pagnini^{b,c}

^a*Department of Mathematics &
Alma Mater Research Center on Applied Mathematics – (AM)²,
University of Bologna, via Saragozza 8, 40123 Bologna, Italy*
^b*BCAM – Basque Center for Applied Mathematics,
Alameda de Mazarredo 14, 48009 Bilbao, Basque Country – Spain*
^c*Ikerbasque – Basque Foundation for Science,
Calle de María Díaz de Haro 3, 48013 Bilbao, Basque Country – Spain*

Abstract

In this paper, a multi-dimensional model is proposed to study the propagation of random fronts in media in which anomalous diffusion takes place. The front position is obtained as the weighted mean of fronts calculated by means of the level set method, using as weight-function the probability density function which characterizes the anomalous diffusion process. Since anomalous diffusion is assumed to be governed by a time-fractional diffusion equation, its fundamental solution is the required probability density function. It is shown that this fundamental solution can be expressed in the multi-dimensional case in terms of the well-known $\mathcal{M}$ -Wright/Mainardi function, as in the one-dimensional case. Making use of this representation for the practical purpose of numerical evaluation, the propagation of random fronts in two-dimensional subdiffusive media is discussed and investigated.

Keywords: random front propagation, Level Set Method, multidimensional time-fractional diffusion, $\mathcal{M}$ -Wright/Mainardi function

2010 MSC: 60G22, 60K37, 35F21, 65M08

Email addresses: andrea.mentrelli@unibo.it (Andrea Mentrelli),
gpagnini@bcamath.org (Gianni Pagnini)

1. Introduction

Front propagation is of interest in several fields of applied science. In many cases, the propagating front is embedded in a medium characterized by a random motion and then by a diffusion process. In typical diffusion processes the mean square displacement of particles is a linear function of time. Nevertheless in some complex media diffusion processes show a non-linear relationship to time, as for example in biological systems [1, 2, 3], in fluids and plasmas [4] and in other systems [5]. In the latter case, the diffusion process is known as *anomalous diffusion* [6, 2, 1, 7].

The aim of the present paper is to extend to the multi-dimensional case a recently developed one-dimensional model useful for the study of the propagation of fronts in media in which anomalous diffusion takes place [8].

The main items of the proposed approach are:

- the *average front* position is modelled by means of the level set method (LSM) [9];
- the spreading of interface particles around the average front is described by the probability density function (PDF) that characterizes the diffusive process;
- the *effective front* is obtained as the weighted mean of average fronts, by using as weight-function the PDF of particle spreading.

This approach to the study of random fronts was first proposed – in a context in which anomalous diffusion was not involved – to investigate the evolution of the burnt mass fraction in turbulent premixed combustion [10], and has recently been applied to wildland fire propagation [11, 12, 13, 14, 15]. In these studies, the chosen PDF was the Gaussian density function [10, 11, 12, 13, 14, 15], or a convolution of the Gaussian with an other PDF [13, 14, 15]. In the present study, since anomalous diffusion is assumed to be modelled by the time-fractional diffusion equation, i.e. a diffusion-like equation in which the first order derivative

in time is replaced by a derivative operator of fractional order, the required PDF
30 to be used as weight-function is the fundamental solution, or Green's function,
of this family of fractional partial differential equations.

It is well known that in the one-dimensional case that Green's function is
related to the $\mathcal{M}$ -Wright/Mainardi function [16, 17, 18], which tends to the
Gaussian function as the fractional order of the time derivative tends to one.
35 The $\mathcal{M}$ -Wright/Mainardi function is a well known function for the evaluation
of which robust numerical algorithms exist [19]. However, the fundamental
solution of the multidimensional time-fractional diffusion equation has a cum-
bersome representation [20] so that it is rather complicated to deal with for any
practical purpose. Due to this difficulty, the extension discussed in this paper
40 of the one-dimensional case investigated in [8] to the multi-dimensional cases
is one of the highlights of the paper.

In the following it is shown that, by using an integral representation formula
of the $\mathcal{M}$ -Wright/Mainardi function [21], which is based on the one-dimensional
Gaussian PDF, Green's function for multi-dimensional cases (including the most
45 interesting two- and three-dimensional cases) can be represented by the same
integral formula involving the corresponding multi-dimensional Gaussian PDF.
This formula strongly simplifies the representation of Green's function provided
in [20].

As previously mentioned, this PDF is integrated in the proposed approach
50 that is based on the well-established LSM.

The LSM [22, 9] is one of the most widely used and successful tools for
computing and analyzing the motion of a front and it has been adopted in many
different problems, including turbulent premixed combustion [23], wildland fire
propagation [24], groundwater infiltration [25], biology [26] and material science
55 [27].

In the framework of the LSM, the location of the propagating front is ob-
tained as a level set (for example the zero level set) of an auxiliary function
solution of a Hamilton-Jacobi equation known as *level set equation*.

In this setting, sharp gradients and cusps can be easily accounted for and
60 the effects of curvature may be easily incorporated. Then the LSM is particularly useful to handle problems in which the speed of the evolving interface is dependent on the interface properties, such as curvature and normal direction, as well as on the boundary conditions at the interface location. Hence, it is suitable for problems in which the topology of the evolving interface changes in
65 time, like when topological merging and splitting occur.

In addition to these advantages, one key-feature of the level set method framework is that it allows to easily improve the physical model by progressively including more detailed interface effects that might influence the front propagation, as the dispersion effects of interface particles due to diffusive phenomena considered in this study.
70

The formalism of the proposed approach can be related to the Smoothed Particle Hydrodynamics (SPH) [28]. This framework is characterized by the robustness of its numerical implementation and it is well suited for many practical problems as fluid-structure interactions. However, the typical problems
75 of SPH, namely the choice of the kernel function and of the smoothing length, are here avoided because they are straightforwardly determined by the particle PDF. This can be considered a positive feature of the present approach.

The presented approach can be related also to approaches used in turbulent premixed combustion, namely the LSM/G-equation [23] or the so-called *turbulent flame closure model* first proposed by Zimont [29, 30]. A discussion about
80 how these frameworks are related to the one presented here might be found in [10].

Furthermore, the present approach is also connected to the so-called Stochastic Level Set Method (SLSM), developed and adopted in computer vision. Typical
85 problems in this research area consist in recovering a certain surface or region through a shape optimization framework. However, classical methods suffer from being sometimes stuck in local minima. To overcome this difficulty, the SLSM has been developed. This method combines stochastic motion and

the classical LSM [31, 32, 33] and adopts a decision mechanism. The SLSM
90 is quite close to the approach here proposed, as in both models the interface
location is obtained by solving the level set equation with a suitably random-
ized front speed. However, the final aims of the two methods are different, and
so is the mathematical construction. In fact, in typical computer vision prob-
lems the final aim is not to compute the average shape after a large number
95 of independent realizations, but to recover a certain shape with the frontline
upon the derivation of a proper stochastic differential equation. In the present
study, the interest is indeed focused on the average properties of the processes
for applications in physics and engineering. Even if it could be possible to have
average properties of the process obtained with the SLSM by performing the
100 ensemble average of many independent realizations, the approach here proposed
uses physical arguments to derive the PDF of the interface particle displacement
instead of focusing on the selection of the correct stochastic differential equa-
tion, so it can be regarded as the averaged counterpart of the method adopted
in computer vision. Despite this difference in the two models, the large avail-
105 able literature regarding SLSM is an important guideline for future derivation
of stochastic models within the approach proposed here.

In analogy with studies on diffusion processes that describe the properties of
the underlying stochastic processes by the particle PDF and averaged quantities,
here the motion of random fronts is described is well in terms of an ensemble
110 average resulting in the mentioned effective front.

The paper is organized as follows. Section 2 contains the description of
the LSM-based model used to study the propagation of a front embedded in a
diffusive environment. In Section 3, it is recalled the time-fractional diffusion
equation governing the dispersion of the interface particles and its solutions in
115 the one-, two- and three-dimensional cases are discussed. In particular, it is
shown a new representation of the Green's functions of the time-fractional dif-
fusion equation, based on the Gaussian PDF, which can conveniently be used
for computational purposes. Section 4 contains a discussion of the numerical

methodology adopted for the calculation of Green's functions as well as the tech-
 120 niques adopted for the computation of the effective average front. Numerical
 results are also presented, including plots of the $\mathcal{M}$ -Wright/Mainardi function
 for the one-, two- and three-dimensional cases as well as a selection of numer-
 ical results concerning two-dimensional front propagation in media in which
 anomalous diffusion takes place, restricting the attention to the case of *subdiffu-*
 125 *sive* processes, i.e. those processes whose particle displacement variance grows
 less than linearly in time. Finally, in Section 5, conclusions are drawn and some
 future developments are proposed.

2. Modelling approach

Let us consider a propagating interface composed of a sufficiently large num-
 130 ber of particles with random motion. Let $\bar{\mathbf{x}}_0$ be a point of the interface at the
 initial instant $t = 0$, and let $\mathbf{X}^\omega(t, \bar{\mathbf{x}}_0)$ be the ω -realization of the trajectory
 of a particle P_0 located in $\bar{\mathbf{x}}_0$ at $t = 0$, such that $\mathbf{X}^\omega(0, \bar{\mathbf{x}}_0) = \bar{\mathbf{x}}_0$. By using
 statistical mechanics formalism [34], the trajectory of particle P_0 is described
 by the one-particle density function $p_d^\omega(\mathbf{x}; t) = \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}}_0))$, where $\delta(\mathbf{x})$ is
 135 the Dirac δ -function and d represents the spatial dimension.

Let N_ω be the number of the independent realizations indexed by ω , denoting
 by $\langle \cdot \rangle$ the *ensemble average*, then the PDF of the displacement of the interface
 particle P_0 is written as

$$p_d(\mathbf{x}; t | \bar{\mathbf{x}}_0) = \frac{1}{N_\omega} \sum_{\omega=1}^{N_\omega} \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}}_0)) = \langle \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}}_0)) \rangle,$$

that can be understood as the PDF counterpart of the *empirical* distribution
 function [35]. If the density of particles embodying the interface is assumed
 to be constant during the interface evolution, then an incompressibility-like
 condition can be stated from the Jacobian J of the transformation $\bar{\mathbf{x}} \equiv \bar{\mathbf{x}}(t, \bar{\mathbf{x}}_0)$:
 $J = d\bar{\mathbf{x}}_0/d\bar{\mathbf{x}} = 1$. Finally, in terms of the average position $\bar{\mathbf{x}}$ of particle P_0 , it
 holds

$$p_d(\mathbf{x}; t | \bar{\mathbf{x}}) = \langle \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}})) \rangle.$$

Let Γ be a simple closed surface, or an ensemble of simple non-intersecting closed surfaces, that represents the propagating interface under consideration and let Ω be the region bounded by Γ . In the case of an interface Γ made of more than one closed surface, the domain Ω is not simply connected, resulting
140 in more than one bounded regions independently evolving. Moreover, let $\gamma : \mathcal{S} \times [0, +\infty[\rightarrow \mathbb{R}$ be a function defined on the domain of interest $\mathcal{S} \subseteq \mathbb{R}^d$ such that the level set $\gamma(\mathbf{x}, t) = \gamma_*$ coincides with the evolving front, i.e. $\Gamma(t) = \{\mathbf{x} \in \mathcal{S} \mid \gamma(\mathbf{x}, t) = \gamma_*\}$. In the case of Γ being an ensemble of n surfaces, the ensemble of the n interfaces is considered as *interface*.

In a deterministic setting, the level set function $\gamma(\mathbf{x}, t)$ may be written as

$$\gamma(\mathbf{x}, t) = \int_{\mathcal{S}} \gamma(\bar{\mathbf{x}}, t) \delta(\mathbf{x} - \bar{\mathbf{x}}) d\bar{\mathbf{x}},$$

and the subsets of the domain $\mathcal{S}$ corresponding to the interface Γ and to the region Ω enclosed by Γ (which represent, respectively, the interface and the domain bounded by it) may be conveniently described by two indicator functions $\mathcal{I}_{\Gamma}, \mathcal{I}_{\Omega} : \mathcal{S} \times [0, +\infty[\rightarrow \{0, 1\}$ defined as follows:

$$\mathcal{I}_{\Gamma}(\mathbf{x}, t) = \begin{cases} 1, & \text{if } \gamma(\mathbf{x}, t) = \gamma_* \\ 0, & \text{elsewhere,} \end{cases}$$

and

$$\mathcal{I}_{\Omega}(\mathbf{x}, t) = \begin{cases} 1, & \text{if } \gamma(\mathbf{x}, t) \leq \gamma_* \\ 0, & \text{elsewhere.} \end{cases} \quad (1)$$

145 The indicator functions at time $t = 0$, i.e. $\mathcal{I}_{\Gamma}(\mathbf{x}, t = 0)$ and $\mathcal{I}_{\Omega}(\mathbf{x}, t = 0)$, are denoted in the following as $\mathcal{I}_{\Gamma_0}(\mathbf{x})$ and $\mathcal{I}_{\Omega_0}(\mathbf{x})$, respectively.

In a random setting, the level set function corresponding to the ω -realization $\gamma^{\omega}(\mathbf{x}, t)$, which embeds the random front Γ^{ω} , for each realization is written as

$$\gamma^{\omega}(\mathbf{x}, t) = \int_{\mathcal{S}} \gamma(\bar{\mathbf{x}}, t) \delta(\mathbf{x} - \mathbf{X}^{\omega}(t, \bar{\mathbf{x}})) d\bar{\mathbf{x}}$$

and, accordingly, $\mathcal{I}_\Gamma$ and $\mathcal{I}_\Omega$ are replaced by random indicator functions $\mathcal{I}_{\Gamma^\omega}, \mathcal{I}_{\Omega^\omega} :$
 $\mathcal{S} \times [0, +\infty[\rightarrow \{0, 1\}$ defined as follows:

$$\begin{aligned}\mathcal{I}_{\Gamma^\omega}(\mathbf{x}, t) &= \int_{\mathcal{S}} \mathcal{I}_{\Gamma_0}(\bar{\mathbf{x}}_0) \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}}_0)) d\bar{\mathbf{x}}_0 \\ &= \int_{\Gamma_0} \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}}_0)) d\bar{\mathbf{x}}_0 \\ &= \int_{\Gamma(t)} \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}})) d\bar{\mathbf{x}},\end{aligned}$$

and

$$\begin{aligned}\mathcal{I}_{\Omega^\omega}(\mathbf{x}, t) &= \int_{\mathcal{S}} \mathcal{I}_{\Omega_0}(\bar{\mathbf{x}}_0) \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}}_0)) d\bar{\mathbf{x}}_0 \\ &= \int_{\Omega_0} \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}}_0)) d\bar{\mathbf{x}}_0 \\ &= \int_{\Omega(t)} \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}})) d\bar{\mathbf{x}}.\end{aligned}$$

In the case of a non-deterministic interface, an *effective indicator* of the region surrounded by a random front, $\varphi_e(\mathbf{x}, t) : \mathcal{S} \times [0, +\infty[\rightarrow [0, 1]$, was defined in [10] (in the context of modelling of the burnt mass fraction in turbulent premixed combustion) as follows:

$$\begin{aligned}\varphi_e(\mathbf{x}, t) &= \langle \mathcal{I}_{\Omega^\omega}(\mathbf{x}, t) \rangle = \left\langle \int_{\Omega(t)} \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}})) d\bar{\mathbf{x}} \right\rangle \\ &= \int_{\Omega(t)} \langle \delta(\mathbf{x} - \mathbf{X}^\omega(t, \bar{\mathbf{x}})) \rangle d\bar{\mathbf{x}} \\ &= \int_{\Omega(t)} p_d(\mathbf{x}; t | \bar{\mathbf{x}}) d\bar{\mathbf{x}}.\end{aligned}\tag{2}$$

It should be noted that the *effective indicator* $\varphi_e(\mathbf{x}, t)$ is not an indicator function in the classical sense. Actually, by using terminology from fuzzy logic, it is a *membership function*, its range being the compact interval $[0, 1]$ rather than the discrete set $\{0, 1\}$. Despite this, the concept of probability which led to Eq. (2) should not be confused with the concept of degree of truth (typical of fuzzy logic), then $\varphi_e(\mathbf{x}, t)$ is classified as an indicator function. Since the range of the effective indicator $\varphi_e(\mathbf{x}, t)$ is the compact interval $[0, 1]$, a criterion to mark the *effective* surrounded region Ω_e has to be stated. The region Ω_e is

155 here defined as the region over which the effective indicator exceeds a threshold value arbitrarily fixed φ_e^{th} , i.e. $\Omega_e(t) = \{\mathbf{x} \in \mathcal{S} \mid \varphi_e(\mathbf{x}, t) \geq \varphi_e^{th}\}$.

Making use of the indicator function $\mathcal{I}_\Omega$ defined in Eq. (1), Eq. (2) can be rewritten as:

$$\varphi_e(\mathbf{x}, t) = \int_{\mathcal{S}} \mathcal{I}_\Omega(\bar{\mathbf{x}}, t) p_d(\mathbf{x}; t \mid \bar{\mathbf{x}}) d\bar{\mathbf{x}}. \quad (3)$$

In the deterministic case, i.e. $\mathbf{X}^\omega(t, \bar{\mathbf{x}}) = \bar{\mathbf{x}}(t)$ for all realizations, it turns out that $p_d(\mathbf{x}; t \mid \bar{\mathbf{x}}) = \delta(\mathbf{x} - \bar{\mathbf{x}}(t))$, and from Eq. (3) it is obtained $\varphi_e(\mathbf{x}, t) = \mathcal{I}_\Omega(\mathbf{x}, t)$. Eq. (3) is similar to the basic equation of SPH method [28], however, as stated in
160 the Introduction, the choices of the kernel function and of the smoothing length are avoided because they are straightforwardly determined by the particle PDF $p_d(\mathbf{x}; t \mid \bar{\mathbf{x}})$.

In conclusion, the indicator function $\mathcal{I}_\Omega(\mathbf{x}, t)$ marks the region $\Omega(t)$ that is bounded by an interface propagating according to the average trajectory $\bar{\mathbf{x}}(t)$. Recalling the definition of $\mathcal{I}_\Omega(\mathbf{x}, t)$ in Eq. (1), the evolution of the level set function $\gamma(\mathbf{x}, t)$ is governed by the following Hamilton–Jacobi equation

$$\frac{\partial \gamma}{\partial t} = \mathcal{V}(\mathbf{x}, t) \|\nabla \gamma\|, \quad (4)$$

where γ_0 is the initial field embedding the interface Γ at $t = 0$, $\Gamma_0 \equiv \Gamma(t = 0)$.

It is worth mentioning here that an alternative representation of the dynamics of the propagating front is possible whenever velocity $\mathcal{V}(\mathbf{x}, t)$ is constant in time and strictly positive (or strictly negative). In this case, let $\tau(\mathbf{x})$ be the *arrival time* function that represents the temporal instant at which the front reaches the point $\mathbf{x}$, then $\|\nabla \tau\|$ is the rate of change of the arrival time with respect to the change in the front distance, i.e.

$$\|\nabla \tau\| = \frac{1}{\mathcal{V}}, \quad \tau(\mathbf{x}) = 0 \quad \forall \mathbf{x} \in \Gamma_0. \quad (5)$$

Equation (5) is known as eikonal equation and it is a time-independent version
165 of the level set equation (4). It has the important advantage of allowing – when applicable – to greatly reduce the computational effort required for solving the front tracking problem by solving a time-independent problem instead of a time-dependent one.

3. Multidimensional time-fractional diffusion equation

170 In the model here discussed for front propagation in complex media, the PDF of the interface particle displacement $p_d(\mathbf{x}; t)$, which appears in Eq. (3), is assumed to be governed by the d -dimensional time-fractional diffusion equation. The latter is obtained from the classical diffusion equation by replacing the first order time derivative with a real order derivative operator [36, 37, 38, 39, 20, 40].
 175 This replacement can be done by using the time fractional derivative operator in the Caputo sense or in the Riemann–Liouville sense. However, these two formulations are equivalent if standard initial condition is used [41]. Recently, in place of this operator replacement, the emergence of fractional kinetics in complex media has been explained within the standard Gaussian diffusion frame-
 180 work by the randomization due to the complexity of the environment of certain characteristic parameters of the medium [7].

Let $0 < \beta \leq 1/2$ be a real parameter, then the time-fractional diffusion equation in the Caputo sense reads

$${}_t^{2\beta} D_t^{\beta} p_d = \nu \nabla^2 p_d, \quad p_d(\mathbf{x}; 0) = \delta(\mathbf{x}), \quad (6)$$

where the coefficient ν is a positive constant with dimensions $[\nu] = [L]^2 [T]^{-2\beta}$ and ${}_t^{\mu} D_t^{\mu}$ is the Caputo fractional derivative that is defined by the Laplace transformation

$$\int_0^{+\infty} e^{-st} \{ {}_t^{\mu} D_t^{\mu} f(t) \} dt = s^{\mu} \tilde{f}(s) - \sum_{k=0}^{m-1} s^{\mu-1-k} f^{(k)}(0^+),$$

with $m - 1 < \mu \leq m$ and $m \in \mathbb{N}$, where $f(t)$ is a sufficiently well-behaved function and $\tilde{f}(s)$ its Laplace transform.

The time-fractional diffusion equation in the Riemann–Liouville sense reads

$$\frac{\partial p_d}{\partial t} = \nu D_t^{1-2\beta} \nabla^2 p_d, \quad p_d(\mathbf{x}; 0) = \delta(\mathbf{x}), \quad (7)$$

where D_t^{μ} is the Riemann–Liouville fractional derivative defined as

$$\int_0^{+\infty} e^{-st} \{ D_t^{\mu} f(t) \} dt = s^{\mu} \tilde{f}(s),$$

provided that all the limiting values $f^{(k)}(0^+)$ are finite with $k \in \mathbb{N}$ such that
185 $0 \leq k \leq m-1$ where $m \in \mathbb{N}$ and $m-1 < \mu \leq m$.

When $\beta = 1/2$, both equations (6) and (7) reduce to the classical diffusion equation

$$\frac{\partial p_d}{\partial t} = \nu \nabla^2 p_d, \quad p_d(\mathbf{x}; 0) = \delta(\mathbf{x}).$$

Applying Laplace and Fourier transformations to Eq. (6) as well as to Eq. (7), the fundamental solution $p_d(\mathbf{x}; t)$ for an arbitrary spatial dimension d results to be:

$$p_d(\mathbf{x}; t) = \frac{1}{(2\pi)^{d+1} i} \int_{-i\infty}^{+i\infty} \left\{ \int_{R^d} \frac{s^{2\beta-1}}{s^{2\beta} + \nu |\mathbf{k}|^2} e^{i\mathbf{k} \cdot \mathbf{x}} d\mathbf{k} \right\} e^{st} ds, \quad (8)$$

which can be written as [20]:

$$p_d(\mathbf{x}; t) = \frac{1}{(2\pi\sqrt{\nu})^{d/2+1} i} \int_{-i\infty}^{+i\infty} s^{\beta(d/2+1)-1} e^{st} \mathcal{K}_{d/2-1} \left(\frac{s^\beta}{\sqrt{\nu}} |\mathbf{x}| \right) ds, \quad (9)$$

where $\mathcal{K}_\theta$ is the Macdonald function, or modified Bessel function of the second kind, of order θ . By setting $s = -i\xi$, $\xi \in \mathbb{R}$, it follows that $s^\beta = e^{-i\pi\beta \operatorname{sgn} \xi/2} |\xi|^\beta$ and $p_d(\mathbf{x}; t)$ is real [20]. However, integral (9) is not straightforwardly evaluable and a less computationally demanding formulation is desirable. It is shown
190 in the following that such an alternative exists, since $p_d(\mathbf{x}; t)$ can be written in terms of the d -dimensional $\mathbb{M}$ -Wright/Mainardi function of order β here denoted by $\mathbb{M}_\beta^d(\mathbf{x}; t)$.

Let us start by reminding the following integral representation of the one-dimensional $\mathcal{M}$ -Wright/Mainardi function [21]:

$$\mathbb{M}_{\eta/2}^1(x; t) = \frac{1}{t^{\eta/2}} \mathcal{M}_{\eta/2} \left(\frac{x}{t^{\eta/2}} \right) = 2 \int_0^\infty \frac{e^{-x^2/(4\tau)}}{\sqrt{4\pi\tau}} \mathcal{M}_\eta \left(\frac{\tau}{t^\eta} \right) \frac{d\tau}{t^\eta}, \quad 0 < \eta < 1, x \in \mathbb{R}, \quad (10)$$

where the $\mathcal{M}$ -Wright/Mainardi function $\mathcal{M}_\eta(z)$ is represented by series as ($z \in \mathbb{C}$) [37, 38, 16, 17]

$$\mathcal{M}_\eta(z) = \sum_{n=0}^{\infty} \frac{(-z)^n}{n! \Gamma(-\eta n + (1-\nu))} = \frac{1}{\pi} \sum_{n=1}^{\infty} \frac{(-z)^{n-1}}{(n-1)!} \Gamma(\eta n) \sin(\pi \eta n).$$

Eq. (10) can be extended to a general spatial dimension d by the extension of the Gaussian function as follows ($0 < \eta \leq 1$, $\mathbf{x} \in \mathbb{R}^d$)

$$\begin{aligned}\mathbb{M}_{\eta/2}^d(\mathbf{x}; t) &= 2 \int_0^{+\infty} \frac{e^{-|\mathbf{x}|^2/(4\tau)}}{(4\pi\tau)^{d/2}} \mathcal{M}_\eta\left(\frac{\tau}{t^\eta}\right) \frac{d\tau}{t^\eta}, \\ &= 2 \int_0^{+\infty} \prod_{i=1}^d \frac{e^{-x_i^2/(4\tau)}}{(4\pi\tau)^{1/2}} \mathcal{M}_\eta\left(\frac{\tau}{t^\eta}\right) \frac{d\tau}{t^\eta}.\end{aligned}\quad (11)$$

When $d = 1$, Eq. (11) coincides with Eq. (10). Since the following Laplace transform pair holds [39]:

$$\int_0^{+\infty} e^{-st} \mathcal{M}_\eta\left(\frac{\tau}{t^\eta}\right) \frac{dt}{t^\eta} = s^{\eta-1} e^{-\tau s^\eta},$$

after the application of Laplace and Fourier transformations to (11) it follows that

$$\begin{aligned}\mathbb{M}_{\eta/2}^d(\mathbf{x}; t) &= \frac{2}{(2\pi)^{d+1} i} \int_{-i\infty}^{+i\infty} \left\{ \int_{\mathbb{R}^d} \left[\int_0^{+\infty} \prod_{i=1}^d e^{-\tau k_i^2} e^{-\tau s^\eta} d\tau \right] e^{i\mathbf{k} \cdot \mathbf{x}} d\mathbf{k} \right\} s^{\eta-1} e^{st} ds \\ &= \frac{2}{(2\pi)^{d+1} i} \int_{-i\infty}^{+i\infty} \left\{ \int_{\mathbb{R}^d} \left[\int_0^{+\infty} e^{-\tau(s^\eta + \nu k^2)} d\tau \right] e^{i\mathbf{k} \cdot \mathbf{x}} d\mathbf{k} \right\} s^{\eta-1} e^{st} ds \\ &= \frac{2}{(2\pi)^{d+1} i} \int_{-i\infty}^{+i\infty} \left\{ \int_{\mathbb{R}^d} \frac{s^{\eta-1}}{s^\eta + k^2} e^{i\mathbf{k} \cdot \mathbf{x}} d\mathbf{k} \right\} e^{st} ds,\end{aligned}$$

The latter, together with (8) and making use of the change of variables $k \rightarrow \sqrt{\nu} k$ and $\mathbf{x} \rightarrow \mathbf{x}/\sqrt{\nu}$, allows to conclude that

$$p_d(\mathbf{x}; t) = \frac{1}{2\sqrt{\nu}} \mathbb{M}_\beta^d\left(\frac{\mathbf{x}}{\sqrt{\nu}}; t\right) = \int_0^{+\infty} \frac{e^{-|\mathbf{x}|^2/(4\nu\tau)}}{(4\pi\nu\tau)^{d/2}} \mathcal{M}_{2\beta}\left(\frac{\tau}{t^{2\beta}}\right) \frac{d\tau}{t^{2\beta}}. \quad (12)$$

Eq. (12) is both analytically and numerically less cumbersome than (8), thus providing a computationally attractive alternative to Eq. (9). Comparing p_1 and p_3 obtained by means of Eq. (12), it holds

$$p_3(\mathbf{x}; t) = -\frac{1}{2\pi r} \frac{\partial}{\partial r} p_1(r; t), \quad r = |\mathbf{x}|, \quad (13)$$

195 which provides a well-known relation between the one-dimensional and three-dimensional PDFs p_1 and p_3 [20].

It is noted that the previously mentioned constraint on parameter β ($0 < \beta \leq 1/2$) guarantees to have a proper diffusion process characterized by an unimodal PDF [37, 39]. In fact, when $1/2 < \beta < 1$ function $\mathcal{M}_\beta(z)$ is bimodal and no diffusion process is modelled by p_3 , because the latter assumes also negative values, see (13). The subdiffusive character of diffusion is highlighted by the particle variance σ_d^2 which in one-, two- and three-dimensions ($d = 1, 2, 3$) is [39, 20]:

$$\sigma_1^2 = \int_{-\infty}^{+\infty} x^2 p_1(x; t) dx = \frac{2\nu t^{2\beta}}{\Gamma(2\beta + 1)}, \quad \sigma_2^2 = 2\sigma_1^2, \quad \sigma_3^2 = 3\sigma_1^2. \quad (14)$$

In Figure 1, the fundamental solutions of time-fractional diffusion equation (7) are presented for the one-, two- and three-dimensional case and for five different values of parameter β , i.e. $\beta = 0.1, 0.2, 0.3, 0.4$ and 0.5 , corresponding the latter to the case of ordinary diffusion and the other values to subdiffusive processes.

Fundamental solutions have been numerically calculated on the basis of the analytical representation given in Eq. (12). In particular the numerical evaluation of the $\mathcal{M}$ -Wright/Mainardi function $\mathcal{M}_\eta(z)$ and of the fundamental solutions in d dimensions $p_d(\mathbf{x}; t)$ have been performed by means of a python library (`pyMlib`) based on state-of-the-art algorithms for the Mittag-Leffler function [42] as well as on standard Laplace inversion and standard integration techniques available in the `Integrate` submodule of the `SciPy` scientific library [43].

In this respect, it is here reminded that the Laplace transform of the $\mathcal{M}$ -Wright/Mainardi function is [39, 16, 17]

$$\int_0^{+\infty} e^{-st} \mathcal{M}_\eta(t) dt = E_\eta(-s), \quad 0 < \eta < 1, \quad \text{Re}(s) > 0,$$

where $E_\eta(z)$ is the Mittag-Leffler function defined by [44]

$$E_\rho(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\rho n + 1)}, \quad \rho > 0, \quad z \in \mathbb{C}.$$

Figure 1: Fundamental solutions of the time-fractional diffusion equation (7) in the (a) one-dimensional, (b) two-dimensional and (c) three-dimensional cases for five values of the parameter β : $\beta = 0.1, 0.2, 0.3, 0.4$ and 0.5 .

210 4. Numerical results

In this Section the profiles of the effective indicator $\varphi_e(\mathbf{x}, t)$ defined in (3) are shown and discussed. In particular, to compute (3) the indicator function $\mathcal{I}_\Omega(\mathbf{x}, t)$ is obtained according to definition (1) by solving the level set equation (4) with constant velocity $\mathcal{V}$, and the PDF $p_d(\mathbf{x}; t|\bar{\mathbf{x}}) = p_d(\mathbf{x} - \bar{\mathbf{x}}; t)$ is the
215 fundamental solution of the d -dimensional time-fractional diffusion given in (12). The algorithm used to compute (12) is described in Section 3.

The evolution of random fronts propagating in subdiffusive and diffusive media has been numerically investigated by means of a software package developed for this goal. The code makes use of a general-purpose library (`liblsm90`)
220 which aims at providing a robust and efficient tool for studying the evolution of co-dimension one fronts propagating in one-, two- and three-dimensional system. The library, written in Fortran2008/OpenMP, is presently still under active development and is intended to be released as open source software under the GNU/GPLv3 licence. Along with standard algorithms useful for the calculation of the front evolution by means of the classical LSM, the library includes
225 Fast Marching Method (FMM) algorithms, which allow to efficiently solve the eikonal equation (5) arising from the classical LSM formulation in place of the standard Hamilton–Jacobi equation whenever it is possible, as in the case here under consideration.

230 All the results presented in the following have been obtained making use of the above-mentioned software and have been post-processed for exposition purposes by means of open source software such as `SciPy` [43] and `Matplotlib` [45] in the IPython framework [46].

All the calculations have been performed by means of the computational
235 server available at the Basque Center for Applied Mathematics (BCAM) in Bilbao, Basque Country – Spain.

A selection of the results concerning the propagation of random fronts in media characterized by anomalous diffusion is presented in this Section. The

discussion is restricted to fronts propagating in two-dimensional space, the one-
240 dimensional case being discussed elsewhere [8] and a thorough analysis including
three-dimensional cases being still under development.

Once the main features of one-dimensional propagation of plane fronts have
been investigated [8], the two-dimensional case has the relevant advantage of
allowing to investigate practical situations of effective interest in which multi-
245 dimensional effects possibly come into play, still avoiding the complexity of a
three-dimensional analysis.

The analysis proposed in this section concerns the investigation of the prop-
agation of random fronts in two-dimensional subdiffusive and diffusive media,
focusing on the effects of the diffusion coefficient ν , the parameter β and the
250 level set velocity $\mathcal{V}$.

First the case of an initial front profile with circular shape is discussed, to
focus on the effects of ν , β and $\mathcal{V}$ avoiding the two-dimensional effects connected
to a non-axisymmetric initial front profile. These effects are then discussed in the
case of a square-shaped initial front profile, with the main aim of investigating
255 the effects of the parameter β on the short-term and long-term evolution of the
random front.

In both cases, the computational domain is the rectangular region $[-2\text{ m}, 2\text{ m}] \times$
 $[-2\text{ m}, 2\text{ m}]$. The circular profile has a radius $r = 0.4\text{ m}$, and the square-shaped
profile has a side ℓ such that in both cases the region enclosed by the front when
260 $t = 0\text{ s}$ has the same area (i.e. $\ell \approx 0.7\text{ m}$). The center of both regions is located
in $(x, y) = (0, 0)$.

In the scenario involving a circular initial front profile, when the diffusion
coefficient ν is *small* enough, it is seen that for any value under investigation
of the parameter β the only outstanding feature of the propagating front is its
265 smoothing with respect to the sharp front that characterizes the deterministic
case, i.e. a non-diffusive system. Profiles representative of this behaviour are
shown in Fig. 2, where the numerical computations of $\varphi_e(\mathbf{x}, t)$ as defined in
(3) are plotted together with the deterministic front (dashed line) at different

time instants for the case $\nu = 10^{-3} \text{m}^2 \text{s}^{-2\beta}$ and two different values of the
 270 level set velocity ($\mathcal{V} = 0.1 \text{ms}^{-1}, 1 \text{ms}^{-1}$). Moreover, from Figure 2(b,d,f) it is
 evident that when $\mathcal{V}$ increases, the value of the parameter β becomes less and
 less relevant for the propagation of the front profile, thus allowing to conclude
 that the knowledge of the exact value of β becomes less and less critical. On
 contrast, as shown in Figure 2(a,c,e), for small values of $\mathcal{V}$ the front profile is
 275 clearly affected, at least qualitatively, by the value of the parameter β at any
 stage of its evolution, the profile being sharper and sharper as the parameter β
 becomes smaller and smaller.

In Fig. 3 it is shown that the maximum value of the effective indicator φ_e is
 initially slightly affected by the diffusion processes and later grows towards the
 280 initial maximum value.

From this analysis, it emerges that for *sufficiently small* values of the diffu-
 sion coefficient ν , the random front propagation differs from the deterministic
 front propagation mainly for the shape of the front profiles as illustrated in
 Fig. 2.

285 When the diffusion coefficient ν is *large*, the qualitative behaviour of the
 random fronts is remarkably different from the behaviour reported in Fig. 2
 and Fig. 3. In fact, as it is shown in Fig. 4, in this case the smoothing of the
 front may reach the point as to affect the existence of the bulk of the region Ω_e
 enclosed by the propagating front, where the “bulk region” is intended as the
 290 region characterized by φ_e close to 1. In this scenario, the maximum value of the
 effective indicator φ_e can be substantially reduced from its maximum unitary
 value, especially when the region Ω_e enclosed by the propagating front is small
 enough. This effect may be referred to as a *weakening* effect of the diffusion on
 the size of region Ω_e . The comparison of the results obtained for two different
 295 values of $\mathcal{V}$ ($\mathcal{V} = 0.1 \text{ms}^{-1}$ and $\mathcal{V} = 1 \text{ms}^{-1}$), respectively shown in Fig. 4(a,c,e)
 and Fig. 4(b,d,f), clearly put into evidence how the weakening effect described
 above is less relevant for the case with larger $\mathcal{V}$.

This behaviour suggests a balance between the diffusion coefficient ν and the

Figure 2: Cross-section at $y = 0$ of the φ_e fields corresponding to five values of the parameter β ($\beta = 0.1, 0.2, 0.3, 0.4, 0.5$) on the two-dimensional domain $[-2\text{ m}, 2\text{ m}] \times [-2\text{ m}, 2\text{ m}]$ at three values of the time t (top, middle and bottom row) and for two values of velocity $\mathcal{V}$ (left and right column). The initial profile at $t = 0\text{ s}$ is circular, with centre $C = (0, 0)$ and radius $r = 0.4\text{ m}$. The diffusion coefficient is $\nu = 0.001\text{ m}^2\text{s}^{-2\beta}$.

Figure 3: Time evolution of the maximum value of φ_e (φ_e^{max}) corresponding to five values of the parameter β ($\beta = 0.1, 0.2, 0.3, 0.4, 0.5$) for the case discussed in Figure 2.

velocity $\mathcal{V}$ from which the timescale τ_w of this weakening effect can be estimated.

In particular, such timescale emerges to be defined as

$$\tau_w = \frac{\sqrt{\nu \tau_u^{2\beta}}}{\mathcal{V}},$$

where τ_u is the unit of measurement for the temporal variable t and then it can be stated equal to 1. The estimation of the timescale τ_w for the weakening effect can be verified from the plots. For the cases showed in Fig. 3(a,b) it holds $\tau_w \simeq 0.3$ s and 0.03 s, respectively. So in Fig. 5(a,b) it holds $\tau_w \simeq 1$ s and 0.1 s, and in Fig. 7(a,b) it holds $\tau_w \simeq 3$ s and 0.3 s. Hereinafter, elapsed times are referred to *small* and *large* with respect to the timescale τ_w .

For *large* times, as the region enclosed by the propagating front increases in size, the qualitative behaviour previously discussed for the case of a *small* diffusion parameter ν can be eventually recovered, as it is seen in Fig. 4(f) for the case with $\mathcal{V} = 1 \text{ ms}^{-1}$. It is worth noting that for the case with $\mathcal{V} = 0.1 \text{ ms}^{-1}$, the same behaviour takes place for time larger than that reported in Fig. 4(e), as can be argued from Fig. 5.

Figure 5 also puts into evidence how the *weakening* effect due to the diffusion process is largely dependent on the parameter β , as expected, as well as on the velocity $\mathcal{V}$. In fact, the comparison of Fig. 5(a) and Fig. 5(b) shows how subdiffusive processes with *small* values of β may be responsible for smaller weakening effect with respect to subdiffusive processes with *larger* values of β (and also with respect to normal diffusive processes) when *small* values of the velocity $\mathcal{V}$ are involved. Interestingly, the situation can be reversed when larger values of $\mathcal{V}$ are involved.

The features of the random fronts described so far for the case with $\nu = 0.01 \text{ m}^2 \text{ s}^{-2\beta}$ remains qualitatively unchanged as the diffusion coefficient ν increases, but an important remark is in order. Depending on the physical interpretation of the indicator φ_e , it may – or may not – be meaningful to introduce the concept of “quencing” of the propagating front. In fact, assuming that the front is located, say, at the position for which the effective indicator φ_e reaches

Figure 4: The same as in Figure 2 but with $\nu = 0.01 \text{ m}^2 \text{ s}^{-2\beta}$.

Figure 5: Time evolution of the maximum value of φ_e (φ_e^{max}) corresponding to five values of the parameter β ($\beta = 0.1, 0.2, 0.3, 0.4, 0.5$) for the case discussed in Figure 4.

the threshold value φ_e^{th} , it might be meaningful to argue that when the effective
 325 indicator φ_e is reduced to the extent as to be less of φ_e^{th} over all the domain Ω_e ,
 the region Ω_e extinguishes and no further front propagation takes place. Such a
 circumstance may, for instance, be the case met in turbulent premixed combus-
 tion [10]: if φ_e marks the position of the propagating front, the value of φ_e may
 drop below the given threshold value and the flame would be regarded as extin-
 330 guished and no further calculation of the front evolution would be meaningful.
 Accordingly to what already noticed, such a circumstance would become more
 and more likely as the diffusion coefficient ν increases.

In Fig. 6, it is shown the propagation of the random and deterministic fronts
 obtained with $\nu = 0.1 \text{ m}^2 \text{ s}^{-2\beta}$. Assuming, for instance, $\varphi_e^{th} = 0.5$, area Ω_e asso-
 335 ciated with the front would be extinguished shortly after the diffusive process
 begins, independently from the value of the coefficient β (this is clearly seen
 also in Fig. 7), and any further calculation concerning the propagation of the
 connected random front would be meaningless, even if at larger times the effects
 of the diffusion would be such that the values of the effective indicator φ_e could
 340 reach values larger than the threshold φ_e^{th} , as it actually happens, as it is shown
 in Fig. 6(d,e,f) and Fig. 7. However, as already noticed, these considerations
 are problem-dependent and strongly connected with the physical process under
 consideration.

In order to investigate the two-dimensional effects connected to a non-
 345 axisymmetric initial front profile for various values of the parameter β , an ini-
 tial square-shaped profile has been considered. The numerical results shown
 in Fig. 8, Fig. 9 and Fig. 10 correspond to diffusive processes characterized by
 $\beta = 0.1$, $\beta = 0.3$ and $\beta = 0.5$, respectively. Figures point out that at *small*
 times when the value of β increases the effects of diffusion on the initial shape
 350 of the profile decreases, the subdiffusive process being in this case responsible
 for a stronger diffusive effects leading to a quicker smearing of the initial profile,
 because, when $t < \tau_w$, the smaller the parameter β the larger the particle vari-
 ance (14). Among the three cases shown in Fig. 8, Fig. 9 and Fig. 10, the case

Figure 6: The same as in Figure 2 but with $\nu = 0.1 \text{ m}^2 \text{ s}^{-2\beta}$.

Figure 7: Time evolution of the maximum value of φ_e (φ_e^{max}) corresponding to five values of the parameter β ($\beta = 0.1, 0.2, 0.3, 0.4, 0.5$) for the case discussed in Figure 6.

Figure 8: Distribution of the φ_e field for the case with $\beta = 0.1$ on the two-dimensional domain $[-2\text{ m}, 2\text{ m}] \times [-2\text{ m}, 2\text{ m}]$ at four instant times. The initial profile at $t = 0\text{ s}$ is a square, with center $C = (0, 0)$ and side $\ell = 0.7\text{ m}$. The diffusion coefficient is $\nu = 0.01\text{ m}^2\text{s}^{-2\beta}$ and $\mathcal{V} = 1\text{ ms}^{-1}$.

characterized by the ordinary diffusion process is clearly the one in which the
 355 initial front profile remain almost unchanged for a longer time, being the profile
 also sharper. For large times – as seen in Fig. 8(d), Fig. 9(d) and Fig. 10(d)
 – the diffusion process leads to similar front profiles in all the discussed cases,
 being the effective indicator field φ_e undistinguishable, as a matter of fact, in
 all the three cases.

360 5. Conclusions

In this paper, a multi-dimensional model to study the evolution of random
 fronts in media in which anomalous diffusion takes place has been presented.

Figure 9: The same as in Figure 8 but for the case with $\beta = 0.3$.

Figure 10: The same as in Figure 8 but for the case with $\beta = 0.5$.

In the proposed model, the position of the random front is obtained as the weighted mean of fronts calculated by means of the LSM, using as weight-function the relevant PDF.

Since anomalous diffusion can be modelled by the time-fractional diffusion equation, the corresponding fundamental solution is the required PDF.

In the one-dimensional case, it is well known that this fundamental solution can be conveniently written in terms of the $\mathcal{M}$ -Wright/Mainardi function, for the numerical evolution of which robust numerical methods are available. In the multi-dimensional case, the only available representation of this fundamental solution – to the best of the authors’ knowledge – was so far the one provided by Haniga [20], which cannot be straightforwardly used for the practical purpose of numerical calculations. It has been shown in this paper that, however, also in the multi-dimensional case the fundamental solution can be written in terms of the $\mathcal{M}$ -Wright/Mainardi function (see Eq. (9)).

Making use of this representation of the PDF for the practical purpose of numerical evaluation, the propagation of random fronts in two-dimensional media in which anomalous diffusion takes place has been numerically studied, focusing the attention on the case of random fronts propagating in two-dimensional subdiffusive media.

The selection of numerical results presented in this paper includes the case of a circular initial front profile and a case with a square-shaped initial front profile. In the first case, the investigation is performed as to point out the effects on the random front propagation of the parameter β (which is half of the order of the time fractional derivative), the diffusion coefficient ν and the deterministic front velocity $\mathcal{V}$. In the case of a square-shaped initial front profile, the primary intent of the investigation is to focus on the effects of the parameter β on the short-term and long-term evolution of the shape profile.

The presented numerical results clearly show that when the intensity of the diffusion (namely, the diffusion coefficient ν) is *small* enough, the propagating front is smoothed-out with respect to the sharp front obtained in the deter-

ministic, i.e. non-diffusive case, but the overall qualitative behaviour of the front propagation is not affected by the diffusive phenomena taking place in the medium. On contrast, when the intensity of the diffusion is *large*, the qualitative behaviour of the random fronts sensibly changes: in this case the smoothing of the front may be so significant as to even compromise the existence of the region enclosed by the propagating front (and the front itself, in turn). This effect, which has been referred to as a *weakening* effect of the diffusion phenomena on the region bounded by the propagating front can be, as shown, a transient effect gradually vanishing as the front propagates, or even a phenomenon which drastically affects the very existence of the front, depending on the intensity of the diffusion itself. In both cases, it has been shown how the velocity of propagation of the deterministic front plays a significant role in the features of the front propagation, as the above-discussed *weakening* effect is reduced as such velocity increases.

Moreover, aside from depending on the intensity of the diffusion phenomena through the diffusion coefficient ν , the above-mentioned *weakening* effects has also been shown to be more pronounced as the subdiffusive nature of the diffusion increases. The timescale τ_w of this weakening process has been estimated.

As far as the analysis of the propagation of an initially square-shaped front is concerned, it has been shown that, in the case under investigation and for *short* elapsed times (i.e. $t < \tau_w$), small values of the parameter β leads to a smearing of the initial front shape remarkably faster than in the case of ordinary diffusion. This occurs because, when $t < \tau_w$, the smaller the coefficient β the larger the particle variance (14).

The LSM library utilized for the purpose of this investigation, featuring optimized FMM algorithms as well, is intended to be released as open source software under the GNU General Public License version 3 (GPLv3).

420 Acknowledgements

This research is supported by GNFM/INdAM Young Researchers Project 2013, by Bizkaia Talent and European Commission through COFUND programme under Grant AYD-000-226, and also by the Basque Government through the BERC 2014-2017 program and by the Spanish Ministry of Economy and 425 Competitiveness MINECO: BCAM Severo Ochoa accreditation SEV-2013-0323. GP would like to thank Prof. F. Mainardi for continuous support, suggestions and encouragement.

References

- [1] E. Barkai, Y. Garini, R. Metzler, Strange kinetics of single molecules in 430 living cells, *Phys. Today* 65 (8) (2012) 29–35.
- [2] F. Höfling, T. Franosch, Anomalous transport in the crowded world of biological cells, *Rep. Prog. Phys.* 76 (2013) 046602.
- [3] R. Metzler, J.-H. Jeon, A. G. Cherstvy, E. Barkai, Anomalous diffusion models and their properties: non-stationarity, non-ergodicity, and ageing 435 at the centenary of single particle tracking, *Phys. Chem. Chem. Phys.* 16 (2014) 24128–24164.
- [4] D. del Castillo-Negrete, Non-diffusive, non-local transport in fluids and plasmas, *Nonlin. Processes Geophys.* 17 (2010) 795–807.
- [5] R. Metzler, J. Klafter, The restaurant at the end of the random walk: recent 440 developments in fractional dynamics descriptions of anomalous dynamical processes, *J. Phys. A: Math. Theor.* 37 (31) (2004) R161–R208.
- [6] A. Blumen, A. A. Gurtovenko, S. Jespersen, Anomalous diffusion and relaxation in macromolecular systems, *J. Non-Cryst. Solids* 305 (2002) 71–80.
- [7] G. Pagnini, Short note on the emergence of fractional kinetics, *Physica A* 445 409 (2014) 29–34.

- [8] A. Mentrelli, G. Pagnini, Random front propagation in fractional diffusive systems, submitted.
- [9] J. A. Sethian, P. Smereka, Level set methods for fluid interfaces, *Ann. Rev. Fluid Mech.* 35 (2003) 341–372.
- 450 [10] G. Pagnini, E. Bonomi, Lagrangian formulation of turbulent premixed combustion, *Phys. Rev. Lett.* 107 (2011) 044503.
- [11] G. Pagnini, L. Massidda, Modelling turbulence effects in wildland fire propagation by the randomized level-set method, Tech. Rep 2012/PM12a, CRS4, revised Version August 2014. arXiv:1408.6129 (July 2012).
- 455 [12] G. Pagnini, L. Massidda, The randomized level-set method to model turbulence effects in wildland fire propagation, in: D. Spano, V. Bacciu, M. Salis, C. Sirca (Eds.), *Modelling Fire Behaviour and Risk. Proceedings of the International Conference on Fire Behaviour and Risk. ICFBR 2011*, Alghero, Italy, October 4–6 2011, 2012, pp. 126–131, ISBN 978-88-904409-7-7.
- 460 [13] G. Pagnini, A. Mentrelli, Modelling wildland fire propagation by tracking random fronts, *Nat. Hazards Earth Syst. Sci.* 14 (2014) 2249–2263.
- [14] G. Pagnini, A model of wildland fire propagation including random effects by turbulence and fire spotting, in: *Proceedings of XXIII Congreso de Ecuaciones Diferenciales y Aplicaciones XIII Congreso de Matemática Aplicada*. Castelló, Spain, 9–13 September 2013, 2013, pp. 395–403.
- 465 [15] G. Pagnini, Fire spotting effects in wildland fire propagation, in: F. Casas, V. Martínez (Eds.), *Advances in Differential Equations and Applications*, Vol. 4 of SEMA SIMAI Springer Series, Springer International Publishing Switzerland, 2014, pp. 203–216, iISBN: 978-3-319-06952-4. (eBook: 978-3-319-06953-1).
- 470 [16] F. Mainardi, A. Mura, G. Pagnini, The M-Wright function in time-fractional diffusion processes: A tutorial survey, *Int. J. Diff. Equations* 2010 (2010) 104505.

- [17] F. Mainardi, Fractional Calculus and Waves in Linear Viscoelasticity, Imperial College Press, London, 2010.
- [18] G. Pagnini, The M-Wright function as a generalization of the Gaussian density for fractional diffusion processes, *Fract. Calc. Appl. Anal.* 16 (2) (2013) 436–453.
- [19] I. Podlubny, Fractional Differential Equations, Academic Press, San Diego, 1999.
- [20] A. Hanyga, Multidimensional solutions of time-fractional diffusion-wave equations, *Proc. R. Soc. Lond. A* 458 (2002) 933–957.
- [21] F. Mainardi, G. Pagnini, R. Gorenflo, Mellin transform and subordination laws in fractional diffusion processes, *Fract. Calc. Appl. Anal.* 6 (2003) 441–459.
- [22] S. J. Osher, J. A. Sethian, Fronts propagating with curvature dependent speed: algorithms based on Hamilton–Jacobi formulations, *J. Comput. Phys.* 79 (1988) 12–49.
- [23] N. Peters, Turbulent Combustion, Cambridge University Press, Cambridge, 2004.
- [24] V. Mallet, D. E. Keyes, F. E. Fendell, Modeling wildland fire propagation with level set methods, *Comput. Math. Appl.* 57 (2009) 1089–1101.
- [25] E. Jøttum, J. O. Helland, M. Prodanović, A level set method for simulating capillary-controlled displacements at the pore scale with nonzero contact angles, *Water Resour. Res.* 49 (2013) 4645–4661.
- [26] M. Machacek, G. Danuser, Morphodynamic profiling of protrusion phenotypes, *Biophys. J.* 90 (2006) 1439–1452.
- [27] W. Guo, H. H. Sawin, Review of profile and roughening simulation in microelectronics plasma etching, *J. Phys. D: Appl. Phys.* 42 (2009) 194014.

- 500 [28] J. J. Monaghan, Smoothed particle hydrodynamics, *Rep. Prog. Phys.* 68 (2005) 1703–1759.
- [29] V. L. Zimont, Gas premixed combustion at high turbulence. turbulent flame closure combustion model, *Exp. Therm. Fluid Sci.* 21 (2000) 179–186.
- [30] V. A. Sabelnikov, A. N. Lipatnikov, Towards an extension of TFC model of
505 premixed turbulent combustion, *Flow Turbul. Combust.* 90 (2013) 387–400.
- [31] T. Ohta, D. Jasnow, K. Kawasaki, Universal scaling in the motion of random interfaces, *Phys. Rev. Lett.* 49 (1982) 1223–1226.
- [32] H. M. Soner, N. Touzi, A stochastic representation for the level set equations, *Commun. Part. Diff. Eq.* 27 (2002) 2031–2053.
- 510 [33] O. Juan, R. Keriven, G. Postelnicu, Stochastic motion and the Level Set Method in Computer Vision: Stochastic active contours, *Int. J. Comput. Vision* 69 (2006) 7–25.
- [34] Y. L. Klimontovich, Nonlinear Brownian motion, *Physics-Uspekhi* 37 (1994) 737–767.
- 515 [35] M. S. Waterman, D. E. Whiteman, Estimation of probability densities by empirical density functions, *Int. J. Math. Educ. Sci. Technol.* 9 (1978) 127–137.
- [36] W. R. Schneider, W. Wyss, Fractional diffusion and wave equations, *J. Math. Phys.* 30 (1) (1989) 134–144.
- 520 [37] F. Mainardi, Fractional relaxation-oscillation and fractional diffusion-wave phenomena, *Chaos Solitons Fract.* 7 (1996) 1461–1477.
- [38] F. Mainardi, The fundamental solutions for the fractional diffusion-wave equation, *Appl. Math. Lett.* 9 (6) (1996) 23–28.
- [39] F. Mainardi, Y. Luchko, G. Pagnini, The fundamental solution of the space-time fractional diffusion equation, *Fract. Calc. Appl. Anal.* 4 (2) (2001)
525 153–192.

- [40] F. Mainardi, G. Pagnini, The Wright functions as solutions of the time-fractional diffusion equation, *Appl. Math. Comput.* 141 (1) (2003) 51–62.
- [41] F. Mainardi, G. Pagnini, R. Gorenflo, Some aspects of fractional diffusion equations of single and distributed order, *Appl. Math. Comput.* 187 (2007) 295–305.
- [42] R. Gorenflo, J. Loutchko, Y. Luchko, Computation of the Mittag–Leffler function $E_{\alpha,\beta}(z)$ and its derivative, *Fract. Calc. Appl. Anal.* 5 (2002) 491–518.
- [43] E. Jones, T. Oliphant, P. Peterson, et al., SciPy: Open source scientific tools for Python (2001–).
URL <http://www.scipy.org/>
- [44] F. Mainardi, R. Gorenflo, On Mittag–Leffler-type functions in fractional evolution processes, *J. Comput. Appl. Math.* 118 (2000) 283–299.
- [45] J. D. Hunter, Matplotlib: A 2D graphics environment, *Comput. Sci. Eng.* 9 (3) (2007) 90–95.
- [46] F. Pérez, B. E. Granger, IPython: a system for interactive scientific computing, *Comput. Sci. Eng.* 9 (3) (2007) 21–29. doi:10.1109/MCSE.2007.53.
URL <http://ipython.org>