Derivation of the homogenized von Kármán plate theory from 3d elasticity

Stefan Neukamm · Igor Velčić

the date of receipt and acceptance should be inserted later

Abstract In this paper we derive, by means of Γ -convergence, the homogenized von Kármán plate model starting from three dimensional nonlinear elasticity. We assume that the thickness of the plate is h , the oscillations of the material is $\varepsilon(h)$, where $\frac{h}{\varepsilon(h)} \rightarrow \gamma \in (0, +\infty)$. We also assume that the strain energy of the plate has the order h^4 , which corresponds to the von Kármán model in the case of the ordinary plate. The obtained model mixes the bending part of the energy with the stretching part.

Keywords homogenization · Gamma convergence · von Kármán plate · two-scale convergence

Mathematics Subject Classification (2010) 74K20 · 74B20

1 Introduction

The study of thin structures is the subject of numerous works in the theory of elasticity. Many authors have proposed two-dimensional shell and plate models and we come to the problem of their justification. There is a vast literature on the subject of plates and shells (see [5, 6]).

The justification of the model of plates and shells, by using Γ -convergence is well established. The first works in that direction are [12, 13]. The thickness of the plate is assumed to be h , a small parameter, and the external loads are assumed to be of the order 0. The obtained model for plate and shells differs from the one obtained by the formal asymptotic expansion in the sense that additional relaxation of the energy functional is done.

Max Planck Institute für Mathematik in Naturwissenschaften, Inselstrasse 22, 04103 Leipzig Germany

E-mail: neukamm@mis.mpg.de

BCAM –Basque Center for Applied Mathematics Address Bizkaia Technology Park, 500, 48160 Derio Basque Country – Spain

E-mail: ivelcic@bcamath.org

From the pioneering work of Friesecke, James, Müller [8] higher order models of plates and shells are justified from three dimensional nonlinear elasticity (see [8–11, 15, 16]). Here, higher order, relates that we assume that the magnitude of the external loads (i.e. of the strain energy) behaves like h^α , $\alpha > 0$ (i.e. h^β , $\beta > 0$). Depending on different parameter α different lower-dimensional models are obtained (see [9]).

In this paper we look for the model of the homogenized plate in the von Kármán regime, assuming that the strain energy is of the order h^4 , where h is the thickness of the plate. We assume that the material oscillates by the order $\varepsilon(h)$ and that $\frac{h}{\varepsilon(h)} \rightarrow \gamma \in (0, +\infty)$. The homogenized cell formula is given by the expression (4.29) and the main result is given in the Theorem 9.

The homogenization of 3D elasticity is done in [4, 17]. The homogenized membrane plate models are discussed in [2, 4]. In these cases in the cell formula, due to non-convexity, we have to include the possible oscillations on the larger scale than the oscillations of the material (see [17] for the example of relaxation on the larger scale corresponding to buckling). However, in small strain regimes, due to the convexity of the linearized stored energy density, this phenomena does not occur. It is the main reason why we can apply the tool of two-scale convergence, considering only the oscillations of the order $\varepsilon(h)$, where $\varepsilon(h)$ is the order of the oscillations of the material. Recently, such analysis is performed to obtain bending-torsion model of rod (see [18, 19]). Partial result are obtained in the bending case of plate (see [19]) where some geometrical difficulties appear, due to the fact that the limiting deformations are exact isometries. These difficulties does not appear in the von Kármán case, since the limiting deformations are only linearized isometries, which have more flexible geometry. A model of periodically wrinkled plate of the von Kármán type is obtained in [23]. In this model the wrinkles are assumed to be of a period h and the thickness of the period h^2 and then the dimensional reduction is done from 3D elasticity. In the cell formula we also can notice the mixing of bending and stretching part of energy as it can be seen here.

Throughout the paper $\bar{A}$ or $\{A\}^-$ denotes the closure of the set. By a domain we call a bounded open set with Lipschitz boundary. $\mathbf{I}$ denotes the identity matrix, by $\text{SO}(3)$ we denote the rotations in $\mathbb{R}^3$. By $\mathbb{R}_{\text{sym}}^{n \times n}$ we denote the set of symmetric matrices of the dimension $n \times n$ and by $\mathbb{R}_{\text{skew}}^{n \times n}$ the set of antisymmetric matrices. $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ are the vectors of the canonical base in $\mathbb{R}^3$. $\rightarrow$ denotes the strong convergence and $\rightharpoonup$ the weak convergence. By $\mathbf{A} \cdot \mathbf{B}$ we denote $\text{tr}(\mathbf{A}^T \mathbf{B})$. We suppose that the Greek indices α, β take the values in the set $\{1, 2\}$ while the Latin indices i, j take the values in the set $\{1, 2, 3\}$.

2 Setting up the problem

Let $\Omega^h = \omega \times [-\frac{h}{2}, \frac{h}{2}]$ where ω is a two-dimensional domain with Lipschitz boundary in the plane spanned by $\mathbf{e}_1, \mathbf{e}_2$. By Ω we denote Ω^1 . By $\hat{x}$ we denote $\hat{x} = (x_1, x_2)$. The canonical cell in $\mathbb{R}^2$ we denote by $Y = [0, 1]^2$; the generic

point in Y is $y = (y_1, y_2)$. Let $W : \Omega \times \mathbb{R}^2 \times \mathbb{R}^{3 \times 3} \rightarrow [0, \infty]$ satisfies the following hypothesis:

- i) $W(x, \cdot; \cdot)$ is continuous for a.e. $x \in \Omega$.
- ii) $W(\cdot, y; \mathbf{F})$ is measurable for all $(y; \mathbf{F}) \in \mathbb{R}^2 \times \mathbb{R}^{3 \times 3}$.
- iii) $W(x, \cdot; \mathbf{F})$ is Y -periodic for almost every $x \in \Omega$ and all $\mathbf{F} \in \mathbb{R}^{3 \times 3}$.
- iv) there exists $\delta > 0$ such that the function $\mathbf{F} \rightarrow W(x, y, \mathbf{F})$ is of class C^2 for a.e. $x \in \Omega$, all $y \in \mathbb{R}^2$ and all $\mathbf{F}$ such that $\text{dist}(\mathbf{F}, \text{SO}(3)) < \delta$. The second derivative $\partial^2 W / \partial \mathbf{F}^2 : \Omega \times \mathbb{R}^2 \times \{\text{dist}(\mathbf{F}, \text{SO}(3)) < \delta\} \rightarrow K_{3 \times 3}$ satisfies the properties i) and ii), where

$$K_{m \times n} := \{q | q \text{ quadratic form on } \mathbb{R}^{m \times n}\}$$

with the standard topology (we can identify $K_{m \times n}$ with the $(\mathbb{R}^{m \times n})^2$). We suppose that $\frac{\partial^2 W}{\partial \mathbf{F}^2}(x, \cdot; \cdot)$ is continuous for a.e. $x \in \Omega$ on the set $\mathbb{R}^2 \times \{\text{dist}(\mathbf{F}, \text{SO}(3)) < \delta\}$ and there exists a constant $\gamma > 0$ such that for a.e. $x \in \Omega$ and all $y \in Y$

$$\left| \frac{\partial^2 W}{\partial \mathbf{F}^2}(x, y; \mathbf{F})[\mathbf{G}, \mathbf{G}] \right| \leq \gamma \|\mathbf{G}\|^2 \text{ for } \text{dist}(\mathbf{F}, \text{SO}(3)) \leq \delta \text{ and } \mathbf{G} \in \mathbb{R}^{3 \times 3}$$

- v) W is frame-indifferent, i.e., $W(x, y; \mathbf{F}) = W(x, y; \mathbf{R}\mathbf{F})$ for a.e. $x \in \Omega$ and every $y \in \mathbb{R}^2$, $\mathbf{F} \in \mathbb{R}^{3 \times 3}$, $\mathbf{R} \in \text{SO}(3)$.
- vi) $W(x, y; \mathbf{F}) = 0$ if $\mathbf{F} \in \text{SO}(3)$ for a.e. $x \in \Omega$ and every $y \in \mathbb{R}^2$. $W(x, y, \mathbf{F}) \geq C \text{dist}^2(\mathbf{F}, \text{SO}(3))$ for a.e. $x \in \Omega$ and every $y \in Y$, $\mathbf{F} \in \mathbb{R}^{3 \times 3}$ where the constant C is independent of $x, y, \mathbf{F}$.

The starting point of our analysis is the minimization problem for the plate with periodically changing heterogeneity. Let $\varepsilon : (0, \infty) \rightarrow (0, \infty)$ satisfies

$$\lim_{h \rightarrow 0} \varepsilon(h) = 0, \quad \text{and} \quad \lim_{h \rightarrow 0} \frac{h}{\varepsilon(h)} = \gamma \quad \text{with} \quad \gamma \in (0, \infty) \quad (2.1)$$

The cases $\gamma = 0$, $\gamma = \infty$ can also be considered, but this one is the most interesting one (see also [18]).

The strain energy of the plate, written on the canonical domain Ω and divided by the order of volume h , is given by

$$I^h(\mathbf{y}) = \int_{\Omega} W(x, \frac{\hat{x}}{\varepsilon(h)}, \nabla_h \mathbf{y}(x)) dx,$$

where $W : \Omega \times Y \times \mathbb{R}^{3 \times 3} \rightarrow [0, \infty]$ satisfies the properties i)-vi) and

$$\nabla_h \mathbf{y} = \left(\partial_1 \mathbf{y} \mid \partial_2 \mathbf{y} \mid \frac{1}{h} \partial_3 \mathbf{y} \right).$$

By $Q_3 : \Omega \times \mathbb{R}^2 \rightarrow K_{3 \times 3}$ we denote the function $Q_3(x, y) = D^2 W(x, y; \mathbf{I})$. From the properties i)-vi) we can conclude

- I) $Q_3(x, \cdot)$ is continuous for a.e. $x \in \Omega$.

- II) $Q_3(\cdot, y)$ is measurable for all $y \in \mathbb{R}^2$.
- III) $Q_3(x, \cdot)$ is Y -periodic for almost every $x \in \Omega$.
- IV) there exists $\gamma > 0$ such that for a.e. $x \in \Omega$ and all $y \in \mathbb{R}^2$ we have

$$Q_3(x, y)[\mathbf{G}] \leq \gamma \|\mathbf{G}\|^2, \quad \forall \mathbf{G} \in \mathbb{R}^{3 \times 3}$$

- V) Q_3 is positive semi-definite, zero on antisymmetric matrices and uniformly positive definite on symmetric matrices i.e. there exists $C > 0$ such that for a.e. $x \in \Omega$ and all $y \in \mathbb{R}^2$ we have

$$Q_3(x, y)[\mathbf{G}] = Q_3(x, y)[\text{sym } \mathbf{G}], \quad Q_3(x, y)[\mathbf{G}] \geq C \|\text{sym } \mathbf{G}\|^2, \quad \forall \mathbf{G} \in \mathbb{R}^{3 \times 3}$$

We shall look for the Γ -limit of the functionals

$$\frac{1}{h^4} I^h(y)$$

The reason why we divide by h^4 is because we are interested in the von Kármán type model of the periodically heterogeneous plate. In the third section we prove some technical results about two scale convergence which we need later, in the forth section we prove the Γ -convergence result (for Γ -convergence techniques see [3, 7]). To prove it we firstly need the compactness result which tells us how the displacements of the energy order h^4 look like and secondly we have to prove lower and upper bound which is standard in Γ -analysis.

3 Two-scale convergence

For the notion of two scale convergence see [1, 20]. Here $\omega \subset \mathbb{R}^n$ is a bounded Lipschitz domain and $Y = [0, 1]^n$. For $k = (k_i)_{i=1, \dots, n} \in \mathbb{Z}^n$ we denote by $|k| = (\sum_{i=1}^n k_i^2)^{1/2}$.

We denote by $C_{\#}^k(Y)$ the space of k -differentiable functions with continuous k -th derivative in $\mathbb{R}^n$ which are periodic of period Y . Then $L_{\#}^2(Y)$ (respectively $H_{\#}^m(Y)$) is the completion for the norm of $L^2(Y)$ (respectively $W^{1,m}(Y)$ of $C_{\#}^{\infty}(Y)$). Remark that $L_{\#}^2(Y)$ actually coincides with the space of functions in $L^2(Y)$ extended by Y -periodicity to the whole of $\mathbb{R}^n$. $W_{\#}^{1,2}(Y)$ coincides with the space of functions in $W^{1,2}(Y)$ which are Y -periodic at the boundary in the sense of traces and $W_{\#}^{2,2}(Y)$ coincides with the space of functions in $W^{2,2}(Y)$ which are with their first derivatives Y -periodic at the boundary in the sense of traces etc. Using the Fourier transform on torus it can be seen that (see e.g. [22])

$$W_{\#}^{1,m}(Y) = \{u, u(y) = \sum_{k \in \mathbb{Z}^n} c^k e^{2\pi i k \cdot y}, \bar{c}^k = c^{-k}, \sum_{k \in \mathbb{Z}^n} |k|^{2m} |c^k|^2 < \infty\}, \quad (3.2)$$

and the norm $\|u\|_{W_{\#}^{1,m}(Y)}$ is equivalent to the norm $\{\sum_{k \in \mathbb{Z}^n} (1 + |k|^{2m}) |c^k|^2\}^{1/2}$. We also set

$$\dot{W}_{\#}^{1,m}(Y) := \{u \text{ of type (3.2), } c_0 = 0\} = \{u, u \in W_{\#}^{1,m}(Y), \int_Y u = 0\}.$$

The restricted norm $\|u\|_{\dot{W}_{\#}^{1,m}(Y)}$ is equivalent to the norm

$$\left\{ \sum_{k \in \mathbb{Z}^2} |k|^{2m} |c^k|^2 \right\}^{1/2}.$$

The space $L^2(\omega; C_{\#}(Y))$ denotes the space of measurable and square integrable functions in $x \in \omega$ with values in the Banach space of continuous functions, Y -periodic in y . In the analogous way one can define $L^2(\omega; W_{\#}^{1,m}(Y))$. Using the representation (3.2) it can be seen that

$$\begin{aligned} L^2(\omega; W_{\#}^{1,m}(Y)) = \{u, u(x, y) = \sum_{k \in \mathbb{Z}^n} c^k(x) e^{2\pi i k \cdot y}, c^k \in L^2(\omega; \mathbb{R}^n), \bar{c}^k = c^{-k}, \\ \sum_{k \in \mathbb{Z}^n} |k|^{2m} \|c^k\|_{L^2(\omega)}^2 < \infty\}, \end{aligned} \quad (3.3)$$

where we have by $\bar{c}^k$ denoted the conjugate of c^k . The norm $\|u\|_{L^2(\omega; H_{\#}^m(Y))}$ is equivalent to the norm

$$\left\{ \sum_{k \in \mathbb{Z}^2} (1 + |k|^{2m}) \|c^k\|_{L^2(\omega)}^2 \right\}^{1/2}.$$

The space $\mathcal{D}(\omega; C_{\#}^{\infty}(Y))$ denotes the space of infinitely differentiable functions which take values in $C_{\#}^{\infty}(Y)$ with compact support in ω . It is easily seen that this space is dense in $L^2(\omega; W_{\#}^{1,m}(Y))$. In fact it can be seen that the space of finite linear combinations

$$\begin{aligned} FL(\omega; C_{\#}^{\infty}(Y)) = \{u, \exists n \in \mathbf{N} \ u(x, y) = \sum_{k \in \mathbb{Z}^n, |k| \leq n} c^k(x) e^{2\pi i k \cdot y}, \\ c^k \in C_0^{\infty}(\omega; \mathbb{R}^2), \bar{c}^k = c^{-k}\} \end{aligned}$$

is dense in $L^2(\omega; W_{\#}^{1,m}(Y))$.

Definition 1 A sequence $(u^{\varepsilon})_{\varepsilon > 0}$ of functions in $L^2(\omega)$ converges two-scale to a function u^0 belonging to $L^2(\omega \times Y)$ if for every $\psi \in L^2(\omega; C_{\#}(Y))$,

$$\int_{\omega} u^{\varepsilon}(x) \psi(x, \frac{x}{\varepsilon}) \rightarrow \int_{\omega} \int_Y u^0(x, y) \psi(x, y) \quad \text{as } \varepsilon \rightarrow 0.$$

By $\rightharpoonup$ we denote the two-scale convergence. The following theorems are given in [1].

Theorem 1 *Let $f \in L^1(\omega; C_\#(Y))$. Then $f(x, \frac{x}{\varepsilon})$ is a measurable function on ω for which it is valid:*

$$\|f(x, \frac{x}{\varepsilon})\|_{L^1(\omega)} \leq \int_{\omega} \sup_{y \in Y} |f(x, y)| dy =: \|f\|_{L^1(\omega; C_\#(Y))}, \quad (3.4)$$

and

$$\lim_{\varepsilon \rightarrow 0} \int_{\omega} f(x, \frac{x}{\varepsilon}) dx = \int_{\omega} \int_Y f(x, y) dy dx. \quad (3.5)$$

Theorem 2 *From each bounded sequence $(u^\varepsilon)_{\varepsilon > 0}$ in $L^2(\omega)$ one can extract a subsequence, and there exists a limit $u^0(x, y) \in L^2(\omega \times Y)$ such that this subsequence two-scale converges to u^0 .*

The following theorem tells us about the form of oscillations of order ε of weakly convergent sequences in $W^{1,2}(\omega)$.

Theorem 3 *Let $(u^\varepsilon)_{\varepsilon > 0}$ be a bounded sequence in $W^{1,2}(\omega)$ which converges weakly to a limit $u \in W^{1,2}(\omega)$. Then u^ε two-scale converges to $u(x)$, and there exists a unique function $u^1(x, y) \in L^2(\omega; \dot{W}_\#^{1,2}(Y))$ such that, up to a subsequence, $(\nabla u^\varepsilon)_{\varepsilon > 0}$ two-scale converges to $\nabla_x u + \nabla_y u^1(x, y)$.*

Remark 1 In the definition of two-scale convergence we have taken the test functions to be in the space $L^2(\omega; C_\#(Y))$. When we are dealing with the sequence of the functions which are bounded in $L^2(\omega)$ it is enough to take the test function to be in the space $\mathcal{D}(\omega; C_\#^\infty(Y))$.

The following lemmas will be needed later.

Lemma 1 *Let $(u^\varepsilon)_{\varepsilon > 0}$ be a bounded sequence in $L^2(\omega)$ which two-scale converges to $u^0(x, y) \in L^2(\omega \times Y)$. Let $(v^\varepsilon)_{\varepsilon > 0}$ be a sequence bounded in $L^\infty(\omega)$ which converges in measure to $v^0 \in L^\infty(\omega)$. Then $v^\varepsilon u^\varepsilon \rightharpoonup v^0(x) u^0(x, y)$.*

Proof.

We know that $(v^\varepsilon u^\varepsilon)_{\varepsilon > 0}$ is bounded in $L^2(\omega)$ and that there exists a subsequence of $(v^\varepsilon)_{\varepsilon > 0}$ such that $v^\varepsilon \rightarrow v^0$ a.e. in ω . Let us take $\psi \in \mathcal{D}(\omega; C_\#(Y))$ and write,

$$\begin{aligned} \int_{\omega} v^\varepsilon(x) u^\varepsilon(x) \psi(x, \frac{x}{\varepsilon}) dx &= \int_{\omega} (v^\varepsilon(x) - v^0(x)) u^\varepsilon(x) \psi(x, \frac{x}{\varepsilon}) dx \\ &\quad + \int_{\omega} v^0(x) u^\varepsilon(x) \psi(x, \frac{x}{\varepsilon}) dx \end{aligned} \quad (3.6)$$

The second converges to

$$\int_{\omega} \int_Y v^0(x) u^0(x, y) \psi(x, y) dy dx,$$

by the definition of two-scale convergence. We have to prove that the first term in (3.6) converges to 0. By Egoroff's theorem for an arbitrary $\delta > 0$ there exists $E \subset \omega$ such that $\text{meas}(E) < \delta$ and $v^\varepsilon \rightarrow v$ uniformly on E^c . We write

$$\begin{aligned} \int_{\omega} (v^\varepsilon(x) - v^0(x)) u^\varepsilon(x) \psi(x, \frac{x}{\varepsilon}) dx &= \int_E (v^\varepsilon(x) - v^0(x)) u^\varepsilon(x) \psi(x, \frac{x}{\varepsilon}) dx \\ &\quad + \int_{E^c} (v^\varepsilon(x) - v^0(x)) u^\varepsilon(x) \psi(x, \frac{x}{\varepsilon}) dx. \end{aligned} \quad (3.7)$$

The second term in (3.7) converges to 0 and can be made arbitrary small. For the first term by the Cauchy inequality we have that there exists $C > 0$ such that

$$\left| \int_E (v^\varepsilon(x) - v^0(x)) u^\varepsilon(x) \psi(x, \frac{x}{\varepsilon}) dx \right| \leq C \sqrt{\delta} \sup_{\varepsilon > 0} \|u^\varepsilon\|_{L^2(\omega)}. \quad (3.8)$$

By the arbitrariness of δ we have the claim.

The following lemma is the special case of Proposition 1.3 in [25].

Lemma 2 *Let $\omega \subset \mathbb{R}^n$ be a domain, $Y = [0, 1]^n$. Let $f : \omega \times \mathbb{R}^n \times \mathbb{R}^m \rightarrow [0, \infty]$ satisfies the following hypothesis:*

- a) $f(x, \cdot; \cdot)$ is continuous for a.e. $x \in \omega$.
- b) $f(\cdot, y; \mathbf{F})$ is measurable for all $(y; \mathbf{F}) \in \mathbb{R}^n \times \mathbb{R}^{3 \times 3}$.
- c) $f(x, \cdot; \mathbf{F})$ is Y -periodic for almost every $x \in \omega$ and all $\mathbf{F} \in \mathbb{R}^{3 \times 3}$.
- d) the function $\mathbf{F} \rightarrow f(x, y, \mathbf{F})$ is convex for a.e. $x \in \omega$ and all $y \in \mathbb{R}^n$.

Let $(U^\varepsilon)_{\varepsilon > 0}$ be a sequence bounded in $(L^2(\omega))^m$ such that $U^\varepsilon \rightharpoonup U \in (L^2(\omega \times Y))^m$. Then we have that

$$\liminf_{\varepsilon \rightarrow 0} \int_{\omega} f(x, \frac{x}{\varepsilon}, U^\varepsilon(x)) dx \geq \int_{\omega} \int_Y f(x, y, U(x, y)) dy dx.$$

Remark 2 One could also define the two scale convergence with respect to the oscillations of the first m components. Namely, let $\omega \subset \mathbb{R}^n$, $Y = [0, 1]^m$ where $m \leq n$. A sequence $(u^\varepsilon)_{\varepsilon > 0}$ of functions in $L^2(\omega)$ converges two-scale to a function u^0 belonging to $L^2(\omega \times Y)$ if for every $\psi \in L^2(\omega; C_\#(Y))$,

$$\int_{\omega} u^\varepsilon(x) \psi(x, \frac{x_1}{\varepsilon}, \dots, \frac{x_m}{\varepsilon}) dx \rightarrow \int_{\omega} \int_Y u^0(x, y) \psi(x, y) \quad \text{as } \varepsilon \rightarrow 0.$$

It can be easily seen that Theorem 1, 2, 3 and Lemma 1, 2 are valid in this case (see also [21] for time dependent problems).

We can prove the following lemma.

Lemma 3 *Let $\omega \subset \mathbb{R}^2$ be a domain, $Y = [0, 1]^2$. Let $\mathbf{u} \in L^2(\omega \times Y; \mathbb{R}^2)$ be such that for each $\psi \in \mathcal{D}(\omega; C_{\#}^{\infty}(Y))$ we have that*

$$\int_{\omega} \int_Y \mathbf{u}(x, y) \operatorname{rot}_y \psi(x, y) dy dx = 0, \quad (3.9)$$

where $\operatorname{rot}_y = (\partial_{y_2} v, -\partial_{y_1} v)$.

Then there exists unique functions $\mathbf{w}^0 \in L^2(\omega; \mathbb{R}^2)$, $w \in L^2(\omega; \dot{W}_{\#}^{1,2}(Y))$ such that $\mathbf{u}(x, y) = \mathbf{w}^0(x) + \nabla_y w(x, y)$.

Proof.

We follow the proof of Lemma 2 in [23]. Let us, as before, identify the space $L^2(\omega; \dot{W}_{\#}^{1,2}(Y))$ with the sequences of functions

$$L^2(\omega; \dot{W}_{\#}^{1,2}(Y)) = \{(b^k)_{k \in \mathbb{Z}^n}, b^k \in L^2(\omega; \mathbb{R}^2), \bar{b}^k = b^{-k}, \\ \int_{\omega} \sum_{k \in \mathbb{Z}^2} |k|^2 |b^k(x)|^2 dx < \infty\},$$

with the norm

$$\|(b^k)_k\|^2 = \int_{\omega} \sum_{k \in \mathbb{Z}^n} |k|^2 |b^k(x)|^2 dx,$$

and the space $L^2(\omega \times Y; \mathbb{R}^2)$ with

$$L^2(\omega \times Y; \mathbb{R}^2) = \{(\mathbf{c}_{\alpha}^k)_{k \in \mathbb{Z}^2, \alpha=1,2}, \mathbf{c}_{\alpha}^k \in L^2(\omega; \mathbb{R}^2), \bar{\mathbf{c}}_{\alpha}^k = \mathbf{c}_{\alpha}^{-k}, \\ \int_{\omega} \sum_{k \in \mathbb{Z}^n} \sum_{\alpha=1,2} |\mathbf{c}_{\alpha}^k(x)|^2 dx < \infty\},$$

with the norm

$$\|(\mathbf{c}_{\alpha}^k)_k\|^2 = \int_{\omega} \sum_{k \in \mathbb{Z}^n} \sum_{\alpha=1,2} |\mathbf{c}_{\alpha}^k(x)|^2 dx$$

The operator rot_y operates in the following way

$$\operatorname{rot}_y(b_k)_k = (k_2 b^k, -k_1 b^k)_{k \in \mathbb{Z}^2}.$$

It is easily seen that rot_y is continuous and one to one. We shall prove that it is enough to demand the condition (3.9) for $\psi \in FL(\omega; C_{\#}^{\infty}(Y))$. Using the properties of the Fourier transform the condition (3.9) can be interpreted in the following way: For given $((\mathbf{u}_{\alpha}^k)_{\alpha=1,2})_{k \in \mathbb{Z}^2} \in L^2(\omega \times Y; \mathbb{R}^2)$ and every $(b^k)_{k \in \mathbb{Z}^2} \in FL(\omega; C_{\#}^{\infty}(Y))$ we have that

$$\int_{\omega} \sum_{k \in \mathbb{Z}^2} (k_2 b^k \mathbf{u}_1^k - k_1 b^k \mathbf{u}_2^k) dx = 0.$$

By fixing $k_0 \in \mathbb{Z}^2, k_0 \neq 0$, and taking $b^{k_0} \in C_0^\infty(\omega; \mathbb{R}^2)$, and defining $b^{-k_0} = \overline{b^{k_0}}, b^{k_0}, \forall k \neq k_0, -k_0$, we conclude that

$$\int_{\omega} \operatorname{Re}(k_2 b^k \mathbf{u}_1^k - k_1 b^k \mathbf{u}_2^k) dx = 0.$$

From this it can be easily seen that there exists $v^{k_0} \in L^2(\omega; \mathbb{R}^2)$ such that $\mathbf{u}_{\alpha}^{k_0} = (k_0)_{\alpha} v^{k_0}$, for $\alpha = 1, 2$. This is valid for an arbitrary $k_0 \in \mathbb{Z}^2$ and we can easily conclude from the fact $((\mathbf{u}_{\alpha}^k)_{\alpha=1,2})_{k \in \mathbb{Z}^2} \in L^2(\omega \times Y; \mathbb{R}^n)$ that $\int_{\omega} \sum_{k \in \mathbb{Z}^n} |k|^2 |v^k(x)|^2 dx < \infty$. Now we have the claim by taking $\mathbf{w}^0(x) = (\mathbf{u}_1^0, \mathbf{u}_2^0) = (\operatorname{Re} \mathbf{u}_1^0, \operatorname{Re} \mathbf{u}_2^0)$, $w(x, y) = (v^k(x))_k \equiv \sum_{k \in \mathbb{Z}^2} v^k(x) e^{2\pi i k \cdot y}$.

The following lemma is crucial for the liminf inequality in Γ -convergence.

Lemma 4 *Let $\varepsilon : (0, \infty) \rightarrow (0, \infty)$ satisfies (2.1). Let $\omega \subset \mathbb{R}^2$ be a domain and $Y = [0, 1]^2$. Let $(s^h)_{h>0} \subset W^{1,2}(\omega)$ be a sequence, $s^h \rightarrow s$ strongly in $W^{1,2}(\omega)$. Let $(z_1^h, z_2^h)_{h>0}$ be a sequence bounded in $(W^{1,2}(\omega))^2$ such that $(\frac{z_1^h + \partial_1 s^h}{h}, \frac{z_2^h + \partial_2 s^h}{h})$ is bounded in $(L^2(\omega))^2$ and we let us assume that $(\frac{z_1^h + \partial_1 s^h}{h}, \frac{z_2^h + \partial_2 s^h}{h})$ two scale converges to $(\mathbf{l}_1, \mathbf{l}_2)$ with respect to $\varepsilon(h)$ oscillations, where $\mathbf{l} \in (L^2(\omega \times Y))^2$. Let us also assume that $\partial_{\beta} z_{\alpha}^h$ two scale converges to $r_{\alpha\beta}^0 \in L^2(\omega \times Y)$ with respect to $\varepsilon(h)$ oscillations, for $\alpha, \beta = 1, 2$. Then we have that $s \in W^{2,2}(\omega)$ and there exists $\mathbf{r} \in (L^2(\omega; \dot{W}_{\#}^{1,2}(Y)))^2$ such that $r_{\alpha\beta}^0 = -\partial_{\alpha\beta} s + \partial_{y\beta} \mathbf{r}_{\alpha}$, for $\alpha, \beta = 1, 2$. Moreover there exists $\mathbf{m} \in (L^2(\omega))^2$ and $p \in L^2(\omega; \dot{W}_{\#}^{1,2}(Y))$ such that $\mathbf{l} = \frac{1}{\gamma} \mathbf{r} + \mathbf{m} + \nabla_y p$.*

Proof.

The first part of the lemma is the direct consequence of the conditions imposed in Lemma and Theorem 3. Let's now prove the second part. By using Lemma 3 it is enough to prove that for every $\psi \in \mathcal{D}(\omega; C_{\#}^\infty(Y))$ we have that

$$\int_{\omega} \int_Y (\mathbf{l}_1 - \frac{1}{\gamma} \mathbf{r}_1, \mathbf{l}_2 - \frac{1}{\gamma} \mathbf{r}_2) \cdot \operatorname{rot}_y \psi(x, y) dy dx = 0, \quad (3.10)$$

Let us calculate

$$\begin{aligned} & \int_{\omega} \int_Y (\mathbf{l}_1(x, y) \partial_{y_2} \psi(x, y) - \mathbf{l}_2(x, y) \partial_{y_1} \psi(x, y)) dy dx \\ &= \lim_{h \rightarrow 0} \int_{\omega} \left(\frac{z_1^h(x) + \partial_1 s^h(x)}{h} \partial_{y_2} \psi(x, \frac{x}{\varepsilon(h)}) - \frac{z_2^h + \partial_2 s^h(x)}{h} \partial_{y_1} \psi(x, \frac{x}{\varepsilon(h)}) \right) dx \\ &= \lim_{h \rightarrow 0} \frac{\varepsilon(h)}{h} \int_{\omega} (\partial_1 z_2^h(x) - \partial_2 z_1^h(x)) \psi(x, \frac{x}{\varepsilon(h)}) dx \\ &\quad - \lim_{h \rightarrow 0} \int_{\omega} (z_1^h(x) \partial_{x_2} \psi(x, \frac{x}{\varepsilon(h)}) - z_2^h(x) \partial_{x_1} \psi(x, \frac{x}{\varepsilon(h)})) dx \\ &\quad - \lim_{h \rightarrow 0} \int_{\omega} \frac{s^h(x)}{h} (\partial_{x_1 y_2} \psi(x, \frac{x}{\varepsilon(h)}) - \partial_{x_2 y_1} \psi(x, \frac{x}{\varepsilon(h)})) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{\gamma} \int_{\omega} \int_Y (r_{21}^0(x, y) - r_{12}^0(x, y)) \psi(x, y) dy dx \\
&- \int_{\omega} \int_Y (\partial_1 v(x) \partial_{x_2} \psi(x, y) - \partial_2 v(x) \partial_{x_1} \psi(x, y)) dy dx \\
&- \lim_{h \rightarrow 0} \frac{\varepsilon(h)}{h} \int_{\omega} (\partial_1 s^h(x) \partial_{x_2} \psi(x, \frac{x}{\varepsilon(h)}) - \partial_2 s^h(x) \partial_{x_1} \psi(x, \frac{x}{\varepsilon(h)})) \\
&= \frac{1}{\gamma} \int_{\omega} \int_Y (\partial_{y_1} \mathbf{r}_2(x, y) - \partial_{y_2} \mathbf{r}_1(x, y)) \psi(x, y) dy dx \\
&= \frac{1}{\gamma} \int_{\omega} \int_Y (\mathbf{r}_1(x, y) \partial_{y_2} \psi(x, y) - \mathbf{r}_2(x, y) \partial_{y_1} \psi(x, y)) dy dx
\end{aligned}$$

From this we have the claim.

The following theorem is the part of [18, Theorem 6.3.3]

Theorem 4 *Let $\Omega = \omega \times S \subset \mathbb{R}^n$ be a domain and let $m \leq n$, $\omega \subset \mathbb{R}^n$, $S \subset \mathbb{R}^{n-m}$ domains, $Y = [0, 1]^m$. Let us for $\mathbf{u} \in W^{1,2}(\Omega; \mathbb{R}^k)$ define*

$$\begin{aligned}
\nabla_{m,h} \mathbf{u} &:= (\nabla_{\hat{x}} \mathbf{u}, \frac{1}{h} \nabla_{\bar{x}} \mathbf{u}), \\
\nabla_{\hat{x}} \mathbf{u} &:= (\partial_1 \mathbf{u}, \dots, \partial_m \mathbf{u}), \\
\nabla_{\bar{x}} \mathbf{u} &:= (\partial_{m+1} \mathbf{u}, \dots, \partial_n \mathbf{u})
\end{aligned}$$

Let $\varepsilon : (0, \infty) \rightarrow (0, \infty)$ satisfy the condition (2.1). Let $(\mathbf{u}^h)_{h>0}$ be a weakly convergent sequence in $W^{1,2}(\Omega; \mathbb{R}^k)$ with limit $\mathbf{u}$ and suppose that

$$\limsup_{h \rightarrow 0} \int_{\Omega} \|\nabla_{m,h} \mathbf{u}^h\|^2 dx < \infty.$$

Then $\mathbf{u}$ is independent of $\bar{x}$ and can be identified with a map in $W^{1,2}(\omega; \mathbb{R}^k)$.

Also there exists a map

$$\mathbf{k} \in L^2(\omega; W_{\#Y}^{1,2}(S \times Y; \mathbb{R}^k))$$

and a subsequence (not relabeled) such that

$$\nabla_{m,h} \mathbf{u}^h \rightharpoonup (\nabla_{\hat{x}} \mathbf{u}(\hat{x})|0) + \tilde{\nabla}_{m,\gamma} \mathbf{k}(x, y).$$

weakly two-scale in $L^2(\omega \times Y; \mathbb{R}^k)$ with respect to $\varepsilon(h)$ -oscillations in the first m -components.

Here we have by $W_{\#Y}^{1,2}(S \times Y; \mathbb{R}^k)$ denoted the space $W^{1,2}(S \times Y; \mathbb{R}^k)$ of functions which are Y -periodic and by $\tilde{\nabla}_{m,\gamma} \mathbf{k}(x, y)$ we have denoted

$$\tilde{\nabla}_{m,\gamma} \mathbf{k} := (\nabla_y \mathbf{k}(x, y), \frac{1}{\gamma} \nabla_{\bar{x}} \mathbf{k}(x, y)).$$

The following theorem is given in [18, Theorem 6.3.8]

Theorem 5 *Let S and Y be as in Theorem 4. There exists a constant c_γ such that for*

$$\mathbf{u} \in \mathcal{V} := \left\{ \mathbf{u} \in W_{\#Y}^{1,2}(S \times Y; \mathbb{R}^n) \mid \int_{S \times Y} \mathbf{u} dy d\bar{x} = 0 \right\} \quad (3.11)$$

the estimate

$$\begin{aligned} \inf_{r \in \mathcal{R}(m)} \|\mathbf{u} + r\|_{W^{1,2}(S \times Y; \mathbb{R}^n)} &\leq c_\gamma \left\| \text{sym } \tilde{\nabla}_{m,\gamma} \mathbf{u} \right\|_{L^2(S \times Y; \mathbb{R}^n)} \\ &\leq c_\gamma^2 \|\mathbf{u}\|_{W^{1,2}(S \times Y; \mathbb{R}^n)}, \end{aligned} \quad (3.12)$$

is valid, where

$$\mathcal{R}(m) := \left\{ \bar{r} \in \mathcal{V} : \bar{r}(y, \bar{x}) = \begin{pmatrix} \mathbf{0}_m \\ \bar{\mathbf{A}}(x - \mathbf{c}_S) \end{pmatrix} \text{ with } \mathbf{A} \in \mathbb{R}_{\text{skew}}^{(n-m) \times (n-m)} \right\} \quad (3.13)$$

Above $\mathbf{0}_m$ denotes the zero vector in $\mathbb{R}^m$, $\mathbf{c}_S := \frac{1}{\mathcal{H}^{n-m}(S)} \int_S \bar{x} d\bar{x}$.

4 Γ -convergence

We shall need the following theorem which can be found in [8].

Theorem 6 (on geometric rigidity)

Let $U \subset \mathbb{R}^m$ be a bounded Lipschitz domain, $m \geq 2$. Then there exists a constant $C(U)$ with the following property: for every $\mathbf{v} \in W^{1,2}(U; \mathbb{R}^m)$ there is associated rotation $\mathbf{R} \in \text{SO}(m)$ such that

$$\|\nabla \mathbf{v} - \mathbf{R}\|_{L^2(U)} \leq C(U) \|\text{dist}(\nabla \mathbf{v}, \text{SO}(m))\|_{L^2(U)}. \quad (4.1)$$

The constant $C(U)$ can be chosen uniformly for a family of domains which are Bilipschitz equivalent with controlled Lipschitz constants. The constant $C(U)$ is invariant under dilatations.

In the same way as in [9, Theorem 10, Remark 11] we can prove the following theorem.

Theorem 7 *Let $\omega \subset \mathbb{R}^2$ be a domain, $\Omega = \omega \times [-\frac{1}{2}, \frac{1}{2}]$. Let $\mathbf{y}^h \in W^{1,2}(\Omega; \mathbb{R}^3)$ be such that*

$$\int_{\Omega} \text{dist}^2(\nabla_h \mathbf{y}^h, \text{SO}(3)) dx \leq Ch^4,$$

for some $C > 0$. Then there exists map $\mathbf{R}^h \in W^{1,2}(\omega, \text{SO}(3))$ such that

$$\|\nabla_h \mathbf{y}^h - \mathbf{R}^h\|_{L^2(\Omega)} \leq Ch^2, \quad (4.2)$$

$$\|\nabla \mathbf{R}^h\|_{L^2(\omega)} \leq Ch. \quad (4.3)$$

Moreover there exist a constant rotation $\bar{\mathbf{Q}}^h \in \text{SO}(3)$ such that

$$\|\nabla_h \mathbf{y}^h - \bar{\mathbf{Q}}^h\|_{L^2(\Omega)} \leq Ch, \quad (4.4)$$

$$\|\mathbf{R}^h - \bar{\mathbf{Q}}^h\|_{L^p(\omega)} \leq C_p h, \quad \forall p < \infty. \quad (4.5)$$

Here all constants depend only on ω (and on p where indicated).

To prove Γ -convergence result we need to prove the compactness result, the lower and the upper bound.

4.1 Compactness result

The following lemma is the same as Lemma 13 in [9].

Lemma 5 *Let $\mathbf{y}^h \in W^{1,2}(\Omega; \mathbb{R}^3)$ be such that*

$$\int_{\Omega} \text{dist}^2(\nabla_h \mathbf{y}^h, \text{SO}(3)) dx \leq Ch^4, \quad (4.6)$$

Then there exists maps $\mathbf{R}^h \in W^{1,2}(\omega, \text{SO}(3))$ and constants $\bar{\mathbf{R}}^h \in \text{SO}(3)$, $\mathbf{c}^h \in \mathbb{R}^3$ such that

$$\tilde{\mathbf{y}}^h := (\bar{\mathbf{R}}^h)^T \mathbf{y}^h - \mathbf{c}^h$$

and the corrected in-plane and the out-of-plane displacements

$$\begin{aligned} \mathbf{u}^h(x_1, x_2) &:= \frac{1}{h^2} \left(\int_{-1/2}^{1/2} \begin{pmatrix} \tilde{\mathbf{y}}_1^h \\ \tilde{\mathbf{y}}_2^h \end{pmatrix} (x_1, x_2, x_3) dx_3 - \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \right) \\ v^h(x_1, x_2) &:= \frac{1}{h} \int_{-1/2}^{1/2} \tilde{\mathbf{y}}_3^h(x_1, x_2, x_3) dx_3 \end{aligned} \quad (4.7)$$

satisfy

$$\|\nabla_h \tilde{\mathbf{y}}^h - \mathbf{R}^h\|_{L^2(\Omega)} \leq Ch^2, \quad (4.8)$$

$$\|\mathbf{R}^h - \mathbf{I}\|_{L^p(\omega)} \leq C_p h \quad \forall p < \infty, \quad \|\nabla \mathbf{R}^h\|_{L^2(\omega)} \leq Ch. \quad (4.9)$$

Moreover every subsequence (not relabeled) has its subsequence (also not relabeled) such that

$$v^h \rightarrow v \quad \text{in } W^{1,2}(\omega), \quad v \in W^{2,2}(\omega) \quad (4.10)$$

$$\mathbf{u}^h \rightharpoonup \mathbf{u} \quad \text{in } W^{1,2}(\omega; \mathbb{R}^2), \quad (4.11)$$

$$\frac{\mathbf{R}^h - \mathbf{I}}{h^2} \rightarrow \mathbf{A} \quad \text{in } W^{1,2}(\omega; \mathbb{R}^{3 \times 3}), \quad (4.12)$$

$$\frac{\nabla_h \tilde{\mathbf{y}}^h - \mathbf{I}}{h} \rightarrow \mathbf{A} \quad \text{in } L^2(\Omega; \mathbb{R}^{3 \times 3}) \quad (4.13)$$

$$\partial_3 \mathbf{A} = 0, \quad \mathbf{A} \in W^{1,2}(\omega; \mathbb{R}^{3 \times 3}), \quad (4.14)$$

$$\mathbf{A} = \mathbf{e}_3 \otimes \nabla v - \nabla v \otimes \mathbf{e}_3, \quad (4.15)$$

$$\frac{\text{sym}(\mathbf{R}^h - \mathbf{I})}{h^2} \rightarrow \frac{\mathbf{A}^2}{2} \quad \text{in } L^2(\omega; \mathbb{R}^{3 \times 3}). \quad (4.16)$$

4.2 Lower bound

In the next lemma we shall characterize the symmetric part of the limiting strain and express it in terms of $\mathbf{u}, v$ and some auxiliary functions by which the problem will be relaxed.

Since the strain is an element of $L^2(\Omega)$ and we want to obtain its two-scale limit which characterizes only oscillations of the first two variables of the order $\varepsilon(h)$ (see Remark 2). By $\hat{x}$ we denote $\hat{x} = (x_1, x_2)$, by y we denote $y = (y_1, y_2)$.

Lemma 6 *Consider $\mathbf{y}^h : \Omega \rightarrow \mathbb{R}^3$, $\mathbf{R}^h \in W^{1,2}(\omega; \text{SO}(3))$ and define $\mathbf{u}^h, v^h$ by (4.7) Suppose that we have a subsequence of $\mathbf{y}^h$ such that (4.8)-(4.16) are valid. Let us denote*

$$\mathbf{G}^h := \frac{(\mathbf{R}^h)^T \nabla_h \mathbf{y}^h - \mathbf{I}}{h^2}, \quad (4.17)$$

Then there exists a subsequence of $\mathbf{G}^h$ (not relabeled) such that

$$\mathbf{G}^h \rightharpoonup \mathbf{G} \quad \text{in } L^2(\Omega \times Y; \mathbb{R}^{3 \times 3}),$$

and a map $\mathbf{w} \in L^2(\omega; W_{\#Y}^{1,2}([-\frac{1}{2}, \frac{1}{2}] \times Y; \mathbb{R}^3))$ such that

$$\text{sym } \mathbf{G}(x, y) = \text{sym} \left(\begin{array}{c|c} \mathbf{G}''(x, y) & \begin{array}{c} \frac{1}{\gamma} \partial_{x_3} \mathbf{w}_1(x, y) \\ \frac{1}{\gamma} \partial_{x_3} \mathbf{w}_2(x, y) \end{array} \\ \hline \begin{array}{c} \partial_{y_1} \mathbf{w}_3(x, y) \quad \partial_{y_2} \mathbf{w}_3(x, y) \end{array} & \frac{1}{\gamma} \partial_{x_3} \mathbf{w}_3(x, y) \end{array} \right),$$

where $\mathbf{G}''$ is a 2×2 submatrix given by

$$\mathbf{G}''(x, y) = \nabla_{\hat{x}} \mathbf{u}(\hat{x}) + \frac{1}{2} \nabla_{\hat{x}} v(\hat{x}) \otimes \nabla_{\hat{x}} v(\hat{x}) - x_3 \nabla_{\hat{x}}^2 v(\hat{x}) + \nabla_y \begin{pmatrix} \mathbf{w}_1(x, y) \\ \mathbf{w}_2(x, y) \end{pmatrix}. \quad (4.18)$$

Proof.

We shall use the help of the proof of Lemma 15 in [9]. By the assumption $\mathbf{G}^h$ is bounded in L^2 , thus a subsequence converges two scale.

Let us write

$$\mathbf{R}^h \mathbf{G}^h = \frac{\nabla_h \mathbf{y}^h - \mathbf{I}}{h^2} - \frac{\mathbf{R}^h - \mathbf{I}}{h^2} \quad (4.19)$$

Since $\mathbf{R}^h \rightarrow \mathbf{I}$ boundedly in measure we have from Lemma 1

$$\mathbf{R}^h \mathbf{G}^h \rightharpoonup \mathbf{G} \quad (4.20)$$

Let us by $\mathbf{z}^h$ denote the displacement

$$\mathbf{z}^h = \begin{pmatrix} x_1 \\ x_2 \\ hx_3 \end{pmatrix} + \begin{pmatrix} h^2 \mathbf{u}_1^h \\ h^2 \mathbf{u}_2^h \\ hv^h \end{pmatrix} + hx_3 \begin{pmatrix} \mathbf{R}_{13}^h \\ \mathbf{R}_{23}^h \\ 0 \end{pmatrix} \quad (4.21)$$

First we want to prove that $\frac{\mathbf{y}^h - \mathbf{z}^h}{h^2}$ is a bounded sequence in $W^{1,2}(\Omega; \mathbb{R}^3)$ such that $\nabla_h \frac{\mathbf{y}^h - \mathbf{z}^h}{h^2}$ is bounded in $L^2(\Omega; \mathbb{R}^{3 \times 3})$. Let us firstly check the boundedness of the derivatives of $\nabla_h \frac{\mathbf{y}^h - \mathbf{z}^h}{h^2}$ in $L^2(\Omega; \mathbb{R}^{3 \times 3})$. Observe for $\alpha = 1, 2, i = 1, 2, 3$:

$$\begin{aligned} \left\| \frac{\partial_\alpha (\mathbf{y}^h - \mathbf{z}^h)_i}{h^2} \right\|_{L^2} &= \left\| \frac{\partial_\alpha \mathbf{y}_i^h - \int_{-1/2}^{1/2} \partial_\alpha \mathbf{y}_i^h}{h^2} \right\|_{L^2} \\ &\leq \left\| \frac{\partial_\alpha \mathbf{y}_i^h - \mathbf{R}_{i\alpha}^h}{h^2} \right\|_{L^2} \\ &\quad + \left\| \int_{-1/2}^{1/2} \frac{(\partial_\alpha \mathbf{y}_i^h - \mathbf{R}_{i\alpha}^h)}{h^2} \right\|_{L^2} \end{aligned} \quad (4.22)$$

The both terms in (4.22) are bounded due to the inequality (4.8). For $\alpha = 1, 2$ we also have

$$\left\| \frac{\frac{1}{h} \partial_3 (\mathbf{y}^h - \mathbf{z}^h)_\alpha}{h^2} \right\|_{L^2} = \left\| \frac{\frac{1}{h} \partial_3 \mathbf{y}_\alpha^h - \mathbf{R}_{\alpha 3}^h}{h^2} \right\|_{L^2}. \quad (4.23)$$

The last term is bounded also as a consequence of (4.8). And finally

$$\begin{aligned} \left\| \frac{\frac{1}{h} \partial_3 (\mathbf{y}^h - \mathbf{z}^h)_3}{h^2} \right\|_{L^2} &= \left\| \frac{\frac{1}{h} \partial_3 \mathbf{y}_3^h - 1}{h^2} \right\|_{L^2} \\ &\leq \left\| \frac{\frac{1}{h} \partial_3 \mathbf{y}_3^h - \mathbf{R}_{33}^h}{h^2} \right\|_{L^2} \\ &\quad + \left\| \frac{\mathbf{R}_{33}^h - 1}{h^2} \right\|_{L^2} \end{aligned} \quad (4.24)$$

The last two terms are bounded due to (4.8) and (4.16). Due to the fact that for $h \leq 1$ we have

$$\|\nabla \mathbf{p}\|_{L^2(\Omega; \mathbb{R}^{3 \times 3})} \leq C \|\nabla_h \mathbf{p}\|_{L^2(\Omega; \mathbb{R}^{3 \times 3})}$$

for some $C > 0$ independent of $\mathbf{y}^h$ and h we directly have that $\frac{\mathbf{y}^h - \mathbf{z}^h}{h^2}$ is bounded in $W^{1,2}(\Omega; \mathbb{R}^3)$ from Poincaré inequality and the fact that

$$\int_{\Omega} \frac{\mathbf{y}^h - \mathbf{z}^h}{h^2} = 0.$$

From the fact that for every $\hat{x} \in \omega$ we also have that

$$\int_{-1/2}^{1/2} \frac{\mathbf{y}^h - \mathbf{z}^h}{h^2} = 0,$$

we can using Theorem 4 conclude that there exist a subsequence (not relabeled) and $\mathbf{k} \in L^2(\omega; W_{\#Y}^{1,2}(S \times Y; \mathbb{R}^3))$ such that

$$\nabla_h \frac{1}{h^2}(\mathbf{y}^h - \mathbf{z}^h) \rightharpoonup (\partial_{y_1} \mathbf{k}(x, y), \partial_{y_2} \mathbf{k}(x, y), \frac{1}{\gamma} \partial_{x_3} \mathbf{k}(x, y)). \quad (4.25)$$

Let us note that the sequences $\left(\frac{h \partial_{\alpha} v^h + \mathbf{R}_{\alpha 3}}{h^2} \right)_{h>0}$ are bounded in $L^2(\omega)$ due to the fact

$$\begin{aligned} \left\| \frac{h \partial_{\alpha} v^h + \mathbf{R}_{\alpha 3}}{h^2} \right\|_{L^2} &\leq \left\| \frac{\int_{-1/2}^{1/2} \partial_{\alpha} \mathbf{y}_3 - \mathbf{R}_{3\alpha}^h}{h^2} \right\|_{L^2} \\ &\quad + \left\| \frac{\mathbf{R}_{\alpha 3}^h + \mathbf{R}_{3\alpha}^h}{h^2} \right\|_{L^2}. \end{aligned} \quad (4.26)$$

The both terms on the right hand side are bounded due to (4.8) and (4.16). On a subsequence (again not relabeled) we have that there exists $\mathbf{l} \in L^2(\omega \times Y; \mathbb{R}^2)$ such that for $\alpha = 1, 2$

$$\frac{h \partial_{\alpha} v^h + \mathbf{R}_{\alpha 3}}{h^2} \rightharpoonup \mathbf{l}_{\alpha} \quad (4.27)$$

Also the sequences $\frac{\mathbf{R}_{\alpha 3}^h}{h}$ are bounded in $W^{1,2}(\omega)$ and from Theorem 3 and the properties (4.12) and (4.16) we have that there exists $\mathbf{r} \in (L^2(\omega; W_{\#}^{1,2}(Y)))^2$ such that

$$\frac{\partial_{\beta} \mathbf{R}_{\alpha 3}^h}{h} \rightharpoonup -\partial_{\alpha\beta} v + \partial_{y_{\beta}} \mathbf{r}_{\alpha} \quad (4.28)$$

By Theorem 3 and (4.11) we conclude that there exists $\mathbf{u}^1 \in (L^2(\omega; \dot{W}_{\#}^{1,2}(Y)))^2$ such that (on a subsequence) we have

$$\nabla \mathbf{u}^h \rightharpoonup \nabla_{\hat{x}} \mathbf{u} + \nabla_y \mathbf{u}^1.$$

By using Lemma 4 where we take $z_\alpha^h = \frac{\mathbf{R}_{\alpha 3}^h}{h}$, $s^h = v^h$ we conclude that there exist $\mathbf{m} \in (L^2(\omega))^2$, $p \in L^2(\omega; \dot{W}_\#^{1,2}(Y))$

$$\text{sym} \frac{\nabla_h \mathbf{z}^h - \mathbf{I}}{h^2} \rightharpoonup \left(\begin{array}{c|c} \text{sym} \nabla_{\hat{x}} \mathbf{u} + \text{sym} \nabla_y \mathbf{u}^1 - x_3 \nabla_{\hat{x}}^2 v + \frac{x_3}{\gamma} \text{sym} \nabla_y \mathbf{r} & \frac{1}{2} (\frac{1}{\gamma} \mathbf{r} + \mathbf{m} + \nabla_y p) \\ \hline \frac{1}{2} (\frac{1}{\gamma} \mathbf{r} + \mathbf{m} + \nabla_y p)^T & 0 \end{array} \right)$$

By taking $\mathbf{w}(x, y) = \mathbf{k}(x, y) + \begin{bmatrix} \mathbf{u}^1(\hat{x}, y) + \frac{x_3}{\gamma} \mathbf{r}(\hat{x}, y) + x_3 \mathbf{m}(\hat{x}) \\ p(\hat{x}, y) - \frac{x_3}{2} \|\nabla_{\hat{x}} v(\hat{x})\|^2 \end{bmatrix}$ we have the claim, from (4.16), (4.19) and (4.20).

Let us by $Q_2^H : \omega \times \mathbb{R}^{2 \times 2} \times \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$ denote the functional

$$Q_2^H(\hat{x}, \mathbf{G}_1'', \mathbf{G}_2'') = \inf_{\mathbf{w} \in W_{\#Y}^{1,2}([-\frac{1}{2}, \frac{1}{2}] \times Y; \mathbb{R}^3)} \int_{-1/2}^{1/2} \int_Y Q_3(\hat{x}, x_3, y) [\mathbf{M}(\mathbf{G}_1'', \mathbf{G}_2'', \tilde{\nabla}_{1,\gamma} \mathbf{w})] dy dx_3, \quad (4.29)$$

where $\mathbf{M}(\mathbf{G}_1'', \mathbf{G}_2'', \tilde{\nabla}_{1,\gamma} \mathbf{w})$ is given by

$$\mathbf{M}(\mathbf{G}_1'', \mathbf{G}_2'', \tilde{\nabla}_{1,\gamma} \mathbf{w}) = \text{sym} \left(\begin{array}{c|c} \mathbf{G}_1'' - x_3 \mathbf{G}_2'' + \nabla_y \begin{pmatrix} \mathbf{w}_1(x, y) \\ \mathbf{w}_2(x, y) \end{pmatrix} & \begin{pmatrix} \frac{1}{\gamma} \partial_{x_3} \mathbf{w}_1(x, y) \\ \frac{1}{\gamma} \partial_{x_3} \mathbf{w}_2(x, y) \end{pmatrix} \\ \hline \partial_{y_1} \mathbf{w}_3(x, y) \quad \partial_{y_2} \mathbf{w}_3(x, y) & \frac{1}{\gamma} \partial_{x_3} \mathbf{w}_3(x, y) \end{array} \right)$$

By $I^L : W^{1,2}(\omega; \mathbb{R}^2) \times W^{2,2}(\omega) \rightarrow \mathbb{R}_0^+$ we denote the functional

$$I^L(\mathbf{u}, v) = \int_\omega Q_2^H(\text{sym} \nabla \mathbf{u} + \frac{1}{2} \nabla v \otimes \nabla v, \nabla^2 v) dx, \quad (4.30)$$

and by $I_E^L : W^{1,2}(\omega; \mathbb{R}^2) \times W^{2,2}(\omega) \times L^2(\omega; \mathcal{V}) \rightarrow \mathbb{R}_0^+$ the functional

$$I_E^L(\mathbf{u}, v, \boldsymbol{\alpha}) = \int_\omega \int_{-1/2}^{1/2} \int_Y Q_3(\hat{x}, x_3, y) \left[\mathbf{M} \left(\text{sym} \nabla_{\hat{x}} \mathbf{u} + \frac{1}{2} \nabla_{\hat{x}} v \otimes \nabla_{\hat{x}} v, \nabla_{\hat{x}}^2 v, \tilde{\nabla}_{1,\gamma} \boldsymbol{\alpha} \right) \right] dy dx_3 d\hat{x}, \quad (4.31)$$

where

$$\mathcal{V} = \left\{ \mathbf{w} \in W_{\#Y}^{1,2}([- \frac{1}{2}, \frac{1}{2}] \times Y; \mathbb{R}^3) \mid \int_{[- \frac{1}{2}, \frac{1}{2}] \times Y} \mathbf{w} dy dx_3 = 0 \right\}.$$

Corollary 1 *Let $\mathbf{y}^h, \mathbf{R}^h, \mathbf{u}^h, \mathbf{u}, v^h, v, \mathbf{G}^h, \mathbf{G}$ be as in Lemma 6. Then we have the following semi-continuity results.*

$$\liminf_{h \rightarrow 0} \frac{1}{h^4} I^h(\mathbf{y}^h) \geq I^L(\mathbf{u}, v). \quad (4.32)$$

Proof.

We shall use the truncation, the Taylor expansion and the weak semi-continuity argument as in the proof of Corollary 16 in [9] (see also Lemma 2.3 in [?]). Consider the good set $\Omega_h := \{x \in \Omega : |\mathbf{G}^h(x)| < h^{-1}\}$. Its characteristic function χ_h is bounded and satisfies $\chi_h \rightarrow 1$ a.e. in Ω . Thus we have $\chi_h \mathbf{G}^h \rightharpoonup \mathbf{G}$ in $L^2(\Omega \times Y; \mathbb{R}^{3 \times 3})$ by Lemma 1. By Taylor expansion

$$\frac{1}{h^4} W(x, \frac{\hat{x}}{\varepsilon(h)}, \mathbf{I} + \mathbf{A}) = \frac{1}{2} \frac{\partial^2 W}{\partial \mathbf{F}^2}(x, \frac{\hat{x}}{\varepsilon(h)}, \mathbf{I} + t\mathbf{A})[\mathbf{A}, \mathbf{A}]. \quad (4.33)$$

where $0 < t < 1$ depends on $x, \frac{\hat{x}}{\varepsilon(h)}, \mathbf{A}$. Using the frame-indifference we have from (4.33)

$$\begin{aligned} \frac{1}{h^4} \int_{\Omega} W(x, \frac{x}{\varepsilon(h)}, \nabla_h \mathbf{y}^h) &\geq \frac{1}{h^4} \int_{\Omega} \chi_h W(x, \frac{\hat{x}}{\varepsilon(h)}, \nabla_h \mathbf{y}^h) dx \\ &= \frac{1}{h^4} \int_{\Omega} \chi_h W(x, \frac{\hat{x}}{\varepsilon(h)}, (\mathbf{R}^h)^T \nabla_h \mathbf{y}^h) dx \\ &= \int_{\Omega} \frac{1}{2} \chi_h \frac{\partial^2 W}{\partial \mathbf{F}^2}(x, \frac{\hat{x}}{\varepsilon(h)}, \mathbf{I} + h^2 t_h \mathbf{G}^h) [\mathbf{G}^h, \mathbf{G}^h] dx \end{aligned} \quad (4.34)$$

By Scorza-Dragoni theorem for every $\varepsilon > 0$ there exists a compact subset $K(\varepsilon) \subset \Omega$ such that $\text{meas}(\Omega \setminus K(\varepsilon)) < \varepsilon$ and $\frac{\partial^2 W}{\partial \mathbf{F}^2} \Big|_{K(\varepsilon) \times Y \times \overline{B_\delta(\mathbf{I})}}$ is continuous (hence uniformly continuous). From this we conclude that for an arbitrary $\delta > 0$ and $h < h(\delta)$ we have

$$\begin{aligned} &\frac{1}{2} \int_{\Omega} \chi_h \frac{\partial^2 W}{\partial \mathbf{F}^2}(x, \frac{\hat{x}}{\varepsilon(h)}, \mathbf{I} + h^2 t_h \mathbf{G}^h) [\mathbf{G}^h, \mathbf{G}^h] dx \\ &= \int_{K(\varepsilon)} \frac{1}{2} \left(\chi_h \frac{\partial^2 W}{\partial \mathbf{F}^2}(x, \frac{\hat{x}}{\varepsilon(h)}, \mathbf{I} + h^2 t_h \mathbf{G}^h) [\mathbf{G}^h, \mathbf{G}^h] - Q_3(x, \frac{\hat{x}}{\varepsilon(h)}) [\chi_h \mathbf{G}^h] \right) dx \\ &\quad + \frac{1}{2} \int_{K(\varepsilon)} Q_3(x, \frac{\hat{x}}{\varepsilon(h)}) [\chi_h \mathbf{G}^h] dx \\ &\geq -\delta \int_{K(\varepsilon)} \|\mathbf{G}^h\|^2 + \frac{1}{2} \int_{K(\varepsilon)} Q_3(x, \frac{\hat{x}}{\varepsilon(h)}) [\chi_h \mathbf{G}^h] dx. \end{aligned} \quad (4.35)$$

By letting $h \rightarrow 0$ and then $\delta \rightarrow 0$, using Lemma 2 (applied to the function $\chi_{K(\varepsilon)} Q_3$) and the fact that $\chi^h \mathbf{G}^h \rightharpoonup \mathbf{G}$ we conclude that

$$\begin{aligned} & \liminf_{h \rightarrow 0} \int_{\Omega} \frac{1}{2} \chi_h \frac{\partial^2 W}{\partial \mathbf{F}^2} \left(x, \frac{\hat{x}}{\varepsilon(h)}, \mathbf{I} + h^2 t_h \mathbf{G}^h \right) [\mathbf{G}^h, \mathbf{G}^h] dx \\ & \geq \frac{1}{2} \int_{K(\varepsilon)} \int_Y Q_3(x, y) [\mathbf{G}] dy dx \end{aligned} \quad (4.36)$$

By letting $\varepsilon \rightarrow 0$ and using the property IV) and Lemma 6 we have the claim.

Lemma 7 *For a.e. $(x_1, x_2) \in \omega$, every $\mathbf{G}_1'', \mathbf{G}_2'' \in \mathbb{R}^{2 \times 2}$ there exists a unique $\mathbf{w}(x_1, x_2, \mathbf{G}_1'', \mathbf{G}_2'') \in \mathcal{V}$ which minimizes the functional Q_2^H . This minimizer satisfies the estimate*

$$\|\mathbf{w}(x_1, x_2, \mathbf{G}_1'', \mathbf{G}_2'')\|_{W_{\#Y}^{1,2}([-\frac{1}{2}, \frac{1}{2}] \times Y; \mathbb{R}^3)}^2 \leq C(\|\mathbf{G}_1''\|^2 + \|\mathbf{G}_2''\|^2), \quad (4.37)$$

for some $C > 0$, where C does not depend on $(x_1, x_2, \mathbf{G}_1'', \mathbf{G}_2'')$. The minimizer is a linear function of $\mathbf{G}_1'', \mathbf{G}_2''$ i.e. for a.e. $(x_1, x_2) \in \omega$, $i = 1, 2, 3$, $\alpha, \beta = 1, 2$ there exist $\mathbf{A}_{i,\alpha\beta}, \mathbf{B}_{i,\alpha\beta} \in L^\infty(\omega; \mathcal{V})$ such that

$$\mathbf{w}_i(\hat{x}, \mathbf{G}_1'', \mathbf{G}_2'') = \mathbf{A}_i(\hat{x}) \cdot \mathbf{G}_1'' + \mathbf{B}_i(\hat{x}) \cdot \mathbf{G}_2''.$$

The functional $Q_2^H : \omega \rightarrow K_{2 \times 4}$ is a quadratic form for a.e. $\hat{x} \in \omega$ and satisfies the properties

- a) $Q_2^H(\hat{x})[\cdot]$ is continuous for a.e. $\hat{x} \in \omega$.
- b) $Q_2^H(\cdot)[\mathbf{G}_1'', \mathbf{G}_2'']$ is measurable for all $\mathbf{G}_1'', \mathbf{G}_2'' \in \mathbb{R}^{2 \times 2}$.
- c) there exists $\gamma_1 > 0$ such that for a.e. $x \in \Omega$ we have

$$Q_2^H(\hat{x})[\mathbf{G}_1'', \mathbf{G}_2''] \leq \gamma_1(\|\mathbf{G}_1''\|^2 + \|\mathbf{G}_2''\|^2), \quad \forall \mathbf{G}_1'', \mathbf{G}_2'' \in \mathbb{R}^{2 \times 2}$$

- d) Q_2^H is positive semi-definite, zero on antisymmetric matrices and uniformly positive definite on symmetric matrices i.e. there exists $C > 0$ such that for a.e. $x \in \Omega$ and all $y \in \mathbb{R}^2$ we have

$$\begin{aligned} Q_2^H(\hat{x})[\mathbf{G}_1'', \mathbf{G}_2''] &= Q_2^H(\hat{x})[\text{sym } \mathbf{G}_1'', \text{sym } \mathbf{G}_2''], \\ Q_2^H(\hat{x})[\mathbf{G}_1'', \mathbf{G}_2''] &\geq C(\|\text{sym } \mathbf{G}_1''\|^2 + \|\text{sym } \mathbf{G}_2''\|^2), \quad \forall \mathbf{G}_1'', \mathbf{G}_2'' \in \mathbb{R}^{2 \times 2} \end{aligned}$$

Proof.

Let us for fixed $\hat{x} \in \omega$, $\mathbf{G}_1'', \mathbf{G}_2'' \in \mathbb{R}^{2 \times 2}$ by $I^H(\hat{x}, \mathbf{G}_1'', \mathbf{G}_2'') : \mathcal{V} \rightarrow \mathbb{R}_+^0$ denote the functional

$$I^H(\mathbf{w}) = \int_{-1/2}^{1/2} \int_Y Q_3(\hat{x}, x_3) [\mathbf{M}(\mathbf{G}_1'', \mathbf{G}_2'', \tilde{\mathbf{V}}_{1,\gamma} \mathbf{w})] dy dx_3$$

The existence of the minimum of the functional I^H is guaranteed by the following facts:

1. for a.e. $\hat{x} \in \omega$ and every $\mathbf{G}_1'', \mathbf{G}_2''$ the functional Q_2^H is sequentially weakly lower semi continuous in $\mathcal{V}$ by the convexity of the form Q_3 on symmetric matrices (see the property V)).
2. for a.e. $\hat{x} \in \omega$, for each $\mathbf{G}_1'', \mathbf{G}_2''$ the functional I^H is coercive in the sense that there exists $C_1 > 0$, $C_2 \in \mathbb{R}$ such that for $\mathbf{w} \in \mathcal{V}$ we have

$$I^H(\hat{x}, \mathbf{G}_1'', \mathbf{G}_2'')(\mathbf{w}) \geq C_1 \|\mathbf{w}\|_{W_{\#Y}^{1,2}([-\frac{1}{2}, \frac{1}{2}] \times Y; \mathbb{R}^3)}^2 + C_2.$$

This follows from Theorem 5 and the fact that for a.e. $\hat{x} \in \omega$, for each $\mathbf{G}_1'', \mathbf{G}_2''$, from the property V) of Q_3 , we have

$$I^H(\hat{x}, \mathbf{G}_1'', \mathbf{G}_2'')(\mathbf{w}) \geq C(\|\text{sym } \tilde{\nabla}_{1,\gamma} \mathbf{w}\|_{L^2([-\frac{1}{2}, \frac{1}{2}] \times Y; \mathbb{R}^{3 \times 3})}^2 - \|\text{sym}(\mathbf{G}_1'' + x_3 \mathbf{G}_2'')\|_{L^2([-\frac{1}{2}, \frac{1}{2}] \times Y; \mathbb{R}^{3 \times 3})}^2).$$

The uniqueness of the minimum is the consequence of the strict convexity of the form Q_3 on symmetric matrices.

Let us by $\mathcal{A}(x, y) : \mathbb{R}_{\text{sym}}^{3 \times 3} \rightarrow \mathbb{R}_{\text{sym}}^{3 \times 3}$ denote the positive definite linear operator which realizes the quadratic functional Q_3 i.e. we have

$$Q_3(x, y)[\mathbf{G}] = (\mathcal{A}(x, y)\mathbf{G}, \mathbf{G}), \forall \mathbf{G} \in \mathbb{R}_{\text{sym}}^{3 \times 3}, \text{ for a.e. } x \in \mathbb{R}^3, y \in \mathbb{R}^2. \quad (4.38)$$

where we have by $(\cdot, \cdot)$ denoted the standard scalar multiplication on $\mathbb{R}^{3 \times 3}$. The property IV) implies that $\|\mathcal{A}\|_{L^\infty(\omega \times Y)} \leq \gamma$. The minimization formulation (4.29) implies that for arbitrary $\hat{x} \in \omega$, $\mathbf{G}_1'', \mathbf{G}_2'' \in \mathbb{R}^{2 \times 2}$ and $\mathbf{t} \in \mathcal{V}$ for the minimizer $\mathbf{w} \in \mathcal{V}$ we have

$$\begin{aligned} & \int_{[-\frac{1}{2}, \frac{1}{2}] \times Y} (\mathcal{A}(\hat{x}, x_3, y) \text{sym } \tilde{\nabla}_{1,\gamma} \mathbf{w}, \text{sym } \tilde{\nabla}_{1,\gamma} \mathbf{t}) dx_3 dy \\ &= \int_{[-\frac{1}{2}, \frac{1}{2}] \times Y} (\mathcal{A}(\hat{x}, x_3, y) (\mathbf{G}_1'' - x_3 \mathbf{G}_2''), \text{sym } \tilde{\nabla}_{1,\gamma} \mathbf{t}) dx_3 dy \end{aligned} \quad (4.39)$$

This implies that $\mathbf{w}$ is the linear function of the pair $(\mathbf{G}_1'', \mathbf{G}_2'')$. By taking $\mathbf{t} = \mathbf{w}$, using Theorem 5, we can conclude the property (4.37). Also from (4.39) it can be easily concluded the property a). The property c) is the direct consequence of the property (4.37). The first part in the property d) is obvious. The second part can be easily seen to be valid. Namely if we take $\mathbf{w} \in \mathcal{V}$, the solution of (4.39), from the property V) of Q_3 and the definition of Q_2^H we conclude

$$\begin{aligned} Q_2^H(\hat{x})[\mathbf{G}_1'', \mathbf{G}_2''] &\geq C \|\text{sym}(\mathbf{G}_1'' - x_3 \mathbf{G}_2'' + \nabla_y(\mathbf{w}_1, \mathbf{w}_2))\|_{L^2([-\frac{1}{2}, \frac{1}{2}] \times Y; \mathbb{R}^{2 \times 2})}^2 \\ &\geq C_1 (\|\text{sym } \mathbf{G}_1''\|^2 + \|\text{sym } \mathbf{G}_2''\|^2). \end{aligned}$$

The property b) can be easily concluded from the definition of Q_2^H and the separability of $W_{\#Y}^{1,2}([-\frac{1}{2}, \frac{1}{2}] \times Y; \mathbb{R}^3)$.

Lemma 8 For arbitrary $\mathbf{u} \in W^{1,2}(\omega; \mathbb{R}^2)$, $v \in W^{2,2}(\omega)$ the following is valid

$$I^L(\mathbf{u}, v) = \inf_{\alpha \in L^2(\omega; \mathcal{V})} I_E^L(\mathbf{u}, v, \alpha) = \min_{\alpha \in L^2(\omega; \mathcal{V})} I_E^L(\mathbf{u}, v, \alpha). \quad (4.40)$$

Proof. That the right hand side is greater or equal to the left hand side is obvious. The opposite inequality follows easily from the estimate (4.37). The existence of the minimizer follows easily from the coercivity and convexity of Q_3 .

4.3 Upper bound

Theorem 8 (optimality of lower bound)

Let $v \in W^{2,2}(\omega)$, $\mathbf{u} \in W^{1,2}(\omega; \mathbb{R}^2)$. Then for each sequence $h \rightarrow 0$ there exists a subsequence, still denoted by h and appropriate $\mathbf{y}^h \in W^{1,2}(\Omega; \mathbb{R}^3)$ such that for $\mathbf{u}^h$, v^h defined by (4.7) (where $\tilde{\mathbf{y}}^h$ should be replaced by $\mathbf{y}^h$) convergence (4.10)-(4.11) are valid and

$$\lim_{h \rightarrow 0} \frac{1}{h^4} I^h(\mathbf{y}^h) = I^L(\mathbf{u}, v). \quad (4.41)$$

Proof.

Let us assume that $\mathbf{u} \in C^\infty(\omega; \mathbb{R}^2)$, $v \in C^\infty(\omega)$. Let us take an arbitrary $\alpha \in (C^\infty(\Omega; C^\infty_\#(Y)))^3$. Then we define

$$\begin{aligned} \mathbf{y}^h(x_1, x_2, x_3) &= \begin{pmatrix} x_1 \\ x_2 \\ hx_3 \end{pmatrix} + \begin{pmatrix} h^2 \mathbf{u}_1(x_1, x_2) \\ h^2 \mathbf{u}_2(x_1, x_2) \\ hv(x_1, x_2) \end{pmatrix} \\ &\quad - h^2 x_3 \begin{pmatrix} \partial_1 v(x_1, x_2) \\ \partial_2 v(x_1, x_2) \\ 0 \end{pmatrix} \\ &\quad + h^2 \varepsilon(h) \alpha(x_1, x_2, x_3, \frac{x_1}{\varepsilon(h)}, \frac{x_2}{\varepsilon(h)}) \\ &\quad - \frac{h^3}{2} \|\nabla_{\hat{x}} v\|^2 \mathbf{e}_3 \end{aligned} \quad (4.42)$$

We calculate

$$\begin{aligned} \nabla_h \mathbf{y}^h &= \mathbf{I} + h \begin{pmatrix} 0 & 0 & -\partial_{x_1} v \\ 0 & 0 & -\partial_{x_2} v \\ \partial_{x_1} v & \partial_{x_2} v & 0 \end{pmatrix} \\ &\quad + h^2 \begin{pmatrix} \partial_{x_1} \mathbf{u}_1 + \partial_{y_1} \alpha_1 & \partial_{x_2} \mathbf{u}_1 + \partial_{y_2} \alpha_1 & \frac{\varepsilon(h)}{h} \partial_{x_3} \alpha_1 \\ \partial_{x_1} \mathbf{u}_2 + \partial_{y_1} \alpha_2 & \partial_{x_2} \mathbf{u}_2 + \partial_{y_2} \alpha_2 & \frac{\varepsilon(h)}{h} \partial_{x_3} \alpha_2 \\ \partial_{y_1} \alpha_1 & \partial_{y_2} \alpha_2 & \frac{\varepsilon(h)}{h} \partial_{x_3} \alpha_3 - \frac{1}{2} \|\nabla_{\hat{x}} v\|^2 \end{pmatrix} \end{aligned}$$

$$-h^2 x_3 \begin{pmatrix} \partial_{x_1 x_1} v & \partial_{x_1 x_2} v & 0 \\ \partial_{x_1 x_2} v & \partial_{x_2 x_2} v & 0 \\ 0 & 0 & 0 \end{pmatrix} + O(h^3), \quad (4.43)$$

where $\|O(h^3)\|_{L^\infty} \leq Ch^3$.

Using the identities $(\mathbf{I} + \mathbf{A})^T(\mathbf{I} + \mathbf{A}) = \mathbf{I} + 2\text{sym } \mathbf{A} + \mathbf{A}^T \mathbf{A}$ and $(\mathbf{e}_3 \otimes \mathbf{a}' - \mathbf{a}' \otimes \mathbf{e}_3)^T(\mathbf{e}_3 \otimes \mathbf{a}' - \mathbf{a}' \otimes \mathbf{e}_3) = \mathbf{a}' \otimes \mathbf{a}' + |\mathbf{a}'|^2 \mathbf{e}_3 \otimes \mathbf{e}_3$ for $\mathbf{a}' \in \mathbb{R}^2$ we obtain

$$\begin{aligned} (\nabla \mathbf{y}^h)^T (\nabla \mathbf{y}^h) &= \mathbf{I} + h^2 [2\text{sym } \mathbf{B} + \nabla_{\hat{x}} v \otimes \nabla_{\hat{x}} v] \\ &\quad - 2h^2 x_3 \nabla_{\hat{x}}^2 v + O(h^3). \end{aligned} \quad (4.44)$$

Here

$$\mathbf{B} = \begin{pmatrix} \partial_{x_1} \mathbf{u}_1 + \partial_{y_1} \boldsymbol{\alpha}_1 & \partial_{x_2} \mathbf{u}_1 + \partial_{y_2} \boldsymbol{\alpha}_1 & \frac{\varepsilon(h)}{h} \partial_{x_3} \boldsymbol{\alpha}_1 \\ \partial_{x_1} \mathbf{u}_2 + \partial_{y_1} \boldsymbol{\alpha}_2 & \partial_{x_2} \mathbf{u}_2 + \partial_{y_2} \boldsymbol{\alpha}_2 & \frac{\varepsilon(h)}{h} \partial_{x_3} \boldsymbol{\alpha}_2 \\ \partial_{y_1} \boldsymbol{\alpha}_1 & \partial_{y_2} \boldsymbol{\alpha}_2 & \frac{\varepsilon(h)}{h} \partial_{x_3} \boldsymbol{\alpha}_3 \end{pmatrix}. \quad (4.45)$$

Taking the square root in (4.44) and using the frame indifference of W , the Taylor expansion and the unfolding isometry (see [24, 25]) we get

$$\frac{1}{h^4} \int_{\Omega} W(\nabla \mathbf{y}^h) = \frac{1}{h^4} \int_{\Omega} W([\nabla \mathbf{y}^h]^T \nabla \mathbf{y}^h)^{1/2} \rightarrow I_E^L(\mathbf{u}, v, \boldsymbol{\alpha}).$$

By Lemma 8 we have to prove that we can construct the appropriate sequence for an arbitrary $\boldsymbol{\alpha} \in L^2(\omega; \mathcal{V})$. But this can be easily justified by the density argument and diagonalization. In the same way we can prove the theorem for arbitrary $\mathbf{u} \in W^{1,2}(\omega; \mathbb{R}^2)$, $v \in W^{2,2}(\omega)$.

Lemma 5, Lemma 6 and Theorem 8 enable us to state the standard theorem on convergence of minimizers. We shall state it without proof (since it is standard), assuming the external loads in $\mathbf{e}_3$ direction. For a more detailed discussion on external loads in the standard Föppl-von Kármán case see [14].

Let $\mathbf{f}_3 \in L^2(\omega)$ be given with the property

$$\int_{\omega} \mathbf{f}_3 dx = 0. \quad (4.46)$$

Let us just mention that it is not necessary to demand that $\mathbf{f}_3$ depends only on x_1, x_2 , but this simplifies the proof.

Let us additionally assume

$$\int_{\omega} x_1 \mathbf{f}_3 dx = 0, \quad \int_{\omega} x_2 \mathbf{f}_3 dx = 0. \quad (4.47)$$

This can be assumed by rotating the coordinate axis (the energy density W is rotational invariant).

We shall assume that the order of the external loads is h^2 and that the total energy functional J^h defined on the space $W^{1,2}(\Omega; \mathbb{R}^3)$, is given by

$$J^h(\mathbf{y}^h) = I^h(\mathbf{y}^h) - \int_{\Omega} h^2 \mathbf{f}_3 \mathbf{y}_3^h. \quad (4.48)$$

The following theorem is the main result and its proof follows the proof of Theorem 2 in [9].

Theorem 9 (Γ -convergence) *Let us suppose that $\mathbf{f}_3 \in L^2(\omega; \mathbb{R})$ is given and satisfies (4.46) and (4.47). Then:*

1. *There exists $C_1, C_2 > 0$ such that for every $h > 0$ we have*

$$0 \geq \inf \left\{ \frac{1}{h^4} J^h(\mathbf{y}^h); \mathbf{y}^h \in W^{1,2}(\Omega; \mathbb{R}^3) \right\} \geq -C_2. \quad (4.49)$$

2. *If $\mathbf{y}^h \in W^{1,2}(\Omega; \mathbb{R}^3)$ is a minimizing sequence of $\frac{1}{h^4} J^h$, that is*

$$\lim_{h \rightarrow 0} \left(\frac{1}{h^4} J^h(\mathbf{y}^h) - \inf \frac{1}{h^4} J^h \right) = 0, \quad (4.50)$$

then we have that there exists $\bar{\mathbf{R}}^h \in \text{SO}(3)$, $\mathbf{c}^h \in \mathbb{R}$ such that the sequence $(\bar{\mathbf{R}}^h, \mathbf{y}^h)$ has its subsequence (also not relabeled) with the following property:

i) *For the sequence $\tilde{\mathbf{y}}^h := (\bar{\mathbf{R}}^h)^T \mathbf{y}^h - \mathbf{c}^h$ and $\mathbf{u}^h, v^h$ defined by (4.7) the following is valid*

$$\begin{aligned} \mathbf{u}^h &\rightharpoonup \mathbf{u} \quad \text{weakly in } W^{1,2}(\omega; \mathbb{R}^2), \\ v^h &\rightarrow v \quad \text{in } W^{1,2}(\omega), \quad v \in W^{2,2}(\omega). \end{aligned}$$

Any accumulation point $(\mathbf{u}, v, \bar{\mathbf{R}})$ of the sequence $(\mathbf{u}^h, v^h, \bar{\mathbf{R}}^h)$ minimizes the functional

$$J^L(\mathbf{u}, v, \bar{\mathbf{R}}) = I^L(\mathbf{u}, v) - \bar{\mathbf{R}}_{33} \int_{\omega} \mathbf{f}_3 v, \quad (4.51)$$

where I^L is defined in (4.30). Moreover, for $\mathbf{f}_3 \neq 0$ we have $\bar{\mathbf{R}}_{33} = 1$ or $\bar{\mathbf{R}}_{33} = -1$.

3. *The minimum of the functional J^L exists in the space $W^{1,2}(\omega; \mathbb{R}^2) \times W^{2,2}(\omega) \times \text{SO}(3)$. If $\mathbf{y}^h \in W^{1,2}(\Omega; \mathbb{R}^3)$ is a minimizing sequence (not relabeled) of $\frac{1}{h^4} J^h$ then we have that*

$$\lim_{h \rightarrow 0} \frac{1}{h^4} J^h(\mathbf{y}^h) = \min_{\mathbf{u} \in W^{1,2}(\omega; \mathbb{R}^2), v \in W^{2,2}(\omega), \bar{\mathbf{R}} \in \text{SO}(3)} J^L(\mathbf{u}, v, \bar{\mathbf{R}}). \quad (4.52)$$

References

1. G. Allaire, *Homogenization and two scale convergence*, SIAM Journal on Mathematical Analysis 23(6) (1992), 1482–1518.
2. J.F. Babadjian, and M. Baía, *3D-2D analysis of a thin film with periodic microstructure*, Proceedings of the Royal Society of Edinburgh, Section A, 136, no. 2 (2006), 223–243.
3. A. Braides, " Γ -convergence for Beginners", Oxford University Press, Oxford, 2002.
4. Braides A., Fonseca, I., Francfort, G.: 3D-2D Asymptotic Analysis for Inhomogeneous Thin Films, Indiana Univ. Math. J., 49, 1367–1404 (2000).
5. P.G. Ciarlet, "Mathematical elasticity. Vol. II. Theory of plates. Studies in Mathematics and its Applications", 27. North-Holland Publishing Co., Amsterdam 1997.
6. P.G. Ciarlet, "Mathematical elasticity. Vol. III. Theory of shells. Studies in Mathematics and its Applications", 29. North-Holland Publishing Co., Amsterdam 2000.
7. G. Dal Maso, "An introduction to Γ -convergence, Progress in Nonlinear Differential Equations and Their Applications", Birkhäuser, Basel 1993.
8. G. Friesecke, R.D. James and S. Müller, *A theorem on geometric rigidity and the derivation of nonlinear plate theory from three-dimensional elasticity*, Comm. Pure Appl. Math. 55 (2002), 1461–1506.
9. G. Friesecke, R.D. James and S. Müller, *A Hierarchy of Plate Models Derived from Nonlinear Elasticity by Γ -Convergence*, Archive for Rational Mechanics and Analysis 180, no.2 (2006), 183–236.
10. G. Friesecke, R.D. James and S. Müller, *The Föppl-von Kármán plate theory as a low energy Γ -limit of nonlinear elasticity*, Comptes Rendus Mathématique 335, no. 2 (2002), 201–206.
11. G. Friesecke, R. James, M.G. Mora and S. Müller, *Derivation of nonlinear bending theory for shells from three-dimensional nonlinear elasticity by Γ -convergence*, C. R. Math. Acad. Sci. Paris, 336, no. 8 (2003), 697–702.
12. H. Le Dret and A. Raoult, *The nonlinear membrane model as a variational limit of nonlinear three-dimensional elasticity*, Journal de Mathématiques Pures et Appliquées 74 (1995), 549–578.
13. H. Le Dret and A. Raoult, *The membrane shell model in nonlinear elasticity: A variational asymptotic derivation*, Journal of Nonlinear Science 6, Number 1 (1996), 59–84.
14. M. Lecumberry and S. Müller, *Stability of slender bodies under compression and validity of the von Kármán theory*, Archive for Rational Mechanics and Analysis, Volume 193, Number 2 (2009), 255–310.
15. M. Lewicka, M.G. Mora and M. Pakzad, *Shell theories arising as low energy Γ -limit of 3d nonlinear elasticity*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. 5, Vol. IX (2010), 1–43.
16. Lewicka, M., Pakzad, M.: The infinite hierarchy of elastic shell models: some recent results and a conjecture, accepted in Fields Institute Communications (2010).
17. S. Müller, *Homogenization of nonconvex integral functionals and cellular elastic materials*, Archive for Rational Mechanics and Analysis 99(1987), 189–212.
18. Neukamm, S.: Homogenization, linearization and dimension reduction in elasticity with variational methods, Dissertation Technische Universität München, 2010.
19. Neukamm, S.: Rigorous derivation of a homogenized bending-torsion theory for inextensible rods from 3d elasticity, preprint.
20. G. Nguetseng, *A general convergence result for a functional related to the theory of homogenization*, SIAM Journal on Mathematical Analysis 20(3) (1989), 608–625.
21. G.A. Pavliotis, and A.M. Stuart, "Multiscale Methods: Averaging and Homogenization", Springer: New York 2008.
22. R. Temam, "Navier Stokes Equations and Nonlinear Functional Analysis", Second Edition, SIAM Society for Industrial and Applied Mathematics: Philadelphia 1995.
23. I. Velčić, *Periodically wrinkled plate of the Föppl-von Kármán type*, to appear in Ann. Scuola Norm. Sup. Pisa Cl. Sci.
24. A. Visintin, *Towards a two-scale calculus*, Control, Optimisation and Calculus of Variations 12(3) (2006), 371–397.
25. A. Visintin, *Two-scale convergence of some integral functionals*, Calculus of Variations and Partial Differential Equations 29 (2) (2007), 239–265.