

FLOW WITH $A_\infty(\mathbb{R})$ DENSITY AND TRANSPORT EQUATION IN $BMO(\mathbb{R})$

RENJIN JIANG¹, KANGWEI LI^{1,2} and JIE XIAO³

¹*Center for Applied Mathematics, Tianjin University, Tianjin, China;
rejiang@tju.edu.cn, kangwei.nku@gmail.com*

²*Basque Center for Applied Mathematics, Alameda de Mazarredo 14, Bilbao, Spain*

³*Dept of Math & Stat, Memorial University, St. John's, NL A1C 5S7, Canada;
jxiao@mun.ca*

Received 18 April 2019

Abstract

We show that, if $b \in L^1(0, T; L^1_{\text{loc}}(\mathbb{R}))$ has spatial derivative in the John-Nirenberg space $BMO(\mathbb{R})$, then it generates a unique flow $\phi(t, \cdot)$ which has an $A_\infty(\mathbb{R})$ density for each time $t \in [0, T]$. Our condition on the map b is not only optimal but also produces a sharp quantitative estimate for the density. As a killer application we achieve the well-posedness for a Cauchy problem of the transport equation in $BMO(\mathbb{R})$.

2010 Mathematics Subject Classification: 42B37 (primary); 34A12 (secondary)

1. Description of the main results

Given an integer $n \geq 1$, a real $T \geq t > 0$ and an evolutionary self-map $b(t, \cdot)$ of \mathbb{R}^n with

$$b \in L^1(0, T; L^1_{\text{loc}}(\mathbb{R}^n)),$$

consider the flow

$$\phi(t, x) = x + \int_0^t b(r, \phi(r, x)) dr.$$

We are motivated by the composition and transportation problems in BMO space to answer the question:

What condition is needed on a vector field such that it generates a flow ϕ that preserves BMO functions?

On \mathbb{R}^n , $n \geq 2$, the question has a satisfactory solution by the seminal work of Reimann [27] via the following (Q)-condition

$$(Q) \quad \sup_{(x,y,z) \in \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^n, |y|=|z|>0} \left| \frac{\langle y, b(x+y) - b(x) \rangle}{|y|^2} - \frac{\langle z, b(x+z) - b(x) \rangle}{|z|^2} \right| < \infty$$

which is equivalent to the anti-conformal part

$$S_A b = \frac{1}{2}(Db + Db^T) - \frac{\operatorname{div} b}{n} I_{n \times n}$$

is bounded - moreover (cf. [27]) -

$$S_A b \in L^\infty(\mathbb{R}^n) \Rightarrow Db \in \operatorname{BMO}(\mathbb{R}^n).$$

More precisely, [27] shows if b satisfies (Q) then it generates a unique flow ϕ , which at each time t is a quasi-conformal mapping; see also [5]. By using the composition result on BMO by Reimann [26], one sees that the flow ϕ preserves BMO; see [10] for an application of Reimann's result to the transportation.

However, less known is the situation on \mathbb{R} . According to Jones [21], a homeomorphism $\phi : \mathbb{R} \rightarrow \mathbb{R}$ preserves BMO, if and only if, ϕ' is an A_∞ weight. Recall that a non-negative locally integrable function w is an A_∞ weight, if

$$[w]_{A_\infty(\mathbb{R}^n)} = \sup_{\text{cubes } I \subset \mathbb{R}^n} \left(\frac{1}{|I|} \int_I w(x) dx \right) \exp \left(-\frac{1}{|I|} \int_I (\log w(x)) dx \right) < \infty.$$

Note that the Reimann's (Q)-condition coincides with the Zygmund condition for a constant $C > 0$:

$$(Z) \quad |b(x+y) + b(x-y) - 2b(x)| \leq C|y| \quad \forall (x,y) \in \mathbb{R} \times \mathbb{R},$$

on the line. Reimann also [27] showed that for functions satisfying (Q) the induced flows are quasi-symmetric mappings. Unfortunately, quasi-symmetric mappings are not necessarily absolutely continuous in \mathbb{R} and a function satisfying (Z) needs not be absolutely continuous (cf. [3, 27] and [14]), in particular, this implies that the induced flows may do not have a density as A_∞ weight.

In view of this, some more restrictions on b seem to be necessary for the generated flow to have an A_∞ density in order to preserve BMO functions. Moreover, we note that Reimann's approach [27] is rather intrinsic for the quasi-conformal /quasi-symmetric mappings, and in \mathbb{R}^n , $n \geq 2$, [27] also obtained rather sharp estimate for the density (see also [5]), but in the line, it does not give enough information on the density of the flow (as the flow may not be absolutely continuous).

In this paper, we show that if b' is of $\text{BMO}(\mathbb{R})$ then b generates a (unique) flow with $A_\infty(\mathbb{R})$ density. To see this clearly, recall that

$$f \in \text{BMO}(\mathbb{R}^n) \Leftrightarrow \|f\|_{\text{BMO}(\mathbb{R}^n)} = \sup_{\text{cubes } I \subset \mathbb{R}^n} |I|^{-1} \int_I |f(x) - f_I| dx < \infty,$$

where

$$f_I = |I|^{-1} \int_I f(x) dx$$

denotes the integral average of f over I whose Lebesgue measure is written as $|I|$. Since all constant functions have zero $\text{BMO}(\mathbb{R}^n)$ -norm, and any constant does effect the flow, we choose the $\text{BMO}(\mathbb{R}^n)$ -norm of f as

$$\|f\|_* = \|f\|_{\text{BMO}(\mathbb{R}^n)} + \int_{B(0,1)} |f(x)| dx,$$

where $B(0, 1)$ is the unit ball of \mathbb{R}^n . Obviously,

$$f \in \text{BMO}(\mathbb{R}^n) \Leftrightarrow \|f\|_* < \infty,$$

however, $\|f\|_*$ is not comparable to $\|f\|_{\text{BMO}(\mathbb{R}^n)}$. In what follows,

$$\frac{\partial}{\partial x} b(t, x) \in L^1(0, T; \text{BMO}(\mathbb{R})) \text{ or } \frac{\partial}{\partial x} b(t, x) \in L^1(0, T; L^\infty(\mathbb{R}))$$

stands for

$$\int_0^T \left\| \frac{\partial}{\partial x} b(t, x) \right\|_* dt < \infty \text{ or } \int_0^T \left\| \frac{\partial}{\partial x} b(t, x) \right\|_{L^\infty(\mathbb{R})} dt < \infty.$$

Our first main result reads as follows.

Theorem 1.1. *Let $b(t, x) : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be in*

$$L^1(0, T; L^1_{\text{loc}}(\mathbb{R})) \text{ with } \frac{\partial b(t, x)}{\partial x} \in L^1(0, T; \text{BMO}(\mathbb{R})). \quad (1)$$

Then there exists a unique flow $\phi(t, x)$ satisfying

$$\begin{cases} \frac{\partial}{\partial t} \phi(t, x) = b(t, \phi(t, x)) & \forall (t, x) \in [0, T] \times \mathbb{R}; \\ \phi(0, x) = x & \forall x \in \mathbb{R}. \end{cases}$$

Moreover, for each $t \in [0, T]$,

$$\left| \frac{\partial}{\partial x} \phi(t, x) \right|$$

is an $A_\infty(\mathbb{R})$ -weight, and there exist constants $C_1, c > 0$ such that

$$\left\| \log \left| \frac{\partial}{\partial x} \phi(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \leq \frac{\int_0^t C_1 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds}{\exp\left(-c \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds\right)}. \quad (2)$$

Some remarks are in order. First, from the well-known fact that the logarithm of an A_∞ weight is a BMO function (see Lemma 2.4) and the formula

$$\log \left| \frac{\partial}{\partial x} \phi(t, x) \right| = \int_0^t \frac{\partial}{\partial x} b(s, \phi(s, x)) ds,$$

we see that our condition (1) is critical, i.e., for each t ,

$$x \mapsto \frac{\partial}{\partial x} b(t, x)$$

is necessarily a $\text{BMO}(\mathbb{R})$ -function. Second, taking

$$b(x) = x \log |x|$$

for example, indicates that b generates a flow ϕ and a constant $C > 0$ with

$$\begin{cases} \phi(t, x) = (\text{sign } x) |x|^{e^t}; \\ \frac{\partial}{\partial x} \phi(t, x) = e^t |x|^{e^t-1} \text{ being an } A_\infty(\mathbb{R}) \text{ weight}; \\ \left\| \left\| \log \left| \frac{\partial}{\partial x} \phi(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \right\| \leq (e^t - 1) \|\log |x|\|_{\text{BMO}(\mathbb{R})} \leq C t e^t. \end{cases}$$

This implies that our estimate (2) is sharp.

For the proof, we shall first provide a version of the result in smooth setting, namely,

$$b \in L^1(0, T; C^1(\mathbb{R})) \text{ with } \frac{\partial b(t, x)}{\partial x} \in L^1(0, T; \text{BMO}(\mathbb{R})), \quad (3)$$

and then use the compactness argument based on the development of non-smooth flows from [2, 9, 12, 13]. Since the Zygmund condition is satisfied for b , existence and uniqueness follow already from Reimann [27]. Accordingly, the key of the proof is to establish (2), which even in the smooth setting seems non-trivial. By the composition result of Jones [21], a homeomorphism ϕ preserves $\text{BMO}(\mathbb{R})$ if and only if ϕ' is an $A_\infty(\mathbb{R})$ weight. However, even we assume that b is smooth on \mathbb{R} , it seems mysteries to us whether one can prove the generated flow carries $A_\infty(\mathbb{R})$ density directly from (1).

In order to overcome the difficulties, we further consider the simpler case

$$b \in L^1(0, T; C^1(\mathbb{R})) \text{ with } \frac{\partial b(t, x)}{\partial x} \in L^1(0, T; L^\infty(\mathbb{R})), \quad (4)$$

where the generated flow carries $A_\infty(\mathbb{R})$ -density from the Cauchy-Lipschitz theory. Then we observe that for a function v with small $\text{BMO}(\mathbb{R})$ -norm, e^v lies in $A_\infty(\mathbb{R})$ with its norm controlled by the $\text{BMO}(\mathbb{R})$ -norm of v linearly. Then by using the flow with $A_\infty(\mathbb{R})$ -density in the smooth setting, a quantitative estimate

for the norm of composition in $\text{BMO}(\mathbb{R})$, and a bootstrap argument, we succeed in showing (2) in the Lipschitz case (4). Finally a truncation argument involving the Arzelá-Ascoli theorem allows us to pass to the case (3); see Section 3.

One may wonder if a quantitative estimate for the $A_\infty(\mathbb{R})$ -norm of

$$\left| \frac{\partial}{\partial x} \phi(t, x) \right|$$

can be established. Although we do not know a positive answer, we doubt it since a quantitative bound for an $A_\infty(\mathbb{R})$ -weight e^v holds only for v with small $\text{BMO}(\mathbb{R})$ -norm; see Lemma 2.3 and Lemma 2.4 below. However, there is a nice result regarding the homeomorphisms preserving $A_p(\mathbb{R})$ -weights by [20].

We next apply the result on flow to study the transportation problem in BMO space. Besides its own interest, this problem and its dual equation also arise naturally from the study of conservation laws (see [6] for instance). In [10] (somewhat related to [25]), a well-posedness of the Cauchy problem of the transport equation in $\text{BMO}(\mathbb{R}^n)$ has been established for $n \geq 2$ and then pushed to the case $n = 1$ in [29]. The main step over there is to use the hypothesis that

$$(t, x) \mapsto \begin{cases} S_A b(t, x) & \forall n \geq 2 \\ \frac{\partial}{\partial x} b(t, x) & \forall n = 1 \end{cases} \text{ belongs to } L^1(0, T; L^\infty(\mathbb{R}^n)),$$

the quasi-conformal flows of [27] and the composition results obtained in [23, 26] for $n \geq 2$ (cf. [22, 28, 30]) and in [21] for $n = 1$. But nevertheless, as our second main result we utilize Theorem 1.1 and [21, Theorem] to discover the following stronger well-posedness of the transport equation in $\text{BMO}(\mathbb{R})$.

Theorem 1.2. *Let $b(t, x) : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be in $L^1(0, T; L^1_{\text{loc}}(\mathbb{R}))$ and satisfy*

$$\frac{\partial b(t, x)}{\partial x} \in L^1(0, T; \text{BMO}(\mathbb{R})).$$

Then for $u_0 \in \text{BMO}(\mathbb{R})$ there exists a unique solution $u \in L^\infty(0, T; \text{BMO}(\mathbb{R}))$ to the Cauchy problem of the transport equation

$$\begin{cases} \left(\frac{\partial u}{\partial t} - b \cdot \nabla u \right) (t, x) = 0 & \forall (t, x) \in (0, T) \times \mathbb{R}; \\ u(0, x) = u_0(x) & \forall x \in \mathbb{R}. \end{cases}$$

Moreover, for each $t \in [0, T]$, it holds that

$$\begin{cases} u(t, x) = u_0(\phi(t, x)); \\ \frac{\partial}{\partial t} \phi(t, x) = b(t, \phi(t, x)), \end{cases}$$

and there exist $C_2, c > 0$ such that

$$\|u\|_{\text{BMO}(\mathbb{R})} \leq C_2 \|u_0\|_{\text{BMO}(\mathbb{R})} \exp \left(c \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds \right). \quad (5)$$

Based on the duality of Hardy space H^1 and BMO by Fefferman and Stein [16], the above theorem provides the existence of a solution in Hardy space H^1 to the continuity equation

$$\begin{cases} \left(\frac{\partial u}{\partial t} - \frac{\partial}{\partial x}(bu) \right)(t, x) = 0 & \forall (t, x) \in (0, T) \times \mathbb{R}; \\ u(0, x) = u_0(x) & \forall x \in \mathbb{R}. \end{cases}$$

See [11] for a study of the equation in higher dimensions and a proof of uniqueness (cf. [11, Theorem 3]).

Note that Mucha gives in [25] a well-posedness of the transport equation in $L^\infty(0, T; L^\infty(\mathbb{R}^n))$ provided $\operatorname{div} b \in L^1(0, T; \text{BMO})$ with compact support. The condition on the vector fields b has been further relaxed in [8]. Nevertheless, let us point out that, the well-posedness in $L^\infty(0, T; L^\infty(\mathbb{R}^n))$ requires much weaker condition on b than one in $L^\infty(0, T; \text{BMO})$. Indeed, given a map $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$, then ϕ preserves L^∞ functions as soon as for any set E with measure zero, the pre-image $\phi^{-1}(E)$ has measure zero. From our previous discussions, the map ϕ preserving BMO functions requires much finer regularity than this.

The paper is organized as follows. In Section 2, we recall and establish some results concerning Muckenhoupt weights, $\text{BMO}(\mathbb{R})$, and continuity estimates. In Section 3, we present the key a priori estimation for the flow, i.e., the version of Theorem 1.1 in the smooth setting. In Section 4, we verify the above main results.

Notation. In the above and below, C, C_1, C_2, \dots and c, c_1, c_2, \dots stand for positive constants.

2. Weights and bounded mean oscillation

For a locally integrable function f and an open interval $I \subset \mathbb{R}$, we denote by f_I the integral average of f on I . We say that a locally integrable nonnegative function w belongs to the Muckenhoupt class $A_p(\mathbb{R})$, $1 < p < \infty$, if

$$[w]_{A_p(\mathbb{R})} = \sup_{\text{intervals } I \subset \mathbb{R}} \left(\frac{1}{|I|} \int_I w(x) dx \right) \left(\frac{1}{|I|} \int_I (w(x))^{\frac{1}{1-p}} dx \right)^{p-1} < \infty,$$

and that $w \in A_\infty(\mathbb{R})$, if

$$[w]_{A_\infty(\mathbb{R})} = \sup_{\text{intervals } I \subset \mathbb{R}} \left(\frac{1}{|I|} \int_I w(x) dx \right) \exp \left(-\frac{1}{|I|} \int_I (\log w(x)) dx \right) < \infty.$$

Note that, if $w > 0$ a.e., then $[w]_{A_\infty(\mathbb{R})} \geq 1$ follows from the Jensen inequality: indeed

$$[w]_{A_\infty(\mathbb{R})} \geq w_I \exp([- \log w]_I) \geq \exp((\log w)_I) \exp([- \log w]_I) = 1,$$

and similarly

$$[w]_{A_p(\mathbb{R})} \geq [w]_{A_\infty(\mathbb{R})} \quad \forall p \in (1, \infty).$$

We need the following quantitative version of the reverse Hölder inequality for $A_\infty(\mathbb{R})$ from [19]; see also [24].

Lemma 2.1. *Let $w \in A_\infty(\mathbb{R})$ and $I \subset \mathbb{R}$ be an arbitrary interval. Then there exist $\tau > 0$, $r_w = 1 + (\tau[w]_{A_\infty(\mathbb{R})})^{-1}$ and $\epsilon_w = (1 + \tau[w]_{A_\infty(\mathbb{R})})^{-1}$ such that*

$$\begin{cases} \left(|I|^{-1} \int_I w^{r_w} dx \right)^{1/r_w} \leq 2|I|^{-1} \int_I w dx; \\ \frac{w(E)}{w(I)} = \frac{\int_E w(x) dx}{\int_I w(x) dx} \leq 2 \left(\frac{|E|}{|I|} \right)^{\epsilon_w} \quad \forall \text{ measurable set } E \subset I. \end{cases}$$

By [21, Theorem], we know that an increasing homeomorphism φ of \mathbb{R} preserves BMO if and only if φ' belongs to $A_\infty(\mathbb{R})$. By using the previous lemma we deduce the following quantitative version; see [1] for an explicit bound in terms of the reverse Hölder index and [4, 15, 17] for the related results.

Lemma 2.2. *Let φ be an increasing homeomorphism on \mathbb{R} with $\varphi' \in A_\infty(\mathbb{R})$. Then*

$$\exists C_3 > 0 \text{ such that } \|f \circ \varphi^{-1}\|_{\text{BMO}(\mathbb{R})} \leq C_3 [\varphi']_{A_\infty(\mathbb{R})} \|f\|_{\text{BMO}(\mathbb{R})}.$$

Proof. Recall that for a $\text{BMO}(\mathbb{R})$ -function f , the John-Nirenberg inequality states that, for all $I \subset \mathbb{R}$, there exists $c_1, c_2 > 0$ such that

$$|\{x \in I : |f(x) - f_I| > \lambda\}| \leq c_1 |I| \exp\left(-\frac{c_2 \lambda}{\|f\|_{\text{BMO}(\mathbb{R})}}\right) \quad \forall \lambda > 0;$$

see [18] for instance.

Suppose that φ is an increasing homeomorphism of \mathbb{R} with $\varphi' \in A_\infty(\mathbb{R})$. By [21, Theorem], we have

$$f \circ \varphi^{-1} \in \text{BMO}.$$

For every interval

$$I = (a, b) \subset \mathbb{R},$$

set

$$E_\lambda = \{x \in I : |f \circ \varphi^{-1}(x) - f_{\varphi^{-1}(I)}| > \lambda\}.$$

Then

$$\varphi^{-1}(E_\lambda) = \{y \in \varphi^{-1}(I) : |f(y) - f_{\varphi^{-1}(I)}| > \lambda\},$$

and hence, by Lemma 2.1 and the John-Nirenberg inequality, we get

$$\frac{|E_\lambda|}{|I|} \leq 2 \left(\frac{|\varphi^{-1}(E_\lambda)|}{|\varphi^{-1}(I)|} \right)^{\epsilon_w} \leq 2c_1 \exp\left(-\frac{c_2 \epsilon_w \lambda}{\|f\|_{\text{BMO}(\mathbb{R})}}\right) \text{ where } \epsilon_w = (1 + \tau[\varphi']_{A_\infty(\mathbb{R})})^{-1},$$

thereby getting via the layer cake representation

$$\|f \circ \varphi^{-1}\|_{\text{BMO}(\mathbb{R})} \leq C(1 + \tau[\varphi']_{A_\infty(\mathbb{R})})\|f\|_{\text{BMO}(\mathbb{R})} \leq C_3[\varphi']_{A_\infty(\mathbb{R})}\|f\|_{\text{BMO}(\mathbb{R})},$$

where we have used the fact that φ is an increasing homeomorphism on \mathbb{R} with

$$[\varphi']_{A_\infty(\mathbb{R})} \geq 1.$$

□

The following result is well-known; see [7, 18] for instance.

Lemma 2.3. *There exists $\alpha < 1 < \beta$ such that for*

$$\begin{cases} f \in \text{BMO}(\mathbb{R}); \\ s \in \mathbb{R}; \\ |s| \leq \alpha\|f\|_{\text{BMO}(\mathbb{R})}^{-1}, \end{cases}$$

it holds that

$$e^{sf} \in A_2(\mathbb{R}) \text{ with } [e^{sf}]_{A_2(\mathbb{R})} \leq \beta^2.$$

Here it is perhaps appropriate to mention that the requirement

$$|s| \leq \alpha\|f\|_{\text{BMO}(\mathbb{R})}^{-1}$$

is critical since

$$x \mapsto f(x) = \log|x|$$

is in $\text{BMO}(\mathbb{R})$ but

$$x \mapsto e^{-f(x)} = |x|^{-1}$$

is not a Muckenhoupt weight.

Lemma 2.4. *If*

$$0 \leq w \in A_\infty(\mathbb{R})$$

then

$$\|\log w\|_{\text{BMO}(\mathbb{R})} \leq 2 \log([w]_{A_\infty(\mathbb{R})} + 1).$$

Conversely, if $v \in \text{BMO}(\mathbb{R})$, then there exists a sufficiently small $\epsilon_0 \in (0, 1]$ such that

$$\|v\|_{\text{BMO}(\mathbb{R})} < \epsilon_0 \Rightarrow e^v \in A_\infty(\mathbb{R}) \text{ with } [e^v]_{A_\infty(\mathbb{R})} \leq 1 + C_4\|v\|_{\text{BMO}(\mathbb{R})}.$$

Proof. On the one hand, for any $0 \leq w \in A_\infty(\mathbb{R})$ we have

$$\begin{aligned} & \int_I \left| \log(w(x)) - (\log w)_I \right| dx \\ &= \int_I [\log(w(x)) - (\log w)_I]_+ dx + \int_I [\log(w(x)) - (\log w)_I]_- dx \\ &= 2 \int_I [\log(w(x)) - (\log w)_I]_+ dx, \end{aligned}$$

where $[f]_+$ and $[f]_-$ denotes the positive and negative parts of f respectively. In virtue of Jensen's inequality we obtain

$$\begin{aligned} & |I|^{-1} \int_I \left| \log(w(x)) - (\log w)_I \right| dx = 2|I|^{-1} \int_I [\log(w(x)) - (\log w)_I]_+ dx \\ & \leq 2 \log \left(|I|^{-1} \int_I \exp[\log(w(x)) - (\log w)_I]_+ dx \right) \\ & \leq 2 \log \left(|I|^{-1} \int_I \exp[\log(w(x)) - (\log w)_I] dx + 1 \right) \\ & \leq 2 \log([w]_{A_\infty(\mathbb{R})} + 1), \end{aligned}$$

whence

$$\|\log w\|_{\text{BMO}(\mathbb{R})} \leq 2 \log([w]_{A_\infty(\mathbb{R})} + 1).$$

On the other hand, note that

$$[e^v]_{A_\infty(\mathbb{R})} = \sup_{I=(a,b) \subset \mathbb{R}} \left(|I|^{-1} \int_I e^{v(x)} dx \right) \exp([-v]_I) = \sup_{I=(a,b) \subset \mathbb{R}} |I|^{-1} \int_I e^{v(x)-v_I} dx. \quad (6)$$

So, if $v \in \text{BMO}(\mathbb{R})$, then the John-Nirenberg inequality gives

$$|\{x \in I : |v(x) - v_I| > \lambda\}| \leq c_1 |I| \exp\left(-\frac{c_2 \lambda}{\|v\|_{\text{BMO}(\mathbb{R})}}\right).$$

Inserting this into (6), we find that if

$$\|v\|_{\text{BMO}(\mathbb{R})} < c_2$$

then

$$\begin{aligned} |I|^{-1} \int_I e^{v(x)-v_I} dx &= \frac{1}{|I|} \int_{x \in I: v(x)-v_I < 0} e^{v(x)-v_I} dx + \frac{1}{|I|} \int_{x \in I: v(x)-v_I \geq 0} e^{v(x)-v_I} dx \\ &\leq 1 + c_1 \int_0^\infty \exp\left(\lambda - \frac{c_2 \lambda}{\|v\|_{\text{BMO}(\mathbb{R})}}\right) d\lambda \end{aligned}$$

$$\leq 1 + \frac{c_1 \|v\|_{\mathbf{BMO}(\mathbb{R})}}{c_2 - \|v\|_{\mathbf{BMO}(\mathbb{R})}}.$$

Accordingly,

$$\|v\|_{\mathbf{BMO}(\mathbb{R})} < 2^{-1}c_2 \Rightarrow [e^v]_{A_\infty(\mathbb{R})} \leq 1 + 2c_1c_2^{-1}\|v\|_{\mathbf{BMO}(\mathbb{R})}.$$

Letting

$$\epsilon_0 = \min\{1, 2^{-1}c_2\}$$

yields the assertion. \square

Proposition 2.5. *Suppose that $b \in L^1_{\text{loc}}(\mathbb{R})$ has its derivative $b' \in \mathbf{BMO}(\mathbb{R})$. Then b satisfies the Zygmund condition with*

$$|b(x+y) + b(x-y) - 2b(x)| \leq 2|y|\|b'\|_{\mathbf{BMO}(\mathbb{R})} \quad \forall (x, y) \in \mathbb{R} \times \mathbb{R}.$$

Proof. This follows from

$$\begin{aligned} & |b(x+y) + b(x-y) - 2b(x)| \\ &= \left| \int_x^{x+y} b'(z) dz - \int_{x-y}^x b'(z) dz \right| \\ &\leq \left| \int_x^{x+y} b'(z) dz - \frac{1}{2} \int_{x-y}^{x+y} b'(z) dz \right| + \left| \frac{1}{2} \int_{x-y}^{x+y} b'(z) dz - \int_{x-y}^x b'(z) dz \right| \\ &\leq \int_x^{x+y} |b'(z) - b'_{[x-y, x+y]}| dz + \int_{x-y}^x |b'(z) - b'_{[x-y, x+y]}| dz \\ &\leq \int_{x-y}^{x+y} |b'(z) - b'_{[x-y, x+y]}| dz \\ &\leq 2|y|\|b'\|_{\mathbf{BMO}(\mathbb{R})}. \end{aligned}$$

\square

Recall that for a $\mathbf{BMO}(\mathbb{R})$ function f we have

$$\|f\|_* = \|f\|_{\mathbf{BMO}(\mathbb{R})} + \int_{[-1,1]} |f(x)| dx < \infty.$$

In what follows, for a positive constant C , denote by

$$\log^+ C = \max\{1, \log C\}.$$

Proposition 2.6. *Suppose that $b \in L^1_{\text{loc}}(\mathbb{R})$ has its derivative $b' \in \mathbf{BMO}(\mathbb{R})$. Then b satisfies*

$$|b(x) - b(0)| \leq C_5 \|b'\|_* |x| (1 + |\log |x||) \quad \forall x \in \mathbb{R}$$

and

$$|b(x+h) - b(x)| \leq C_5 \|b'\|_* (\log^+ |x|) (|h|(1 + |\log |h||)) \quad \forall (x, h) \in \mathbb{R} \times \mathbb{R}.$$

Proof. From [27, Proposition 5] and Proposition 2.5 it follows that if

$$y \neq 0; z \neq 0; x \in \mathbb{R},$$

then

$$\begin{aligned} & \left| \frac{(y, b(x+y) - b(x))}{|y|^2} - \frac{(z, b(x+z) - b(x))}{|z|^2} \right| \\ & \leq 5\|b'\|_{BMO(\mathbb{R})} + \frac{\|b'\|_{BMO(\mathbb{R})}}{\log 2} \left| \log \frac{|y|}{|z|} \right|. \end{aligned} \quad (7)$$

Letting $x = 0$ and $z = 1$ in (7) gives the first inequality in Proposition 2.6 via

$$\begin{aligned} |b(y) - b(0)| & \leq |y| \left(|b(1) - b(0)| + 5\|b'\|_{BMO(\mathbb{R})} + \frac{\|b'\|_{BMO(\mathbb{R})}}{\log 2} |\log |y|| \right) \\ & \leq C_5 \|b'\|_* |y| (1 + |\log |y||). \end{aligned}$$

Also, by using the structure of $BMO(\mathbb{R})$ (cf. [18, Exercise 7.1.6]) we see that if $x \in \mathbb{R}$ then

$$\begin{aligned} |b(x+1) - b(x)| & = \left| \int_x^{x+1} b'(y) dy - \int_0^1 b'(y) dy \right| + \left| \int_0^1 b'(y) dy \right| \\ & \leq 2(\log^+ |x|) \|b'\|_{BMO(\mathbb{R})} + \left| \int_0^1 b'(y) dy \right| \\ & \leq 2(\log^+ |x|) \|b'\|_*. \end{aligned}$$

This, along with (7), derives the second inequality in Proposition 2.6 via

$$\begin{aligned} |b(x+h) - b(x)| & \leq |h| \left(|b(x+1) - b(x)| + 5\|b'\|_{BMO(\mathbb{R})} + \frac{\|b'\|_{BMO(\mathbb{R})}}{\log 2} |\log |h|| \right) \\ & \leq C_5 \|b'\|_* (\log^+ |x|) |h| (1 + |\log |h||). \end{aligned}$$

□

3. Key a priori estimates for the flow

We say that ϕ is a forward flow associated to b if for each $s \in [0, T]$ and almost every $x \in \mathbb{R}^n$ the map

$$t \mapsto |b(t, \phi_s(t, x))| \text{ belongs to } L^1(s, T)$$

and

$$\phi_s(t, x) = x + \int_s^t b(r, \phi_s(r, x)) dr.$$

If the flow starts at $s = 0$, then we simply denote $\phi_0(t, x)$ by $\phi(t, x)$.

Meanwhile, we say that $\tilde{\phi}$ is a backward flow associated to $b(t, x)$ if for each $t \in [0, T]$ and almost every $x \in \mathbb{R}^n$ the map

$$s \mapsto |b(s, \tilde{\phi}_t(s, x))| \text{ belongs to } L^1(0, t)$$

and

$$\tilde{\phi}_t(s, x) = x - \int_s^t b(r, \tilde{\phi}_t(r, x)) dr.$$

Theorem 3.1. *Let*

$$b(t, x) : [0, T] \times \mathbb{R} \rightarrow \mathbb{R} \text{ be in } L^1(0, T; C^1(\mathbb{R})) \text{ with } \int_0^T \left\| \frac{\partial b(t, x)}{\partial x} \right\|_{L^\infty(\mathbb{R})} dt < \infty.$$

Then there exists a unique flow $\phi(t, x)$ satisfying

$$\begin{cases} \frac{\partial}{\partial t} \phi(t, x) = b(t, \phi(t, x)) & \forall (t, x) \in [0, T] \times \mathbb{R}; \\ \phi_0(x) = x & \forall x \in \mathbb{R}. \end{cases}$$

Moreover, for each $t \in [0, T]$, it holds that

$$\left\| \log \left| \frac{\partial}{\partial x} \phi(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \leq \frac{\int_0^t C_6 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds}{\exp\left(-C_7 \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds\right)}.$$

Proof. The argument is divided into four steps.

Step 1 - initialing argument. Since

$$b(t, x) : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$$

satisfies

$$b \in L^1(0, T; C^1(\mathbb{R})) \text{ with } \frac{\partial b(t, x)}{\partial x} \in L^1(0, T; L^\infty(\mathbb{R})),$$

the classical Cauchy-Lipschitz theory produces a unique flow $\phi_s(t, x)$ with

$$\begin{cases} \frac{\partial}{\partial t} \phi_s(t, x) = b(t, \phi_s(t, x)) & \forall (t, x) \in [s, T] \times \mathbb{R}; \\ \phi_s(s, x) = x & \forall x \in \mathbb{R}. \end{cases}$$

Moreover, for each $t \in [s, T]$, $\phi_s(t, \cdot)$ is a bi-Lipschitz map on \mathbb{R} . Differentiating the equation with respect to the spatial direction, we have

$$\begin{cases} \frac{\partial}{\partial x} \left(\frac{\partial}{\partial t} \phi_s(t, x) \right) = \left(\frac{\partial}{\partial x} b(t, \phi_s(t, x)) \right) \frac{\partial}{\partial x} \phi_s(t, x); \\ \frac{\partial}{\partial t} \log \left| \frac{\partial}{\partial x} \phi_s(t, x) \right| = \frac{\partial}{\partial x} b(t, \phi_s(t, x)). \end{cases}$$

As $\phi_s(t, \cdot)$ is a bi-Lipschitz map on \mathbb{R} for each $t \in [s, T]$, its x -derivative has lower and upper bounds, i.e.,

$$e^{-\int_s^t A(r) dr} \leq \left| \frac{\partial}{\partial x} \phi_s(t, x) \right| \leq e^{\int_s^t A(r) dr},$$

where

$$A(r) = \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{L^\infty(\mathbb{R})}.$$

In particular, this implies that for each t , the function

$$\left| \frac{\partial}{\partial x} \phi_s(t, x) \right|$$

is an $A_\infty(\mathbb{R})$ -weight with

$$\left[\left| \frac{\partial}{\partial x} \phi_s(t, x) \right| \right]_{A_\infty(\mathbb{R})} \leq e^{2 \int_s^t A(r) dr}.$$

Note that the same estimate holds for the backward flow $\tilde{\phi}_t(s, x)$, which is the inverse of $\phi_s(t, x)$.

Upon applying Lemma 2.2, we achieve

$$\begin{aligned} \left\| \log \left| \frac{\partial}{\partial x} \phi_s(t, x) \right| \right\|_{BMO(\mathbb{R})} &= \left\| \int_s^t \frac{\partial}{\partial x} b(r, \phi_s(r, x)) dr \right\|_{BMO(\mathbb{R})} \\ &\leq \int_s^t \left\| \frac{\partial}{\partial x} b(r, \phi_s(r, x)) \right\|_{BMO(\mathbb{R})} dr \\ &\leq \int_s^t C_3 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{BMO(\mathbb{R})} \left[\frac{\partial}{\partial x} \tilde{\phi}_r(s, x) \right]_{A_\infty(\mathbb{R})} dr \\ &\leq \int_s^t C_3 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{BMO(\mathbb{R})} e^{2 \int_s^r A(z) dz} dr. \end{aligned} \quad (8)$$

Step 2 - starting from short time. By letting $T_0 > s \geq 0$ be small enough with

$$\int_s^{T_0} C_3 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{BMO(\mathbb{R})} e^{2 \int_s^r A(z) dz} dr < \epsilon_0,$$

where ϵ_0 is as in Lemma 2.4, we utilize (8) to get

$$\sup_{s \leq t \leq T_0} \left\{ \left\| \log \left| \frac{\partial}{\partial x} \phi_s(t, x) \right| \right\|_{BMO(\mathbb{R})}, \left\| \log \left| \frac{\partial}{\partial x} \tilde{\phi}_t(s, x) \right| \right\|_{BMO(\mathbb{R})} \right\} < \epsilon_0.$$

Hence, by applying Lemma 2.4, we see

$$\left[\frac{\partial}{\partial x} \phi_s(t, x) \right]_{A_\infty(\mathbb{R})} < 1 + C_4 \left\| \log \left| \frac{\partial}{\partial x} \phi_s(t, x) \right| \right\|_{BMO(\mathbb{R})}.$$

Inserting this estimate into (8), we conclude

$$\begin{aligned} \left\| \log \left| \frac{\partial}{\partial x} \phi_s(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} &\leq \int_s^t C_3 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})} \left[\left\| \frac{\partial}{\partial x} \tilde{\phi}_t(s, x) \right\|_{A_\infty(\mathbb{R})} \right] dr \\ &\leq \int_s^t C_3 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})} \left(1 + C_4 \left\| \log \left| \frac{\partial}{\partial x} \tilde{\phi}_r(s, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \right) dr. \end{aligned}$$

Set

$$I_s(t) = \sup_{s \leq r \leq t} \left\{ \left\| \log \left| \frac{\partial}{\partial x} \phi_s(r, \cdot) \right| \right\|_{\text{BMO}(\mathbb{R})}, \left\| \log \left| \frac{\partial}{\partial x} \tilde{\phi}_r(s, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \right\}.$$

The above estimates yield

$$I_s(t) \leq \int_s^t C_3 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})} (1 + C_4 I_s(r)) dr \quad \forall t \in [s, T_0].$$

The Gronwall inequality then implies

$$I_s(t) \leq \frac{\int_s^t C_3 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})} dr}{\exp\left(-\int_s^t C_3 C_4 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})} dr\right)} \quad \forall t \in [s, T_0]. \quad (9)$$

Step 3 - removing the dependence of Lipschitz constant. Let $T_1 \in (s, T]$ obey

$$\int_s^{T_1} C_3 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})} dr \exp\left(C_3 C_4 \int_s^{T_1} \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})} dr\right) \leq 2^{-1} \epsilon_0. \quad (10)$$

We claim that (9) holds for all $t \in (s, T_1]$.

If $T_1 \leq T_0$, then the claim follows from (9).

Suppose now $T_0 < T_1$. Assume that for some $t_0 \in [T_0, T_1)$, (9) holds for all $t \in (s, t_0]$. Then

$$I_s(t_0) \leq \frac{\int_s^{t_0} C_3 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})} dr}{\exp\left(-\int_s^{t_0} C_3 C_4 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})} dr\right)} \leq 2^{-1} \epsilon_0.$$

Since

$$\frac{\partial b(t, x)}{\partial x} \in L^1(0, T; L^\infty(\mathbb{R})),$$

we can choose $t_1 \in (t_0, T_1]$ such that

$$\int_{t_0}^{t_1} C_3 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})} e^{2 \int_{t_0}^r A(z) dz} dr < \epsilon_0 \quad (11)$$

and

$$\frac{\int_{t_0}^{t_1} C_3 \left\| \left| \frac{\partial}{\partial x} b(r, x) \right| \right\|_{BMO(\mathbb{R})} dr}{\exp\left(-\int_{t_0}^{t_1} C_3 C_4 \left\| \left| \frac{\partial}{\partial x} b(r, x) \right| \right\|_{BMO(\mathbb{R})} dr\right)} < \frac{\epsilon_0}{2C_3(1 + C_4 2^{-1} \epsilon_0)}. \quad (12)$$

The same argument as in proving (9) then implies

$$I_{t_0}(t) \leq \frac{\int_{t_0}^t C_3 \left\| \left| \frac{\partial}{\partial x} b(r, x) \right| \right\|_{BMO(\mathbb{R})} dr}{\exp\left(-\int_{t_0}^t C_3 C_4 \left\| \left| \frac{\partial}{\partial x} b(r, x) \right| \right\|_{BMO(\mathbb{R})} dr\right)} < \frac{\epsilon_0}{2C_3(1 + C_4 2^{-1} \epsilon_0)} \quad \forall t \in (t_0, t_1]. \quad (13)$$

For any $t \in (t_0, t_1]$, we have via the semigroup property of the flow that

$$\phi_s(t, x) = \phi_{t_0}(t, \phi_s(t_0, x)).$$

By applying Lemma 2.2, Lemma 2.4 and (13), we find

$$\begin{aligned} & \left\| \log \left| \frac{\partial}{\partial x} \phi_s(t, x) \right| \right\|_{BMO(\mathbb{R})} \\ &= \left\| \log \left| \frac{\partial}{\partial x} \phi_{t_0}(t, \phi_s(t_0, x)) \right| \right\|_{BMO(\mathbb{R})} \\ &\leq \left\| \log \left| \frac{\partial}{\partial z} \phi_{t_0}(t, z) \Big|_{z=\phi_s(t_0, \cdot)} \right| \right\|_{BMO(\mathbb{R})} + \left\| \log \left| \frac{\partial}{\partial x} \phi_s(t_0, x) \right| \right\|_{BMO(\mathbb{R})} \\ &\leq C_3 \left\| \log \left| \frac{\partial}{\partial x} \phi_{t_0}(t, x) \right| \right\|_{BMO(\mathbb{R})} \left[\frac{\partial}{\partial x} \tilde{\phi}_{t_0}(s, x) \right]_{A_\infty(\mathbb{R})} + \left\| \log \left| \frac{\partial}{\partial x} \phi_s(t_0, x) \right| \right\|_{BMO(\mathbb{R})} \\ &< \frac{\epsilon_0 C_3(1 + C_4 2^{-1} \epsilon_0)}{2C_3(1 + C_4 2^{-1} \epsilon_0)} + \frac{\epsilon_0}{2} \\ &= \epsilon_0. \end{aligned}$$

This derives

$$\sup_{s \leq t \leq t_1} \left\{ \left\| \log \left| \frac{\partial}{\partial x} \phi_s(t, x) \right| \right\|_{BMO(\mathbb{R})}, \left\| \log \left| \frac{\partial}{\partial x} \tilde{\phi}_t(s, x) \right| \right\|_{BMO(\mathbb{R})} \right\} < \epsilon_0.$$

Using this estimate in **Step 2**, we further have the following estimate

$$\begin{aligned} & \sup_{s \leq t \leq t_1} \left\{ \left\| \log \left| \frac{\partial}{\partial x} \phi_s(t, x) \right| \right\|_{BMO(\mathbb{R})}, \left\| \log \left| \frac{\partial}{\partial x} \tilde{\phi}_t(s, x) \right| \right\|_{BMO(\mathbb{R})} \right\} \\ &\leq \int_s^{t_1} C_3 \left\| \left| \frac{\partial}{\partial x} b(r, x) \right| \right\|_{BMO(\mathbb{R})} \exp\left(C_3 C_4 \int_s^{t_1} \left\| \left| \frac{\partial}{\partial x} b(z, x) \right| \right\|_{BMO(\mathbb{R})} dz\right) dr < 2^{-1} \epsilon_0, \end{aligned}$$

which implies that (9) holds for all $t \in (s, t_1]$.

Since in (11) and (12) the extension of time only depends on b itself, we may iterate this argument finite times and conclude that (9) holds for all $t \in (s, T_1]$.

Step 4 - completing argument. Since b satisfies

$$\frac{\partial b(t, x)}{\partial x} \in L^1(0, T; L^\infty(\mathbb{R})),$$

we may choose a sequence of increasing numbers $\{T_i\}_{i=1, \dots, k_0}$ such that $T_1 = 0$, $T_{k_0} = T$ and

$$\frac{\int_{T_i}^{T_{i+1}} C_3 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds}{\exp\left(-\int_{T_i}^{T_{i+1}} C_3 C_4 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds\right)} = 2^{-1} \epsilon_0 \quad \forall i \in \{1, \dots, k_0 - 2\},$$

and

$$\frac{\int_{T_{k_0-1}}^{T_{k_0}} C_3 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds}{\exp\left(-\int_{T_{k_0-1}}^{T_{k_0}} C_3 C_4 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds\right)} \leq 2^{-1} \epsilon_0.$$

If $t \in (T_1, T_2]$, then **Step 3** gives

$$\left\| \log \left| \frac{\partial}{\partial x} \phi(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \leq \int_0^t \frac{C_3 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})}}{\exp\left(-C_3 C_4 \int_0^r \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{\text{BMO}(\mathbb{R})} dr\right)} ds. \quad (14)$$

Suppose that t belongs to

$$\text{some } (T_i, T_{i+1}] \text{ with } 2 \leq i \leq k_0 - 1.$$

By using the semigroup property of the flow ϕ , we have

$$\phi(t, x) = \phi_{T_i}(t, \cdot) \circ \phi_{T_{i-1}}(T_i, \cdot) \circ \dots \circ \phi_{T_1}(T_2, x).$$

By using Lemma 2.2, Lemma 2.4 and **Step 3**, we conclude

$$\begin{aligned} & \left\| \log \left| \frac{\partial}{\partial x} \phi(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \\ &= \left\| \log \left| \frac{\partial}{\partial x} \phi_{T_2}(t, \phi_{T_1}(T_2, x)) \right| \right\|_{\text{BMO}(\mathbb{R})} \\ &\leq \left\| \log \left| \frac{\partial}{\partial z} \phi_{T_2}(t, z) \Big|_{z=\phi_{T_1}(T_2, x)} \right| \right\|_{\text{BMO}(\mathbb{R})} + \left\| \log \left| \frac{\partial}{\partial x} \phi_{T_1}(T_2, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \end{aligned}$$

$$\begin{aligned}
&\leq C_3 \left\| \log \left| \frac{\partial}{\partial x} \phi_{T_2}(t, x) \right| \right\|_{BMO(\mathbb{R})} \left[\left\| \frac{\partial}{\partial x} \tilde{\phi}_{T_2}(T_1, x) \right\|_{A_\infty(\mathbb{R})} \right] + \left\| \log \frac{\partial}{\partial x} \phi_{T_1}(T_2, x) \right\|_{BMO(\mathbb{R})} \\
&\leq C_3(1 + C_4 2^{-1} \epsilon_0) \left\| \log \left| \frac{\partial}{\partial x} \phi_{T_2}(t, x) \right| \right\|_{BMO(\mathbb{R})} + 2^{-1} \epsilon_0 \\
&\leq C_3(1 + C_4) \left\| \log \left| \frac{\partial}{\partial x} \phi_{T_2}(t, x) \right| \right\|_{BMO(\mathbb{R})} + 1 \\
&\leq (C_3(1 + C_4))^2 \left\| \log \left| \frac{\partial}{\partial x} \phi_{T_3}(t, x) \right| \right\|_{BMO(\mathbb{R})} + C_3(1 + C_4) + 1 \\
&\leq \dots \\
&\leq (C_3(1 + C_4))^{i-1} \left\| \log \left| \frac{\partial}{\partial x} \phi_{T_i}(t, x) \right| \right\|_{BMO(\mathbb{R})} + \sum_{j=0}^{i-2} (C_3(1 + C_4))^j \\
&\leq (C_3(1 + C_4) + 1)^i.
\end{aligned}$$

Let $\delta_0 > 0$ obey

$$C_3 \delta_0 e^{C_3 C_4 \delta_0} = 2^{-1} \epsilon_0.$$

As

$$\begin{cases} \epsilon_0 \leq 1; \\ \delta_0 < 1; \\ t \in (T_i, T_{i+1}], \end{cases}$$

by our choice of $\{T_i\}$ we find

$$(i-1)\delta_0 < \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds \leq i\delta_0,$$

whence

$$\left\| \log \left| \frac{\partial}{\partial x} \phi(t, x) \right| \right\|_{BMO(\mathbb{R})} \leq \frac{\frac{1}{\delta_0} \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds}{\exp\left(-C \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds\right)}.$$

This, together with (14), implies

$$\left\| \log \left| \frac{\partial}{\partial x} \phi(t, x) \right| \right\|_{BMO(\mathbb{R})} \leq \frac{\int_0^t \frac{C_3}{\delta_0} \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{BMO(\mathbb{R})} ds}{\exp\left(-C \int_0^t \left\| \frac{\partial}{\partial x} b(r, x) \right\|_{BMO(\mathbb{R})} dr\right)} ds,$$

as desired. \square

Rather surprisingly, the hypothesis

$$\int_0^T \left\| \frac{\partial b(t, x)}{\partial x} \right\|_{L^\infty(\mathbb{R})} dt < \infty$$

in Theorem 3.1 can be replaced by a weaker one

$$\int_0^T \left\| \frac{\partial b(t, x)}{\partial x} \right\|_* dt < \infty$$

in the following assertion.

Theorem 3.2. *Let*

$$b(t, x) : [0, T] \times \mathbb{R} \rightarrow \mathbb{R} \text{ be in } L^1(0, T; C^1(\mathbb{R})) \text{ with } \int_0^T \left\| \frac{\partial b(t, x)}{\partial x} \right\|_* dt < \infty.$$

Then there exists a unique flow $\phi(t, x)$ satisfying

$$\begin{cases} \frac{\partial}{\partial t} \phi(t, x) = b(t, \phi(t, x)) & \forall (t, x) \in [0, T] \times \mathbb{R}; \\ \phi_0(x) = x & \forall x \in \mathbb{R}. \end{cases}$$

Moreover

$$\left\| \log \left| \frac{\partial}{\partial x} \phi(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \leq \frac{\int_0^t 2C_6 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds}{\exp\left(-2C_7 \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds\right)} \quad \forall t \in [0, T].$$

Proof. The existence and uniqueness has essentially been established in [27]. So it remains to verify the last $\text{BMO}(\mathbb{R})$ -size estimate.

For each $(k, t) \in \mathbb{N} \times [0, T]$ set

$$\begin{cases} v_k(t, x) = \min \left\{ \max \left\{ -k, \frac{\partial}{\partial x} b(t, x) \right\}, k \right\}; \\ b_k(t, x) = b(t, 0) + \int_0^x v_k(t, y) dy. \end{cases}$$

Then

$$\begin{cases} \frac{\partial}{\partial x} b_k(t, x) \in L^1(0, T; L^\infty(\mathbb{R})); \\ \|v_k(t, \cdot)\|_{\text{BMO}(\mathbb{R})} \leq 2\|\partial_x b(t, \cdot)\|_{\text{BMO}(\mathbb{R})}; \\ \|v_k(t, \cdot)\|_* \leq 2\|\partial_x b(t, \cdot)\|_*. \end{cases} \quad (15)$$

In accordance with Propositions 2.5-2.6, we see that $\{b_k\}$ and b satisfy the Zygmund condition with a uniform constant.

Let $\{\phi_k, \phi\}$ be the unique flow pair generated by $\{b_k(t, x), b(t, x)\}$. Then by [27, Proposition 4], we see that $\phi(t, \cdot)$ and $\phi_k(t, \cdot)$ are locally Hölder continuous

on \mathbb{R} for each $t \in [0, T]$. Moreover for each compact set $K \subset \mathbb{R}$, both $\phi(t, \cdot)$ and $\phi_k(t, \cdot)$ are Hölder continuous on K for each $t \in [0, T]$ with the Hölder exponent and constant depending only on

$$\int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_* ds.$$

On the other hand, by the construction of b_k and Proposition 2.6 we have

$$|b_k(t, x) - b(t, 0)| \leq C_5 \|v_k(t, \cdot)\|_* |x| (1 + |\log |x||) \leq 2C_5 \|\partial_x b(t, \cdot)\|_* |x| (1 + |\log |x||),$$

thereby getting that

$$\{|\phi_k(t, x)| : (t, x) \in [0, T] \times K\}$$

is uniformly bounded. Denote by

$$C_8(K) := \sup \{|\phi_k(t, x)| + |\phi(t, x)| : (t, x, k) \in [0, T] \times K \times \mathbb{N}\}.$$

Then it holds for each $x \in K$ and all $0 \leq s < t \leq T$ that

$$\begin{aligned} |\phi_k(t, x) - \phi_k(s, x)| &\leq \int_s^t |b_k(r, \phi_k(r, x))| dr \\ &\leq \int_s^t \left(|b(r, 0)| + 2C_5 \left\| \frac{\partial}{\partial x} b(r, x) \right\|_* C_8(K) \left(1 + |\log |C_8(K)||\right) \right) dr. \end{aligned}$$

This, together with the previous discussion on the Hölder continuity in the spatial direction, implies that $\{\phi_k\}_k$ are equicontinuous on $[0, T] \times K$. Applying the Arzelá-Ascoli theorem, we conclude that there is a subsequence of $\{\phi_k\}_k$, denoted by $\{\phi_{K,k}\}_k$, such that $\phi_{K,k}$ converges uniformly on $[0, T] \times K$.

By construction we have

$$b_k(t, x) \rightarrow b(t, x) \text{ as } k \rightarrow \infty,$$

thereby concluding that if $(t, x) \in [0, T] \times K$ then

$$\begin{aligned} \lim_{k \rightarrow \infty} \phi_{K,k}(t, x) &= x + \lim_{k \rightarrow \infty} \int_0^t b_{K,k}(s, \phi_{K,k}(s, x)) ds \\ &= x + \lim_{k \rightarrow \infty} \int_0^t \int_0^{\phi_{K,k}(s, x)} \left[v_{K,k}(s, y) - \frac{\partial}{\partial x} b(s, y) \right] dy ds \\ &\quad + \lim_{k \rightarrow \infty} \int_0^t b(s, \phi_{K,k}(s, x)) ds. \end{aligned}$$

Since

$$|\phi_k(s, x)| \leq C_8(K),$$

one has

$$\left| \int_0^t \int_0^{\phi_{K,k}(s,x)} [v_{K,k}(s,y) - \partial_x b(s,y)] dy ds \right| \leq \int_0^T \int_{-C_8(K)}^{C_8(K)} \left| \frac{\partial}{\partial x} b(s,x) \right| dx ds < \infty,$$

and hence the dominated convergence theorem and continuity of $b(t, \cdot)$ guarantee

$$\lim_{k \rightarrow \infty} \phi_{K,k}(t, x) = x + \int_0^t b(s, \lim_{k \rightarrow \infty} \phi_{K,k}(s, x)) ds.$$

By choosing a sequence of increasing compacts K_j such that $\mathbb{R} = \cup_j K_j$ and passing to further subsequences, we see that there is a subsequence of $\{\phi_k\}$, still denoted by $\{\phi_{K,k}\}$, such that $\phi_{K,k}(t, x)$ converges on $[0, T] \times \mathbb{R}$, and uniformly on any compact subset $[0, T] \times \tilde{K}$, and consequently,

$$\lim_{k \rightarrow \infty} \phi_{K,k}(t, x) = x + \int_0^t b(s, \lim_{k \rightarrow \infty} \phi_{K,k}(s, x)) ds \quad \forall (t, x) \in [0, T] \times \mathbb{R}.$$

By the uniqueness, we see that

$$\phi(t, x) = \lim_{k \rightarrow \infty} \phi_{K,k}(t, x), \quad \forall (t, x) \in [0, T] \times \mathbb{R},$$

and the convergence is uniform on any compact set.

Since

$$b(t, x) \in L^1(0, T; C^1(\mathbb{R})),$$

and so is any $b_k(t, x)$. Accordingly, the proof of Theorem 3.1 yields that if $(t, x) \in [0, T] \times \mathbb{R}$ then

$$\begin{aligned} \log \left| \frac{\partial}{\partial x} \phi(t, x) \right| &= \int_0^t \frac{\partial}{\partial x} b(s, \phi(s, x)) ds \\ &= \int_0^t \lim_{k \rightarrow \infty} v_k(s, \phi_k(s, x)) ds \\ &= \lim_{k \rightarrow \infty} \log \left| \frac{\partial}{\partial x} \phi_k(t, x) \right|. \end{aligned}$$

By (15) and Theorem 3.1, we see that for each $k \in \mathbb{N}$,

$$\left\| \log \left| \frac{\partial}{\partial x} \phi_k(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \leq \frac{\int_0^t 2C_6 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds}{\exp \left(-2C_7 \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds \right)}.$$

By this, the weak-* compactness in $\text{BMO}(\mathbb{R})$, and the pointwise convergence of

$$\frac{\partial}{\partial x} \phi_k(t, x),$$

we conclude that the last estimation holds also for

$$\log \left| \frac{\partial}{\partial x} \phi(t, x) \right|,$$

thereby completing the proof. \square

4. Verification of the main results

Proof of Theorem 1.1. The argument consists of three steps.

Step 1 - an Orlicz space estimate. Let μ denote the Gaussian measure on \mathbb{R} , i.e.,

$$\mu(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{|x|^2}{2}\right),$$

and $\operatorname{div}_\mu b$ denotes the distributional divergence of b with respect to μ . We say that a measurable function

$$f \in \operatorname{Exp}_\mu\left(\frac{L}{\log L}\right)$$

provided

$$\|f\|_{\operatorname{Exp}_\mu\left(\frac{L}{\log L}\right)} = \inf \left\{ \lambda > 0 : \int_{\mathbb{R}} \left[\exp\left(\frac{|f(x)|/\lambda}{1 + \log^+(|f(x)|/\lambda)}\right) - 1 \right] d\mu \leq 1 \right\}.$$

Let $b(t, x)$ obey (1). Then

$$\begin{cases} \frac{b(t, x)}{1 + |x| \log^+ |x|} \in L^1(0, T; L^\infty(\mathbb{R})); \\ \operatorname{div}_\mu b(t, x) \in L^1(0, T; \operatorname{Exp}_\mu\left(\frac{L}{\log L}\right)). \end{cases} \quad (16)$$

As a matter of fact, the first estimate of (16) follows from Proposition 2.6 as

$$\frac{|b(t, x)|}{1 + |x| \log^+ |x|} \leq \frac{|b(t, x) - b(t, 0) + b(t, 0)|}{1 + |x| \log^+ |x|} \leq |b(t, 0)| + C \left\| \frac{\partial}{\partial x} b(t, x) \right\|_*.$$

To verify the second relation in (16), set

$$\beta(t) = |b(t, 0)| + C \left\| \frac{\partial}{\partial x} b(t, x) \right\|_{BMO(\mathbb{R})}.$$

Noting that

$$\begin{aligned} & \int_{\mathbb{R}} \exp\left(\frac{c|xb(t, x)|}{1 + \log^+(c|xb(t, x)|)}\right) d\mu(x) \\ & \leq \int_{\mathbb{R}} \exp\left(\frac{c|x|(1 + |x| \log^+ |x|)\beta(t)}{1 + \log^+(c|x|(1 + |x| \log^+ |x|)\beta(t))}\right) d\mu(x), \end{aligned}$$

we obtain

$$\|xb(t, x)\|_{\operatorname{Exp}_\mu\left(\frac{L}{\log L}\right)} \leq C\beta(t).$$

On the other hand, for a $\text{BMO}(\mathbb{R})$ -function f , we utilize the John-Nirenberg inequality:

$$\{|x \in I : |f(x) - f_I| > \lambda|\} \leq c_1 |I| \exp\left(-\frac{c_2 \lambda}{\|f\|_{\text{BMO}(\mathbb{R})}}\right), \quad \forall \text{ interval } I \subset \mathbb{R}$$

to obtain that if

$$\begin{cases} I = [x - r, x + 1]; \\ (x, r) \in \mathbb{R} \times [1, \infty); \\ \gamma(t) = \left\| \frac{\partial}{\partial x} b(t, x) \right\|_*; \\ \alpha = c_2 (2\gamma(t))^{-1}, \end{cases}$$

then

$$|f_I| \leq |f_I - f_{[-1,1]}| + |f_{[-1,1]}| \leq C(1 + \log^+ |x|) \|f\|_*,$$

and hence

$$\begin{aligned} & \int_{\mathbb{R}} \exp\left(\alpha \left| \frac{\partial}{\partial x} b(t, x) \right| \right) d\mu(x) \\ & \leq \int_{-1}^1 \exp\left(\alpha \left| \frac{\partial}{\partial x} b(t, x) \right| \right) d\mu(x) + \sum_{k=1}^{\infty} \left(\int_{2^{k-1}}^{2^k} + \int_{-2^k}^{-2^{k-1}} \right) \exp\left(\alpha \left| \frac{\partial}{\partial x} b(t, x) \right| \right) d\mu(x) \\ & \leq e^{2\alpha\gamma(t)} \sum_{k=0}^{\infty} \alpha 2^k e^{-2^{2k-1} + ck} \left(\frac{\alpha\gamma(t)}{c_2 - \alpha\gamma(t)} \right) \leq C. \end{aligned}$$

Consequently we achieve the desired inequality

$$\left\| \frac{\partial}{\partial x} b(t, x) \right\|_{\text{Exp}_{\mu}\left(\frac{L}{\log L}\right)} \leq \left\| \frac{\partial}{\partial x} b(t, x) \right\|_{\text{Exp}_{\mu}(L)} \leq C \left\| \frac{\partial}{\partial x} b(t, x) \right\|_*.$$

Step 2 - existence-uniqueness-size of flow. Under (1) we conclude via Proposition 2.5 for a.e. t , that b is in the Zygmund class, which implies that the flow exists and is unique; see [27] for instance.

Moreover, from **Step 1** above it follows that b satisfies requirements from [9, Main Theorem] and so that $\phi(t, x)$ is absolutely continuous and differentiable. Indeed, by using [9, Theorem 1.2] and that $b(t, \cdot)$ is in the Zygmund class, one can deduce that

$$\left| \frac{\partial}{\partial x} \phi(t, x) \right| \left(1 + \log^+ \left| \frac{\partial}{\partial x} \phi(t, x) \right| \right)^q \in L^1_{\text{loc}}(\mathbb{R})$$

for any $q \in [1, \infty)$. As $\partial_x b(t, x) \in \text{BMO}(\mathbb{R})$ is locally exponentially integrable, we deduce that

$$\frac{\partial}{\partial t} \left(\frac{\partial}{\partial x} \phi(t, x) \right) = \left(\frac{\partial}{\partial z} b(t, z) \Big|_{z=\phi(t,x)} \right) \frac{\partial}{\partial x} \phi(t, x)$$

and

$$\log \left| \frac{\partial}{\partial x} \phi(t, x) \right| = \int_0^t \frac{\partial}{\partial x} b(s, \phi(s, x)) ds. \quad (17)$$

For $\epsilon > 0$ and $x \in \mathbb{R}$ set

$$\begin{cases} 0 \leq \rho \in C_c^\infty(\mathbb{R}); \\ \text{supp } \rho \subset (-1, 1); \\ \int_{\mathbb{R}} \rho(x) dx = 1; \\ \rho_\epsilon(x) = \frac{1}{\epsilon} \rho\left(\frac{x}{\epsilon}\right); \\ b_\epsilon(t, x) = b(t, \cdot) * \rho_\epsilon(x). \end{cases}$$

Note that

$$\begin{aligned} \frac{\partial}{\partial x} b(t, x) &\in L^1(0, T; \text{BMO}(\mathbb{R})) \\ \Rightarrow \frac{\partial}{\partial x} b_\epsilon(t, x) &\in L^1(0, T; \text{BMO}(\mathbb{R})) \cap L^1(0, T; C^\infty(\mathbb{R})). \end{aligned}$$

Thus we have

$$\int_0^t \left\| \frac{\partial}{\partial x} b_\epsilon(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds \leq \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds, \quad \forall t \in (0, T]$$

and so for any $\epsilon \in (0, 1)$

$$\left\| \frac{\partial}{\partial x} b_\epsilon(t, x) \right\|_* \leq 2 \left\| \frac{\partial}{\partial x} b(t, x) \right\|_* \quad \text{for a.e. } t \in (0, T].$$

Let $\phi_\epsilon(t, x)$ be the flow generated by b_ϵ , i.e.,

$$\frac{\partial}{\partial t} \phi_\epsilon(t, x) = b_\epsilon(t, \phi_\epsilon(t, x)).$$

Then Theorem 3.2 is utilized to imply

$$\begin{aligned} \left\| \log \left| \frac{\partial}{\partial x} \phi_\epsilon(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} &\leq \frac{\int_0^t 2C_6 \left\| \frac{\partial}{\partial x} b_\epsilon(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds}{\exp\left(-2C_7 \int_0^t \left\| \frac{\partial}{\partial x} b_\epsilon(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds\right)} \\ &\leq \frac{\int_0^t 2C_6 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds}{\exp\left(-2C_7 \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds\right)}, \quad \forall \epsilon > 0. \end{aligned}$$

The proof of [9, Main Theorem] infers that, up to a subsequence $\{\epsilon_k\}_{k \in \mathbb{N}}$,

$$\lim_{k \rightarrow \infty} \phi_{\epsilon_k}(t, x) = \phi(t, x), \quad \forall t \in (0, T].$$

From this, (17) and the weak-* compactness in $\text{BMO}(\mathbb{R})$, we conclude that $\frac{\partial}{\partial x} \phi(t, x)$ is the weak-* limit of $\frac{\partial}{\partial x} \phi_{\epsilon_k}(t, x)$ for each $t \in (0, T]$. This implies

$$\left\| \log \left| \frac{\partial}{\partial x} \phi(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \leq \frac{\int_0^t 2C_6 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds}{\exp\left(-2C_7 \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds\right)},$$

namely, the size estimate (2) holds.

Step 3 - $A_\infty(\mathbb{R})$ density of flow. It remains to show that for each $t \in [0, T]$,

$$\left| \frac{\partial}{\partial x} \phi(t, x) \right|$$

is an $A_\infty(\mathbb{R})$ -weight. But, from Theorem 1.2 (to be proved later on), we see that

$$u_0 \in \text{BMO}(\mathbb{R}) \Rightarrow u_0 \circ \phi(t, \cdot) \in \text{BMO}(\mathbb{R}) \quad \forall t \in (0, T].$$

Then we apply [21, Theorem] to conclude that for each $t \in [0, T]$,

$$\left| \frac{\partial}{\partial x} \phi(t, x) \right|$$

is an $A_\infty(\mathbb{R})$ -weight. □

Proof of Theorem 1.2. The argument consists of three steps.

Step 1 - existence of solution. Let ϕ be the flow generated by b , i.e.,

$$\begin{cases} \frac{\partial}{\partial t} \phi(t, x) = b(t, \phi(t, x)) & \forall (t, x) \in (0, T] \times \mathbb{R}; \\ \phi_0(x) = x & \forall x \in \mathbb{R}. \end{cases}$$

Then the same proof of [10, Theorem 1] derives that $u_0 \circ \phi$ is a solution to the transport equation.

Step 2 - size of solution. Let ϵ_0 be the same as in Lemma 2.4, and

$$\delta_0 > 0 \quad \& \quad 2C_6 \delta_0 e^{2C_7 \delta_0} = 2^{-1} \epsilon_0.$$

We choose a sequence of increasing numbers

$$0 = T_0 < T_1 < \cdots < T_{k_0} = T$$

such that

$$\frac{\int_{T_{i-1}}^{T_i} 2C_6 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds}{\exp\left(-\int_{T_{i-1}}^{T_i} 2C_7 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds\right)} = 2^{-1} \epsilon_0 \quad \forall i \in \{1, \dots, k_0 - 1\},$$

and

$$\frac{\int_{T_{k_0-1}}^{T_{k_0}} 2C_6 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds}{\exp\left(-\int_{T_{k_0-1}}^{T_{k_0}} 2C_7 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds\right)} \leq 2^{-1} \epsilon_0.$$

Suppose that t belongs to

some interval $(T_i, T_{i+1}]$ where $i = 0, \dots, k_0 - 1$.

If $i = 0$, then by Lemma 2.2 and Lemma 2.4, we obtain

$$\begin{aligned} \|u(t, x)\|_{BMO(\mathbb{R})} &\leq C_3 \|u_0\|_{BMO(\mathbb{R})} \left(1 + C_4 \left\| \log \left| \frac{\partial}{\partial x} \tilde{\phi}_t(0, x) \right| \right\|_{BMO(\mathbb{R})}\right) \\ &\leq C_3 \|u_0\|_{BMO(\mathbb{R})} \left(1 + \frac{2C_4 C_6 \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds}{\exp\left(-\int_0^t 2C_7 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds\right)}\right) \\ &\leq C_3 \|u_0\|_{BMO(\mathbb{R})} \exp\left(\int_0^t C \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds\right). \end{aligned} \quad (18)$$

Suppose next $i \geq 1$. By the semigroup property of the flow, we may write

$$u(t, x) = u_0 \circ \phi_{T_i}(t, \cdot) \circ \dots \circ \phi_{T_0}(T_1, x).$$

By Theorem 1.1, for all $t \in (T_i, T_{i+1}]$ we have

$$\left\| \log \left| \frac{\partial}{\partial x} \tilde{\phi}_t(T_i, x) \right| \right\|_{BMO(\mathbb{R})} \leq \frac{\int_{T_i}^t 2C_6 \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds}{\exp\left(-2C_7 \int_{T_i}^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{BMO(\mathbb{R})} ds\right)} \leq 2^{-1} \epsilon_0. \quad (19)$$

A combination of (19) and Lemma 2.4 derives

$$\left\{ \begin{array}{l} \left| \frac{\partial}{\partial x} \tilde{\phi}_t(T_i, x) \right| \in A_\infty(\mathbb{R}); \\ \left\| \left| \frac{\partial}{\partial x} \tilde{\phi}_t(T_i, x) \right| \right\|_{A_\infty(\mathbb{R})} \leq 1 + C_4 \left\| \log \left| \frac{\partial}{\partial x} \tilde{\phi}_t(T_i, x) \right| \right\|_{BMO(\mathbb{R})}. \end{array} \right.$$

Then Lemma 2.2 implies for all $v \in \text{BMO}(\mathbb{R})$ that

$$\|v \circ \phi_{T_i}(t, x)\|_{\text{BMO}(\mathbb{R})} \leq C_3 \|v\|_{\text{BMO}(\mathbb{R})} \left(1 + C_4 \left\| \log \left| \frac{\partial}{\partial x} \tilde{\phi}_{T_i}(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \right).$$

Upon repeating this argument for i times more, we gain

$$\begin{aligned} \|u(t, x)\|_{\text{BMO}(\mathbb{R})} &= \|u_0 \circ \phi_{T_i}(t, x) \circ \cdots \circ \phi_{T_0}(T_1, x)\|_{\text{BMO}(\mathbb{R})} \\ &\leq C_3^{i+1} \|u_0\|_{\text{BMO}(\mathbb{R})} \frac{\prod_{j=1}^i \left(1 + C_4 \left\| \log \left| \frac{\partial}{\partial x} \tilde{\phi}_{T_j}(T_{j-1}, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \right)}{\left(1 + C_4 \left\| \log \left| \frac{\partial}{\partial x} \tilde{\phi}_{T_i}(t, x) \right| \right\|_{\text{BMO}(\mathbb{R})} \right)^{-1}} \\ &\leq C_3^{i+1} \left(1 + C_4 2^{-1} \epsilon_0 \right)^{i+1} \|u_0\|_{\text{BMO}(\mathbb{R})} \\ &\leq \|u_0\|_{\text{BMO}(\mathbb{R})} \exp \left(C \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds \right), \end{aligned}$$

where in the last inequality we have used

$$i\delta_0 < \int_0^t \left\| \frac{\partial}{\partial x} b(s, x) \right\|_{\text{BMO}(\mathbb{R})} ds \leq (i+1)\delta_0.$$

This, together with (18), gives the desired size estimate.

Step 3 - uniqueness of solution. This follows easily as an application of the renormalized property of solutions established by DiPerna-Lions [13] and the well-posedness of solutions in $L^\infty(0, T; L^\infty(\mathbb{R}))$ established in [8]; see the proof of [10, Theorem 1] for instance. \square

Acknowledgement. This work is a continuation of R. Jiang's project joint with A. Clop, J. Mateu and J. Orobitg at Department of Mathematics, Autonomous University of Barcelona and R. Jiang would like to thank the department for its warm hospitality. In addition, R. Jiang was supported in part by National Natural Science Foundation of China (11671039, 11771043 & 11922114); K. Li was supported by Juan de la Cierva - Formaci3n 2015 FJCI-2015-24547, by the Basque Government via the BERC 2018-2021 program and by Spanish Ministry of Economy and Competitiveness MINECO through both BCAM Severo Ochoa excellence accreditation SEV-2013-0323 and project MTM2017-82160-C2-1-P funded by (AEI/FEDER, UE) and acronym "HAQMEC"; J. Xiao was supported by NSERC of Canada (#20171864).

References

- [1] T. Alberico, R. Corporente and C. Sbordone, ‘Explicit bounds for composition operators preserving $BMO(\mathbb{R})$ ’, *Georgian Math. J.* **14** (2007), 21-32.
- [2] L. Ambrosio, ‘Transport equation and Cauchy problem for BV vector fields’, *Invent. Math.* **158** (2004), 227-260.
- [3] A. Beurling and L. Ahlfors, ‘The boundary correspondence under quasiconformal mappings’, *Acta Math.* **96** (1956), 125-142.
- [4] S. Bloom, ‘Sharp weights and BMO-preserving homeomorphisms’, *Studia Math.* **96** (1990), 1-10.
- [5] M. Bonk, J. Heinonen and E. Saksman, ‘Logarithmic potentials, quasiconformal flows, and Q-curvature’, *Duke Math. J.* **142** (2008), 197-239.
- [6] F. Bouchut and F. James, ‘Duality solutions for pressureless gases, monotone scalar conservation laws, and uniqueness’, *Comm. Partial Differential Equations* **24** (1999), 2173-2189.
- [7] D. Chung, C. Pereyra and C. Pérez, ‘Sharp bounds for general commutators on weighted Lebesgue spaces’, *Trans. Amer. Math. Soc.* **364** (2012), 1163-1177.
- [8] A. Clop, R. Jiang, J. Mateu and J. Orobitg, ‘Linear transport equations for vector fields with subexponentially integrable divergence’, *Calc. Var. Partial Differential Equations* **55** (2016), Art. 21, 30 pp.
- [9] A. Clop, R. Jiang, J. Mateu and J. Orobitg, ‘Flows for non-smooth vector fields with subexponentially integrable divergence’, *J. Differential Equations* **261** (2016), 1237-1263.
- [10] A. Clop, R. Jiang, J. Mateu and J. Orobitg, ‘A note on transport equations in quasiconformally invariant spaces’, *Adv. Calc. Var.* **11** (2018), 193-202.
- [11] A. Clop, H. Jylhä, J. Mateu and J. Orobitg, ‘Well-posedness for the continuity equation for vector fields with suitable modulus of continuity’, *J. Funct. Anal.* **276** (2019), 45-77.
- [12] G. Crippa and C. De Lellis, ‘Estimates and regularity results for the DiPerna-Lions flow’, *J. Reine Angew. Math.* **616** (2008), 15-46.
- [13] R. J. DiPerna and P. L. Lions, ‘Ordinary differential equations, transport theory and Sobolev spaces’, *Invent. Math.* **98** (1989), 511-547.
- [14] J.J. Donaire, J.G. Llorente and A. Nicolau, ‘Differentiability of functions in the Zygmund class’, *Proc. Lond. Math. Soc.* (3) **108** (2014), 133-158.
- [15] Y. Fan, Y. Hu and Y. Shen, ‘A note on BMO map induced by strongly quasisymmetric homeomorphism’, *Proc. Amer. Math. Soc.* **145** (2017), 2505-2512.
- [16] C. Fefferman and E. M. Stein, ‘ H^p spaces of several variables’, *Acta Math.* **129** (1972), 137-193
- [17] M. A. Fominykh, ‘Admissible changes of variables in the class of BMO functions’, *Math. Notes* **43** (1988), 366-371.
- [18] L. Grafakos, *Modern Fourier analysis*. Second edition. Graduate Texts in Mathematics, 250. Springer, New York, 2009. xvi+504pp.
- [19] T. Hytönen and C. Pérez, ‘Sharp weighted bounds involving A_∞ ’, *Anal. PDE* **6** (2013), 777-818.
- [20] R. Johnson and C.J. Neugebauer, ‘Homeomorphisms preserving $A_p(\mathbb{R})$ ’, *Rev. Mat. Iberoica.* **3** (1987), 249-273.
- [21] P. Jones, ‘Homeomorphisms of the line which preserve BMO’, *Ark. Mat.* **21** (1983), 229-231.
- [22] H. Koch, P. Koskela, E. Saksman and T. Soto, ‘Bounded compositions on scaling invariant Besov spaces’, *J. Funct. Anal.* **266** (2014), 2765-2788.
- [23] P. Koskela, D. Yang and Y. Zhou, ‘Pointwise characterizations of Besov and Triebel-

- Lizorkin spaces and quasiconformal mappings', *Adv. Math.* **226** (2011), 3579-3621.
- [24] K. Li, S. Ombrosi and C. Pérez, 'Proof of an extension of E. Sawyer's conjecture about weighted mixed weak-type estimates', *Math. Ann.* **374** (2019), 907-929.
- [25] P. B. Mucha, 'Transport equation: extension of classical results for $\operatorname{div} b \in BMO$ ', *J. Differential Equations* **249** (2010), 1871-1883.
- [26] H. M. Reimann, 'Functions of bounded mean oscillation and quasiconformal mappings', *Comment. Math. Helv.* **49** (1974), 260-276.
- [27] H. M. Reimann, 'Ordinary differential equations and quasiconformal mappings', *Invent. Math.* **33** (1976), 247-270.
- [28] S.K. Vodop'yanov, 'Mappings of homogeneous groups and imbeddings of functional spaces', *Siberian Math. Zh.* **30** (1989), 25-41.
- [29] J. Xiao, 'The transport equation in scaling invariant Besov or Essén-Janson-Peng-Xiao space', *J. Differential Equations* **266** (2019), 7124-7151.
- [30] D. Yang, W. Yuan and Y. Zhou, 'Sharp boundedness of quasiconformal composition operators on Triebel-Lizorkin type spaces', *J. Geom. Anal.* **27** (2017), 1548-1588.