

Global stability for a discrete epidemic model for disease with immunity and latency spreading in a heterogeneous host population

Yoshiaki Muroya^{a,*}, Alfredo Bellen^b, Yoichi Enatsu^c, Yukihiro Nakata^d

^a*Department of Mathematics, Waseda University 3-4-1 Ohkubo, Shinjuku-ku, Tokyo 169-8555, Japan*

^b*Dipartimento di Matematica e Informatica, University of Trieste, Italy*

^c*Department of Pure and Applied Mathematics, Waseda University, 3-4-1 Ohkubo, Shinjuku-ku, Tokyo, 169-8555, Japan*

^d*Basque Center for Applied Mathematics, Bizkaia Technology Park, Building 500 E-48160 Derio, Spain*

Abstract

In this paper, we propose a discrete epidemic model for disease with immunity and latency spreading in a heterogeneous host population which is derived from the continuous case by using the well-known backward Euler method and applying a Lyapunov functional technique which is a discrete version to that in the paper [Prüss, Pajo-Menjouet, Webb and Zacher, Analysis of a model for the dynamics of prions, *Discrete and Continuous Dynamical Systems-Series B* **6** (2006), 225-235]. It is shown that the global dynamics of this discrete epidemic model with latency are fully determined by a single threshold parameter.

Keywords: Epidemic model; latency; heterogeneous host; permanence; global asymptotic stability; Lyapunov functional;

1. Introduction

The application of theories of functional differential/difference equations in mathematical biology has been developed rapidly. Various mathematical models have been proposed in the literature of population dynamics, ecology and

*Corresponding author

Email addresses: ymuroya@waseda.jp (Yoshiaki Muroya), bellen@units.it (Alfredo Bellen), yo1.gc-rw.docomo@akane.waseda.jp (Yoichi Enatsu), nakata@bcamath.org (Yukihiro Nakata)

^aResearch was supported by Scientific Research (c), No.21540230 of Japan Society for the Promotion of Science.

^cResearch was supported by JSPS Fellows, No.237213 of Japan Society for the Promotion of Science.

^dResearch was supported by the Grant MTM2010-18318 of the MICINN, Spanish Ministry of Science and Innovation.

epidemiology. Many authors have studied the dynamical behavior of several epidemic models (see [1]-[15] and the references therein).

Consider the following integro-differential system:

$$\begin{cases} S'(t) = B - \delta S(t) - \beta S(t)I(t) + \sigma I(t), \\ I'(t) = \beta \int_0^t S(u)I(u)g(t-u)e^{-\delta(t-u)}du - (\delta + \epsilon + \gamma + \sigma)I(t), \\ R'(t) = \gamma I(t) - \delta R(t), \end{cases} \quad (1.1)$$

where $S(t)$, $I(t)$ and $R(t)$ denote the numbers of susceptible, infectious, and recovered of individuals at time t , respectively. The non-negative constant β is the transmission rate due to the contact of susceptible individuals with infectious individuals. The non-negative constants δ , ϵ , γ and σ are natural death rates, disease-caused death rates, recovery rates and immigration rates of infectives, respectively. The function $g(t)$ is the probability density function for the time (a random variable) it takes for an infected individual to become infectious and we choose the gamma distribution:

$$g(u) = g_{n,b}(u) \equiv \frac{u^{n-1}}{(n-1)!b^n} e^{-u/b}, \quad (1.2)$$

where $b > 0$ is a real number and $n > 1$ is an integer.

By using “linear chain trick” to transfer from this model (see Yuan and Zou [15]), we can derive the following system of ordinary differential equations for a disease not only with a latent period and but also with an immigration of infectives:

$$\begin{cases} S'(t) = B - \delta S(t) - \beta S(t)I(t) + \sigma I(t), \\ y_1'(t) = c(S(t))y_{n+1}(t) - dy_1(t), \\ y_j'(t) = dy_1(t) - dy_j(t), \quad j = 2, 3, \dots, n, \\ y_{n+1}'(t) = dy_n(t) - (e + \sigma)y_{n+1}(t), \end{cases} \quad (1.3)$$

where

$$\begin{cases} c(S) = \frac{\beta S}{(1+\delta b)^n}, \quad d = \frac{1}{b}, \quad e = \delta + \epsilon + \gamma, \\ \hat{b} = \frac{b}{1+\delta b}, \quad y_{n+1}(t) = I(t). \end{cases} \quad (1.4)$$

Put

$$S^0 = \frac{B}{\mu}. \quad (1.5)$$

We know that system (1.3) always has the disease-free equilibrium $E^0 = (S^0, 0, \dots, 0) \in \mathbb{R}^{n+2}$, where S^0 is given in (1.5). The reproduction number of system (1.3) becomes

$$R_0 = \frac{c(S^0)}{e + \sigma} = \frac{\beta S^0}{(1 + \delta b)^n (\delta + \epsilon + \gamma + \sigma)}. \quad (1.6)$$

Apart from the disease-free equilibrium E^0 , system (1.3) allows a unique endemic equilibrium $E^* = (S^*, y_1^*, \dots, y_n^*, I^*) \in \mathbb{R}^{n+2}$ under the conditions $R_0 > 1$, with $y_l^* = \hat{b}(\delta + \epsilon + \gamma + \sigma)I^* > 0$, $l = 1, 2, \dots, n$. For the case $\sigma = 0$,

Yuan and Zou [15] established a complete analysis of the global asymptotic stability of system (1.3) with a single threshold parameter R_0 .

On the other hand, if $n = 1$, then (1.3) is corresponding to the following continuous SEIS epidemic model with immigration of infectives:

$$\begin{cases} \frac{dS(t)}{dt} = B - \mu_1 S(t) - \beta S(t)I(t) + \sigma I(t), \\ \frac{dE(t)}{dt} = \beta S(t)I(t) - (\mu_2 + \lambda)E(t), \\ \frac{dI(t)}{dt} = \lambda E(t) - (\mu_3 + \sigma)I(t). \end{cases} \quad (1.7)$$

System (1.7) always has a disease-free equilibrium $P_0 = (B/\mu_1, 0, 0)$. Furthermore, if $R_0 > 1$, then system (1.7) has a unique endemic equilibrium $P_* = (S_*, E_*, I_*)$ (see Prüss *et al.* [11, Theorem 2.2]), where

$$\begin{cases} 0 < S_* = \frac{(\mu_2 + \lambda)(\mu_3 + \sigma)}{\lambda\beta} < \frac{B}{\mu_1}, & 0 < E_* = \frac{(\mu_3 + \sigma)\{\lambda\beta B - \mu_1(\mu_2 + \lambda)(\mu_3 + \sigma)\}}{\lambda\beta(\lambda\mu_3 + \sigma\mu_2 + \mu_2\mu_3)} < \frac{B}{\mu_1}, \\ 0 < I_* = \frac{\lambda\beta B - \mu_1(\mu_2 + \lambda)(\mu_3 + \sigma)}{\beta(\lambda\mu_3 + \sigma\mu_2 + \mu_2\mu_3)} < \frac{B}{\mu_1}, \end{cases} \quad (1.8)$$

and R_0 is the reproduction number defined by

$$R_0 = \frac{\lambda\beta B}{\mu_1(\mu_2 + \lambda)(\mu_3 + \sigma)}. \quad (1.9)$$

By using a geometric approach developed in Li and Muldowney [8], Fan *et al.* [2] have first proved Theorem A for the case $\mu_1 = \mu_3 \leq \mu_2$ below, and later, by using appropriate Lyapunov functionals, Prüss *et al.* [11] established complete analysis of a mathematical model for the dynamics of prion proliferation whose result is also applicable to the system (1.7) for $R_0 \leq 1$ and $R_0 > 1$ as follows (see Prüss *et al.* [11, Theorem 2.2]).

Theorem A *For system (1.7), there is precisely one disease-free equilibrium $P_0 = (\frac{B}{\mu_1}, 0, 0)$, which is globally asymptotically stable if and only if $R_0 \leq 1$. On the other hand, if $R_0 > 1$, then there is a unique endemic equilibrium $P_* = (S_*, E_*, I_*)$, which is globally asymptotically stable in $R_+^3 \setminus [\mathbf{R}_+ \times \{0\} \times \{0\}]$.*

In those cases, how to choose the discrete schemes which guarantee the global asymptotic stability for the endemic equilibrium of the models, is very important. A complete solution of this problem has been elusive until recent paper Enatsu *et al.* [1].

Motivated by the above results, in this paper we propose the following discrete epidemic model which is derived from system (1.3) by applying the well-known backward Euler method (cf. Izzo and Vecchio [5]).

$$\begin{cases} s(p+1) = s(p) + B - \delta s(p+1) - \beta s(p+1)y_{n+1}(p+1) + \sigma y_{n+1}(p+1), \\ y_1(p+1) = y_1(p) + c(s(p+1))y_{n+1}(p+1) - dy_1(p+1), \\ y_j(p+1) = y_j(p) + dy_{j-1}(p+1) - dy_j(p+1), \quad j = 2, 3, \dots, n, \\ y_{n+1}(p+1) = y_{n+1}(p) + dy_n(p+1) - (e + \sigma)y_{n+1}(p+1), \\ \text{and } i(p+1) = y_{n+1}(p+1) > 0, \quad p = 0, 1, 2, \dots, \end{cases} \quad (1.10)$$

where the initial condition of system (1.10) is

$$s(0) > 0, y_j(p) > 0, \quad \text{and} \quad j = 1, \dots, n+1. \quad (1.11)$$

Remark 1.1 To prove the positivity of $s(p), y_j(p)$, $1 \leq j \leq n$ and $i(p)$ for any $p \geq 0$ and to apply both Lyapunov functional techniques in Enatsu *et al.* [1] and a discrete time analogue to Prüss *et al.* [11], we need to use the backward Euler discretization (see, e.g., Izzo and Vecchio [5] and Izzo *et al.* [6]) which is different from that of Jang and Elaydi [7] and Sekiguchi [12]. Moreover, to consider only the positive solution $(s(p+1), y_1(p+1), y_2(p+1), \dots, i(p+1))$ in (1.10) for any obtained positive solution $(s(p), y_1(p), y_2(p), \dots, y_n(p), i(p))$, we need the restriction $i(p+1) > 0$ in (1.10), because without the condition $i(p+1) > 0$ in (1.10), there exist just two solutions $(s(p+1), y_1(p+1), y_2(p+1), \dots, y_n(p+1), i(p+1))$ of (1.10), one is $i(p+1) < 0$ and the other is $i(p+1) > 0$ for any obtained positive solution $(s(p), y_1(p), y_2(p), \dots, y_n(p), i(p))$ (see Proof of Lemma 2.1).

Using the same threshold $R_0 = \frac{c(S^0)}{e+\sigma} = \frac{\beta S^0}{(1+\delta b)^n(\delta+\epsilon+\gamma+\sigma)}$ to the continuous system (1.3), system (1.10) always has a disease-free equilibrium $E^0 = (S^0, 0, 0, \dots, 0, 0)$. Furthermore, if $R_0 > 1$, then system (1.10) has a unique endemic equilibrium $E^* = (S^*, y_1^*, y_2^*, \dots, y_n^*, I^*)$ (see Lemma 2.3). Applying the techniques of Lyapunov functions in Prüss *et al.* [11] to both cases for $R_0 \leq 1$ and $R_0 > 1$, we establish a complete analysis of the global asymptotic stability for this discrete SIR epidemic model (1.10) with immigration of infectives and latency spreading in a heterogeneous host population. In particular, we apply techniques of Lyapunov functionals in McCluskey [9] (see Lemma 5.2) to prove the global asymptotic stability for the endemic equilibrium of system (1.10) for the case $R_0 > 1$ which is simple and no longer need to use any of the theory of non-negative matrices and graph theory (cf. Guo *et al.* [4]). Moreover, we extend a simplified proof in Enatsu *et al.* [1] for the permanence of system (1.10) than Sekiguchi [12] and Sekiguchi and Ishiwata [13] to system (1.10).

Our main result in this paper, is as follows.

Theorem 1.1 *For system (1.10), there exists a unique disease-free equilibrium E^0 which is globally asymptotically stable, if and only if, $R_0 \leq 1$, and there exists a unique endemic equilibrium E^* which is globally asymptotically stable, if and only if, $R_0 > 1$.*

Remark 1.2 Theorem 1.1 for system (1.10) with $\sigma = 0$, is just a discrete analogue of the result in Yuan and Zou [15] for system (1.3).

The organization of this paper is as follows. In Section 2, we offer some basic results for system (1.10). In Section 3, we give a proof of the first part of Theorem 1.1 for $R_0 \leq 1$. In Section 4, by applying Lemmas 4.1-4.4, we offer a new proof to obtain lower positive bounds for the permanence of system (1.10) for $R_0 > 1$ (see Enatsu *et al.* [1] and cf. Thieme [14]). In Section 5, we prove the

second part of Theorem 1.1 for $R_0 > 1$ by extending a discrete time analogue of the Lyapunov function proposed by Prüss *et al.* [11] to system (1.10). Finally, short conclusions is offered in Section 6.

2. Basic properties

The following lemma is a basic result in this paper (cf. Izzo and Vecchio [5] and Izzo *et al.* [6]).

Lemma 2.1 *Let $s(p)$, $y_j(p)$, $j = 1, \dots, n$ and $y_{n+1}(p) = i(p)$ be the solutions of system (1.10) with the initial condition (1.11). Then $s(p) > 0$ and $y_j(p) > 0$, $j = 1, \dots, n+1$ for any $p \geq 0$, and (1.10) is equivalent to the following iteration system.*

$$\begin{cases} s(p+1) = \frac{B + s(p) + \sigma i(p+1)}{1 + \delta + \beta i(p+1)}, \\ y_1(p+1) = \frac{c(s(p+1))i(p+1) + y_1(p)}{1 + d}, \\ y_j(p+1) = \frac{dy_{j-1}(p+1) + y_j(p)}{1 + d}, \quad j = 2, 3, \dots, n, \\ i(p+1) = \frac{dy_n(p+1) + i(p)}{1 + e + \sigma}, \quad \text{and } i(p+1) > 0, \quad p = 0, 1, 2, \dots, \end{cases} \quad (2.1)$$

which is equivalent to

$$i(p+1) = \frac{-\tilde{B}_p + \sqrt{\tilde{B}_p^2 + 4\tilde{A}\tilde{C}_p}}{2\tilde{A}} = \frac{2\tilde{C}_p}{\tilde{B}_p + \sqrt{\tilde{B}_p^2 + 4\tilde{A}\tilde{C}_p}}, \quad (2.2)$$

where

$$\begin{cases} \tilde{A} = \beta \left((1 + e + \sigma)(1 + \delta b)^n(1 + d)^{n+1} - \delta d^n \right), \\ \tilde{B}_p = \left[(1 + \delta)(1 + e + \sigma)(1 + \delta b)^n(1 + d)^{n+1} - \beta \left\{ d^n \{ B + s(p) \} \right. \right. \\ \quad \left. \left. + (1 + \delta b)^n(1 + d)^{n+1} \left(\frac{d^n y_1(p)}{(1+d)^n} + \frac{d^{n-1} y_2(p)}{(1+d)^{n-1}} + \dots + \frac{dy_n(p)}{1+d} + i(p) \right) \right\} \right], \\ \tilde{C}_p = (1 + \delta)(1 + \delta b)^n(1 + d)^{n+1} \left(\frac{d^n y_1(p)}{(1+d)^n} + \frac{d^{n-1} y_2(p)}{(1+d)^{n-1}} + \dots + \frac{dy_n(p)}{1+d} + i(p) \right). \end{cases} \quad (2.3)$$

Proof. It is evident that the first $(n+1)$ equations of (1.10) are equivalent to the second- $(n+2)$ -th equations of (2.1). The $(n+2)$ -th equation with the first $(n+1)$ equations of (1.10) is equivalent to the first $(n+1)$ equations of (2.1)

and

$$\begin{aligned}
(1+e+\sigma)i(p+1) &= dy_n(p+1) + i(p) \\
&= d \frac{dy_{n-1}(p+1) + y_n(p)}{1+d} + i(p) \\
&= \frac{d^2 y_{n-1}(p+1)}{1+d} + \frac{dy_n(p)}{1+d} + i(p) \\
&\dots \\
&= \frac{d^n y_1(p+1)}{(1+d)^{n-1}} + \frac{d^{n-1} y_2(p)}{(1+d)^{n-1}} + \dots + \frac{dy_n(p)}{1+d} + i(p) \\
&= \frac{d^n c(s(p+1))i(p+1)}{(1+d)^{n+1}} + \frac{d^n y_1(p)}{(1+d)^n} + \frac{d^{n-1} y_2(p)}{(1+d)^{n-1}} + \dots + \frac{dy_n(p)}{1+d} + i(p) \\
&= \frac{\beta d^n}{(1+\delta b)^n (1+d)^{n+1}} \frac{B + s(p) + \sigma i(p+1)}{1 + \delta + \beta i(p+1)} i(p+1) \\
&\quad + \left(\frac{d^n y_1(p)}{(1+d)^n} + \frac{d^{n-1} y_2(p)}{(1+d)^{n-1}} + \dots + \frac{dy_n(p)}{1+d} + i(p) \right),
\end{aligned}$$

and $i(p+1) > 0$, $p = 0, 1, 2, \dots$,

which is equivalent to the following quadratic equation $P(x) = 0$ with $x = i(p+1) > 0$ such that

$$\begin{aligned}
P(x) &= (1+e+\sigma)(1+\delta b)^n (1+d)^{n+1} \{1 + \delta + \beta x\} x - \beta d^n \{B + s(p) + \sigma x\} x \\
&\quad - (1+\delta b)^n (1+d)^{n+1} \left(\frac{d^n y_1(p)}{(1+d)^n} + \frac{d^{n-1} y_2(p)}{(1+d)^{n-1}} + \dots + \frac{dy_n(p)}{1+d} + i(p) \right) (1 + \delta + \beta x) \\
&= \beta \left((1+e+\sigma)(1+\delta b)^n (1+d)^{n+1} - \sigma d^n \right) x^2 \\
&\quad + \left[(1+\delta)(1+e+\sigma)(1+\delta b)^n (1+d)^{n+1} - \beta \left\{ d^n \{B + s(p)\} \right. \right. \\
&\quad \left. \left. + (1+\delta b)^n (1+d)^{n+1} \left(\frac{d^n y_1(p)}{(1+d)^n} + \frac{d^{n-1} y_2(p)}{(1+d)^{n-1}} + \dots + \frac{dy_n(p)}{1+d} + i(p) \right) \right\} \right] x \\
&\quad - (1+\delta)(1+\delta b)^n (1+d)^{n+1} \left(\frac{d^n y_1(p)}{(1+d)^n} + \frac{d^{n-1} y_2(p)}{(1+d)^{n-1}} + \dots + \frac{dy_n(p)}{1+d} + i(p) \right)
\end{aligned} \tag{2.4}$$

and for $s(p) > 0$ and $i(p) > 0$, it is evident that $i(p+1)$ defined by the first equation of (2.1), is a unique positive solution of the quadratic equation $P(x) = 0$.

Assume that $s(p) > 0$ and $y_j(p) > 0$ $j = 1, \dots, n+1$ for some $p \geq 0$. Suppose that $s(p+1) < S^0$. Then, we have $B - \delta s(p+1) > 0$. Then, system (1.10) becomes

$$\begin{cases} (1 + \beta y_{n+1}(p+1))s(p+1) = s(p) + \{B - \delta s(p+1)\} + \sigma y_{n+1}(p+1) > 0, \\ (1 + d)y_1(p+1) = y_1(p) + c(s(p+1))y_{n+1}(p+1) > 0, \\ (1 + d)y_j(p+1) = y_j(p) + dy_{j-1}(p+1) > 0, \quad j = 2, 3, \dots, n, \\ (1 + e + \sigma)y_{n+1}(p+1) = y_{n+1}(p) + dy_n(p+1) > 0. \end{cases} \tag{2.5}$$

Then, from the first equation of (2.5), $s(p+1) > 0$. For the other case $s(p+1) \geq S^0$, it is evident that $s(p+1) > 0$. Then, from the second equation of (2.5), we have $y_1(p+1) > 0$, and similarly we obtain $y_2(p+1)$, $y_3(p+1)$, \dots , $y_{n+1}(p+1) = i(p+1) > 0$. Hence by induction of $p \geq 0$, we complete the proof of this lemma. \square

Hereafter, in order to simplify the proofs of remaining sections, let us set $y_0(p) = y_{n+2}(p) = s(p)$ and

$$\begin{cases} \underline{s} = \underline{y}_0 = \underline{y}_{n+2} = \liminf_{p \rightarrow +\infty} s(p), & \bar{s} = \bar{y}_0 = \bar{y}_{n+2} = \limsup_{p \rightarrow +\infty} s(p), \\ \underline{y}_j = \liminf_{p \rightarrow +\infty} y_j(p), & \bar{y}_j = \limsup_{p \rightarrow +\infty} y_j(p), \quad j = 1, 2, \dots, n+1, \end{cases} \quad (2.6)$$

and put

$$\begin{cases} \bar{\kappa} = (1 + \delta b)^n, & \kappa = \frac{1}{1 + \delta b}, \\ V(p) = s(p) + \bar{\kappa}\{y_1(p) + \kappa y_2(p) + \dots + \kappa^n y_{n+1}(p)\}. \end{cases} \quad (2.7)$$

Then, by (1.4),

$$\bar{\kappa} > 1 > \kappa > 0, \quad \bar{\kappa}\kappa^n = 1, \quad (1 - \kappa)d = \delta, \quad \text{and} \quad e > \delta, \quad (2.8)$$

and by (1.10) and $i(p+1) = y_{n+1}(p+1)$, one can see that

$$\begin{aligned} V(p+1) - V(p) &= B - \delta s(p+1) + \sigma y_{n+1}(p+1) \\ &\quad - \bar{\kappa}\{(1 - \kappa)dy_1(p+1) - \kappa(1 - \kappa)dy_2(p+1) + \dots \\ &\quad + \kappa^{n-1}(1 - \kappa)dy_n(p+1)\} - \bar{\kappa}\kappa^n(e + \sigma)y_{n+1}(p+1) \\ &\leq B - \delta V(p+1). \end{aligned} \quad (2.9)$$

Then, we easily obtain the following basic lemma of the boundedness of $s(p+1)$ and $y_j(p+1)$, $j = 1, 2, \dots, n+1$.

Lemma 2.2 *Let $s(p)$ and $y_j(p)$, $j = 1, \dots, n+1$ be the solutions of system (1.10) with the initial condition (1.11). Then*

$$\limsup_{p \rightarrow +\infty} V(p) \leq S^0 = \frac{B}{\delta}, \quad (2.10)$$

and

$$\bar{s} \leq S^0 = \frac{B}{\delta}, \quad \text{and} \quad \bar{y}_j \leq \frac{S^0}{\bar{\kappa}\kappa^{j-1}}, \quad j = 1, 2, \dots, n+1. \quad (2.11)$$

Proof. Let $\bar{V} = \limsup_{p \rightarrow +\infty} V(p)$. First, we suppose that $\bar{V} = +\infty$. Then, there exists a sequence $\{p_l\}_{l=1}^{\infty}$ such that $p_l < p_{l+1}$, $l = 1, 2, \dots$, $\lim_{l \rightarrow +\infty} p_l = +\infty$ and

$$V(p) < V(p_l), \quad \text{for any } p < p_l, \quad \text{and} \quad \lim_{l \rightarrow +\infty} V(p_l) = +\infty, \quad (2.12)$$

and by (2.9), we have

$$0 < V(p_l) - V(p_l - 1) \leq B - \delta V(p_l),$$

from which it holds $V(p_l) < \frac{B}{\delta} = S^0$ for any $l \geq 1$. This is a contradiction. Thus, we have $\bar{V} < +\infty$. If there exists a sequence $\{q_l\}_{l=1}^{\infty}$ such that $q_l < q_{l+1}$, $l = 1, 2, \dots$, $\lim_{l \rightarrow +\infty} q_l = +\infty$ and

$$V(q_l - 1) \leq V(q_l), \quad \text{for any } l = 1, 2, \dots, \text{ and } \lim_{l \rightarrow +\infty} V(q_l) = \bar{V}. \quad (2.13)$$

Then, similarly, we obtain that

$$B - \delta V(q_l) \geq V(q_l) - V(q_l - 1) \geq 0,$$

from which we obtain $\bar{V} \leq S^0$. For the other case that $V(p) \geq 0$ is eventually monotone decreasing. Then, there exists a $\lim_{p \rightarrow +\infty} V(p) = \bar{V} \geq 0$, and hence, $\lim_{p \rightarrow +\infty} \{V(p+1) - V(p)\} = 0$, and by (2.9), we obtain that $0 \leq B - \delta \bar{V}$, and $\bar{V} \leq S^0$, from which we get (2.11). Hence, the proof of this lemma is complete. \square

Lemma 2.3 *System (1.10) has an equilibrium $E^0 = (S^0, 0, 0, \dots, 0)$, and if $R_0 \leq 1$, then E^0 is a unique equilibrium, but if $R_0 > 1$, then there exists another equilibrium $E^* = (S^*, y_1^*, y_2^*, \dots, y_{n+1}^*)$, where $\beta S^* - \sigma > 0$ and*

$$\begin{cases} 0 < S^* = \frac{(1+\delta b)^n(e+\sigma)}{\beta} < S^0, & c(S^*) = e + \sigma, \\ y_1^* = y_2^* = \dots = y_n^* = \frac{e+\sigma}{d} y_{n+1}^*, & y_{n+1}^* = \frac{B-\mu S^*}{\beta S^* - \sigma}. \end{cases} \quad (2.14)$$

Proof. By Lemma 2.1, positivity of the sequences $\{s(p)\}_{p=1}^\infty$ and $\{y_j(p)\}_{p=1}^\infty$, $j = 1, 2, \dots, n+1$ is assured. Then, the equilibrium $\hat{E} = (\hat{S}, \hat{y}_1, \hat{y}_2, \dots, \hat{y}_{n+1})$ of (1.10) satisfy the following equations.

$$\begin{cases} B - \delta \hat{S} = (\beta \hat{S} - \sigma) \hat{y}_{n+1}, & c(\hat{S}) \hat{y}_{n+1} = d \hat{y}_1, \\ \hat{y}_1 = \hat{y}_2 = \dots = \hat{y}_n, & \text{and } d \hat{y}_n = (e + \sigma) \hat{y}_{n+1}, \end{cases} \quad (2.15)$$

that is,

$$B - \delta \hat{S} = (\beta \hat{S} - \sigma) \hat{y}_{n+1}, \quad \text{and} \quad (c(\hat{S}) - e - \sigma) \hat{y}_{n+1} = 0. \quad (2.16)$$

Then,

$$\hat{y}_{n+1} = 0, \quad \text{or} \quad c(\hat{S}) = e + \sigma. \quad (2.17)$$

If $\hat{y}_{n+1} = 0$, then by (2.15), we have that

$$\hat{S} = S^0, \quad \text{and} \quad \hat{y}_1 = \hat{y}_2 = \dots = \hat{y}_{n+1} = 0. \quad (2.18)$$

If $R_0 = \frac{c(S^0)}{e+\sigma} < 1$ then $\hat{y}_{n+1} = 0$, and by (2.15), we have (2.18). If $R_0 = 1$, then $c(S^0) = e + \sigma$. Then, by (2.17), $\hat{S} = S^0$ and by (2.15), we also have (2.18). If $R_0 > 1$, then $c(S^0) > e + \sigma$. Then, by (1.4), there exists a $0 < \frac{\sigma}{\beta} < \hat{S} = S^* = \frac{(1+\delta b)^n(e+\sigma)}{\beta} < S^0$ such that $c(S^*) = e + \sigma$, and by (2.15), it holds that $\hat{y}_{n+1} = y_{n+1}^* = \frac{B-\delta S^*}{\beta S^* - \sigma}$ and $\hat{y}_j = y_j^* = \frac{(e+\sigma)(B-\delta S^*)}{d(\beta S^* - \sigma)}$, $j = 1, 2, \dots, n$. Therefore, system (1.10) has an equilibrium E^0 . If $R_0 \leq 1$, then E^0 is a unique equilibrium, but if $R_0 > 1$, then there exists another equilibrium E^* . This completes the proof of this lemma. \square

3. Global stability of the disease-free equilibrium $E^0 = (S^0, 0, 0, \dots, 0)$ for $R_0 \leq 1$

In this section we assume that $R_0 \leq 1$ and, by applying the similar Lyapunov functional techniques to Prüss *et al.* [11] for the continuous SEIS model without delays, we prove the first part of Theorem 1.1 for system (1.10).

Proof of the first part in Theorem 1.1. Suppose that $R_0 \leq 1$. By means of a Lyapunov functional, we show that in this case the disease-free equilibrium $E^0 = (S^0, 0, 0, \dots, 0)$ is globally asymptotically stable in R_+^{n+2} . For this purpose, we first consider the case $S^0 - \frac{\sigma}{\beta} \geq 0$ and set

$$W(p) = \frac{\{s(p) - S^0\}^2}{2} + \frac{\beta S^0}{e + \sigma} \left(S^0 - \frac{\sigma}{\beta} \right) \sum_{j=1}^{n+1} y_j(p). \quad (3.1)$$

Then, we obtain

$$\begin{aligned} & W(p+1) - W(p) \\ = & \frac{\{s(p+1) - S^0\}^2}{2} - \frac{\{s(p) - S^0\}^2}{2} \\ & + \frac{\beta S^0}{e + \sigma} \left(S^0 - \frac{\delta}{\beta} \right) \left(\{c(s(p+1))y_{n+1}(p+1) - dy_1(p+1)\} \right. \\ & \left. + d \sum_{j=2}^n \{y_{j-1}(p+1) - y_j(p+1)\} + \{dy_n(p+1) - (e + \sigma)y_{n+1}(p+1)\} \right) \\ = & \frac{\{s(p+1) - S^0\}^2}{2} - \frac{\{s(p) - S^0\}^2}{2} \\ & + \frac{\beta S^0}{e + \sigma} \left(S^0 - \frac{\delta}{\beta} \right) \{c(s(p+1))y_{n+1}(p+1) - (e + \sigma)y_{n+1}(p+1)\}. \end{aligned}$$

By the relations $R_0 = \frac{\beta S^0}{(1+\delta b)^n(e+\sigma)} \leq 1$ and

$$c(s(p+1)) = \frac{\beta s(p+1)}{(1+\delta b)^n} = \frac{(e + \sigma)R_0 s(p+1)}{S^0},$$

we have

$$\begin{aligned}
W(p+1) - W(p) &= \frac{[\{s(p+1) - S^0\} + \{s(p) - S^0\}]\{s(p+1) - s(p)\}}{2} \\
&\quad + \beta \left(S^0 - \frac{\delta}{\beta} \right) \{R_0 s(p+1) - S^0\} y_{n+1}(p+1) \\
&\leq -\frac{\{s(p+1) - s(p)\}^2}{2} + \{s(p+1) - S^0\} \{s(p+1) - s(p)\} \\
&\quad + \beta \left(S^0 - \frac{\sigma}{\beta} \right) \{s(p+1) - S^0\} i(p+1) \\
&= -\frac{\{s(p+1) - s(p)\}^2}{2} + \{s(p+1) - S^0\} \\
&\quad \times \{B - \beta s(p+1) i(p+1) - \delta s(p+1) + \sigma i(p+1)\} \\
&\quad + \beta \left(S^0 - \frac{\sigma}{\beta} \right) \{s(p+1) - S^*\} i(p+1) \\
&= -\frac{\{s(p+1) - s(p)\}^2}{2} - \delta \{s(p+1) - S^0\}^2 \\
&\quad - \beta \{s(p+1) - S^0\} \{s(p+1) - \frac{\sigma}{\beta}\} i(p+1) \\
&\quad + \beta \left(S^0 - \frac{\sigma}{\beta} \right) \{s(p+1) - S^0\} i(p+1) \\
&= -\frac{\{s(p+1) - s(p)\}^2}{2} - \delta \{s(p+1) - S^0\}^2 \\
&\quad - \beta \{s(p+1) - S^0\} \left\{ \{s(p+1) - S^0\} + \left(S^0 - \frac{\sigma}{\beta} \right) \right\} i(p+1) \\
&\quad + \beta \left(S^0 - \frac{\sigma}{\beta} \right) \{s(p+1) - S^0\} i(p+1) \\
&= -\frac{\{s(p+1) - s(p)\}^2}{2} - \delta \{s(p+1) - S^0\}^2 \\
&\quad - \beta \{s(p+1) - S^0\} \left\{ \{s(p+1) - S^0\} + \left(S^0 - \frac{\sigma}{\beta} \right) \right\} i(p+1) \\
&\quad + \beta \left(S^0 - \frac{\sigma}{\beta} \right) \{s(p+1) - S^0\} i(p+1) \\
&= -\frac{\{s(p+1) - s(p)\}^2}{2} - \{\delta + \beta i(p+1)\} \{s(p+1) - S^0\}^2 \\
&\leq 0.
\end{aligned}$$

Thus, $W(p+1) \leq W(p) \leq W(0)$ for all $p \geq 0$ and $\lim_{p \rightarrow +\infty} W(p) = 0$.

If $S^0 - \frac{\sigma}{\beta} > 0$, then $\lim_{p \rightarrow +\infty} W(p) = 0$, if and only if $\lim_{p \rightarrow +\infty} s(p) = S^0 = B/\delta$ and $y_j(p+1) = 0$, $j = 1, 2, \dots, n+1$, and by $W(p+1) \leq W(p) \leq W(0)$ for all $p \geq 0$,

If $S^0 - \frac{\sigma}{\beta} = 0$, then $\lim_{p \rightarrow +\infty} W(p) = 0$, if and only if $\lim_{p \rightarrow +\infty} s(p) = S^0 = B/\delta$.

By (2.7) and (2.9), we obtain that $V(p+1) - S^0 \leq \frac{1}{1+\delta} \{V(0) - S^0\}$, $V(p+1) \leq \left(\frac{1}{1+\delta} \right)^p \{V(0) - S^0\}$ and

$$\sum_{j=1}^{n+1} \kappa^{j-1} y_j(p) \leq \left(\frac{1}{1+\delta} \right)^p \{V(0) - S^0\} - \frac{1}{\kappa} \{s(p) - S^0\}.$$

Therefore, for the case $S^0 - \frac{\sigma}{\beta} \geq 0$, the disease-free equilibrium $E^0 = (S^0, 0, 0, \dots, 0)$ is uniformly stable, and hence, it is globally asymptotically stable in R_+^{n+2} .

Next, consider the other case $S^0 - \frac{\sigma}{\beta} < 0$. Then, by Lemma 2.2, $\bar{s} \leq S^0$, and hence, there exists a sufficiently large $p_0 > 0$ such that $s(p) - \frac{\sigma}{\beta} < 0$ for any $p \geq p_0$. For $\liminf_{p \rightarrow +\infty} s(p) = \underline{s}$, we first suppose that $\underline{s} = 0$. Then, there exists a sequence $\{p_l\}_{l=1}^\infty$ such that $p_0 \leq p_l < p_{l+1}$, $l = 1, 2, \dots$, $\lim_{l \rightarrow +\infty} p_l = 0$ and

$$s(p) > s(p_l), \quad \text{for any } p < p_l, \quad \text{and} \quad \lim_{l \rightarrow +\infty} s(p_l) = 0, \quad (3.2)$$

and by (1.10), we have

$$0 > s(p_l) - s(p_l - 1) \geq B - \delta s(p_l) - \{\beta s(p_l) - \sigma\}i(p_l) \geq B - \delta s(p_l),$$

from which it holds $s(p_l) > \frac{B}{\delta} = S^0$ for any $l \geq 1$. This is a contradiction. Thus, we have $\underline{s} > 0$. If there exists a sequence $\{q_l\}_{l=1}^\infty$ such that $q_l < q_{l+1}$, $l = 1, 2, \dots$, $\lim_{l \rightarrow +\infty} q_l = +\infty$ and

$$s(q_l - 1) \geq s(q_l), \quad \text{for any } l = 1, 2, \dots, \text{ and } \lim_{l \rightarrow +\infty} s(q_l) = \underline{s}. \quad (3.3)$$

Then, similarly, we obtain that

$$B - \delta s(q_l) \leq s(q_l) - s(q_l - 1) \leq 0,$$

from which we obtain $\underline{s} \geq S^0 \geq \bar{s}$. Thus, $\lim_{p \rightarrow +\infty} s(p) = S^0$. Next, consider the other case that $s(p) > 0$ is eventually monotone increasing. Then, by Lemma 2.2, there exists a $\lim_{p \rightarrow +\infty} s(p) = \underline{s} > 0$, and hence, $\lim_{p \rightarrow +\infty} \{s(p+1) - s(p)\} = 0$, and by (1.10), we obtain that $0 \geq B - \delta \underline{s}$, and $\underline{s} \geq S^0 \geq \bar{s}$, from which we also get $\lim_{p \rightarrow +\infty} s(p) = S^0$. Then, by (2.10), we obtain that $\lim_{p \rightarrow +\infty} y_j(p) = 0$, $j = 1, 2, \dots, n+1$, which implies that the disease-free equilibrium $E^0 = (S^0, 0, 0, \dots, 0)$ is globally asymptotically stable in R_+^{n+2} . Hence, the proof of the first part of Theorem 1.1 for system (1.10) is complete. \square

4. Permanence of system (1.10) for $R_0 > 1$

In this section, we assume that $R_0 > 1$ and we will prove the permanence of system (1.10) for $R_0 > 1$.

By Lemma 2.3, the endemic equilibrium $E^* = (S^*, y_1^*, y_2^*, \dots, y_{n+1}^*)$ exists. We have basic lemmas as follow.

Lemma 4.1 *For $E^* = (S^*, y_1^*, y_2^*, \dots, y_{n+1}^*)$, it holds that*

$$\frac{c(S^*)y_{n+1}^*}{dy_1^*} = 1, \quad \frac{y_{j-1}^*}{y_j^*} = 1, \quad j = 2, 3, \dots, n, \quad \text{and} \quad \frac{dy_n^*}{(e + \sigma)y_{n+1}^*} = 1, \quad (4.1)$$

and

$$\frac{c(S^*)}{e + \sigma} = 1. \quad (4.2)$$

Proof. By (2.14), we can easily prove this lemma. \square

By Lemma 4.1, we put

$$\tilde{y}_j(p) = \frac{y_j(p)}{y_j^*}, \quad j = 1, 2, \dots, n+1. \quad (4.3)$$

Then, (1.10) becomes that

$$\begin{cases} s(p+1) - s(p) = B - \delta s(p+1) - y_{n+1}^*(\beta s(p+1) - \sigma)\tilde{y}_{n+1}(p+1), \\ \tilde{y}_1(p+1) - \tilde{y}_1(p) = \frac{c(s(p+1))y_{n+1}^*}{y_1^*}\tilde{y}_{n+1}(p+1) - d\tilde{y}_1(p+1), \\ \tilde{y}_j(p+1) - \tilde{y}_j(p) = \frac{dy_{j-1}^*}{y_j^*}\tilde{y}_{j-1}(p+1) - d\tilde{y}_j(p+1), \quad j = 2, 3, \dots, n, \\ \tilde{y}_{n+1}(p+1) - \tilde{y}_{n+1}(p) = \frac{dy_n^*}{y_{n+1}^*}\tilde{y}_n(p+1) - (e + \sigma)\tilde{y}_{n+1}(p+1), \quad p \geq 0, \end{cases}$$

and by (4.1), one can obtain that

$$\begin{cases} s(p+1) - s(p) = B - \delta s(p+1) - y_{n+1}^*(\beta s(p+1) - \sigma)\tilde{y}_{n+1}(p+1), \\ \tilde{y}_1(p+1) - \tilde{y}_1(p) = d\{\frac{c(s(p+1))}{c(S^*)}\tilde{y}_{n+1}(p+1) - \tilde{y}_1(p+1)\}, \\ \tilde{y}_j(p+1) - \tilde{y}_j(p) = d\{\tilde{y}_{j-1}(p+1) - \tilde{y}_j(p+1)\}, \quad j = 2, 3, \dots, n, \\ \tilde{y}_{n+1}(p+1) - \tilde{y}_{n+1}(p) = (e + \sigma)\{\tilde{y}_n(p+1) - \tilde{y}_{n+1}(p+1)\}, \quad p \geq 0. \end{cases} \quad (4.4)$$

Lemma 4.2 *If*

$$\min_{1 \leq j \leq n+1} \tilde{y}_j(p+1) < \min_{1 \leq j \leq n+1} \tilde{y}_j(p), \quad (4.5)$$

then

$$\tilde{y}_1(p+1) = \min_{1 \leq j \leq n+1} \tilde{y}_j(p+1) < \min_{1 \leq j \leq n+1} \tilde{y}_j(p) \quad \text{and} \quad s(p+1) < S^*. \quad (4.6)$$

Inversely,

$$\tilde{y}_1(p+1) > \min_{1 \leq j \leq n+1} \tilde{y}_j(p+1) \quad \text{or} \quad s(p+1) \geq S^*, \quad (4.7)$$

then

$$\min_{1 \leq j \leq n+1} \tilde{y}_j(p+1) \geq \min_{1 \leq j \leq n+1} \tilde{y}_j(p). \quad (4.8)$$

Proof. Assume that there exists a positive integer $1 \leq j_0 \leq n+1$ such that $0 < \tilde{y}_{j_0}(p+1) = \min_{1 \leq j \leq n+1} \tilde{y}_j(p+1) < \min_{1 \leq j \leq n+1} \tilde{y}_j(p)$. Then, $\tilde{y}_{j_0}(p+1) - \tilde{y}_{j_0}(p) < 0$ and $\tilde{y}_{j_0-1}(p+1) - \tilde{y}_{j_0}(p+1) \geq 0$. Suppose that $2 \leq j_0 \leq n+1$. Then, for the case $2 \leq j_0 \leq n$, by (4.4), we have that

$$0 > \tilde{y}_{j_0}(p+1) - \tilde{y}_{j_0}(p) = d\{\tilde{y}_{j_0-1}(p+1) - \tilde{y}_{j_0}(p+1)\} \geq 0.$$

which is a contradiction. If $j_0 = n + 1$, then by (4.4),

$$0 > \tilde{y}_{n+1}(p+1) - \tilde{y}_{n+1}(p) = (e + \sigma)\{\tilde{y}_n(p+1) - \tilde{y}_{n+1}(p+1)\} \geq 0,$$

which is also a contradiction. Thus, $j_0 = 1$. Then, $\tilde{y}_1(p+1) < \tilde{y}_{n+1}(p)$ and by the second equation of (4.4), we have

$$0 > \tilde{y}_1(p+1) - \tilde{y}_1(p) \geq d \left\{ \frac{c(s(p+1))}{c(S^*)} - 1 \right\} \tilde{y}_1(p+1),$$

from which we obtain that $s(p+1) < S^*$. Hence, if $\min_{1 \leq j \leq n+1} \tilde{y}_j(p+1) < \min_{1 \leq j \leq n+1} \tilde{y}_j(p)$, then we conclude that $s(p+1) < S^*$. Inversely, the remained part of this lemma is evident. \square

Lemma 4.3 *For (4.4), it holds that*

$$\begin{cases} \tilde{y}_j(p+1) \geq \frac{1}{1+d} \tilde{y}_j(p), & j = 1, 2, \dots, n, \\ \tilde{y}_{n+1}(p+1) \geq \frac{1}{1+e+\sigma} \tilde{y}_{n+1}(p), \\ \tilde{y}_1(p+1) \geq \frac{d}{1+d} \frac{c(s(p+1))}{c(S^*)} \tilde{y}_{n+1}(p+1), \\ \tilde{y}_j(p+1) \geq \frac{d}{1+d} \tilde{y}_{j-1}(p+1), & j = 2, \dots, n, \\ \tilde{y}_{n+1}(p+1) \geq \frac{e+\sigma}{1+e+\sigma} \tilde{y}_n(p+1), & p \geq 0. \end{cases} \quad (4.9)$$

In particular, if (4.5) holds for some $p \geq 0$, then

$$\tilde{y}_1(p+1) = \min_{1 \leq j \leq n+1} \tilde{y}_j(p+1) \geq \frac{c(s(p+1))}{c(S^*)} \tilde{y}_{n+1}(p). \quad (4.10)$$

Proof. The proof of this lemma is evident from (4.4) and (4.6). \square

Hereafter, in order to simplify the proofs of remaining sections, let us set $y_0(p) = y_{n+2}(p) = s(p)$ and

$$\begin{cases} \underline{S} = \liminf_{p \rightarrow +\infty} s(p), & \bar{S} = \limsup_{p \rightarrow +\infty} s(p), \\ \underline{y}_j = \liminf_{p \rightarrow +\infty} y_j(p), & \bar{y}_j = \limsup_{p \rightarrow +\infty} y_j(p), & j = 1, 2, \dots, n+1, \\ \underline{I} = \underline{y}_{n+1}, & \bar{I} = \bar{y}_{n+1}. \end{cases} \quad (4.11)$$

Lemma 4.4 *If $R_0 > 1$, then for any solution of system (1.10), it holds that*

$$\begin{cases} \underline{S} \geq v_0 \equiv \frac{B}{\delta + \beta B / \delta} > 0, \\ \frac{e}{d} \underline{y}_{n+1} \geq \underline{y}_n \geq \underline{y}_{n-1} \geq \dots \geq \underline{y}_1 \geq \frac{c(v_0)}{d} \underline{y}_{n+1}, \\ \text{and } \underline{I} = \underline{y}_{n+1} \geq v_{n+1}(q) \equiv \frac{d^n}{(1+d)^n} \left(\frac{1}{1+e+\sigma} \right)^{l_0(q)+1} I^* > 0, \end{cases} \quad (4.12)$$

where for any $0 < q < 1$, the integer $l_0(q) \geq 0$ is sufficiently large such that

$$S^* < \frac{B}{k_q} \left\{ 1 - \left(\frac{1}{1+k_q} \right)^{l_0(q)} \right\} \quad \text{and} \quad k_q = \delta + \beta I^*. \quad (4.13)$$

Proof. By Lemmas 2.1 and 2.2, we obtain that every sequences $\{s(p)\}_{p=0}^{\infty}$, $\{y_j(p)\}_{p=0}^{\infty}$, $j = 1, 2, \dots, n$ and $\{i(p)\}_{p=0}^{\infty}$ are positive and eventually bounded, and $\bar{S} \leq S^0 = \frac{B}{\delta}$ and $\bar{I} = \bar{y}_{n+1} \leq \frac{B}{\delta}$. Then, by the first equation of (1.10), we have that $0 \geq B - \delta \bar{S} - \beta \bar{S} \bar{I}$, from which we obtain that $\bar{S} \geq \frac{B}{\delta + \beta \bar{I}} \geq \frac{B}{\delta + \beta B / \delta}$. Thus, we obtain the first equation of (4.12).

By (1.10), we also obtain the second equation of (4.12).

Now, we show the last equation of (4.12).

First, we prove the claim that any solution $(s(p), y_1(p), y_2(p), \dots, y_n(p), i(p))$ of system (1.10) does not have the following property: for any $0 < q < 1$, there exists a nonnegative integer p_0 such that $y_j(p) \leq qy_j^*$, $j = 1, 2, \dots, n$ and $i(p) \leq qI^*$ for all $p \geq p_0$. Suppose on the contrary that there exist a solution $(s(p), y_1(p), y_2(p), \dots, y_n(p), i(p))$ of system (1.10) and a nonnegative integer p_0 such that $y_j(p) \leq qy_j^*$, $j = 1, 2, \dots, n$ and $i(p) \leq qI^*$ for all $p \geq p_0$. Then, $\tilde{y}_j(p) \leq q$, $j = 1, 2, \dots, n+1$ for all $p \geq p_0$.

Consider the sequence $\{w(p)\}_{p=0}^{\infty}$ defined by

$$w(p) = \sum_{j=1}^n \frac{\tilde{y}_j(p)}{d} + \frac{\tilde{y}_{n+1}(p)}{e + \sigma}. \quad (4.14)$$

Then, by (4.4), we have that

$$\begin{aligned} w(p+1) - w(p) &= \left(\frac{c(s(p+1))}{c(S^*)} \tilde{y}_{n+1}(p+1) - \tilde{y}_1(p+1) \right) \\ &\quad + \sum_{j=2}^n \{ \tilde{y}_{j-1}(p+1) - \tilde{y}_j(p+1) \} + \{ \tilde{y}_n(p+1) - \tilde{y}_{n+1}(p+1) \} \\ &= \left(\frac{c(s(p+1))}{c(S^*)} - 1 \right) \tilde{y}_{n+1}(p+1). \end{aligned} \quad (4.15)$$

i) Consider the case that $\{s(p)\}_{p=0}^{\infty}$ is eventually monotone increasing. Then, there is a limit of $\lim_{p \rightarrow \infty} s(p) = \hat{S} \leq \frac{B}{\delta}$. We show that $\hat{S} = S^*$.

Suppose that $\beta \hat{S} - \sigma < 0$, then $\hat{S} < \frac{\sigma}{\beta}$ and there exists an integer $p_1 \geq p_0$ such that $\beta s(p+1) - \sigma < 0$ for any $p \geq p_1$, and by the first equation of (1.10) and (4.19), we have that

$$s(p+1) - s(p) = B - \delta s(p+1) - \{\beta s(p+1) - \sigma\} i(p+1) > B - \delta s(p+1),$$

and by $p \rightarrow +\infty$, we have that $\hat{S} - \hat{S} \geq B - \delta \hat{S}$, that is, $\hat{S} \geq \frac{B}{\delta}$, which implies that $\frac{\sigma}{\beta} > \hat{S} \geq \frac{B}{\delta}$. On the other hand, since by (1.9) and Lemma 2.3, $S^* = \frac{(1+\delta b)(e+\sigma)}{\beta}$ and $R_0 = \frac{c(S^0)}{e+\sigma} = \frac{S^0}{S^*} > 1$, we have that $S^0 = \frac{B}{\delta} > S^*$, and by $S^* - \frac{\sigma}{\beta} = \frac{(1+\delta b)e + \delta b \sigma}{\beta} > 0$, we have that $\frac{B}{\delta} > S^* > \frac{\sigma}{\beta}$, which is a contradiction.

Thus, we prove that $\beta \hat{S} - \sigma \geq 0$.

Then, by the first equation of (1.10), we have that

$$\begin{aligned} B - \delta s(p+1) &> s(p+1) - s(p) = B - \delta s(p+1) - \{\beta s(p+1) - \sigma\} i(p+1) \\ &\geq B - \delta s(p+1) - \{\beta s(p+1) - \sigma\} qI^*, \end{aligned}$$

and by $p \rightarrow +\infty$, we have that $B - \delta\hat{S} \geq 0 \geq B - \delta\hat{S} - \{\beta\hat{S} - \sigma\}qI^*$, that is, $\frac{B}{\delta} > \hat{S} \geq \frac{B+\sigma qI^*}{\delta+\beta qI^*} > \frac{\sigma}{\beta}$.

Consider the following $\tilde{S} > 0$ such that

$$B - \delta\tilde{S} - \beta\tilde{S}qI^* + \sigma qI^* = 0, \quad \text{that is} \quad \tilde{S} = \frac{B + \sigma qI^*}{\delta + \beta qI^*}. \quad (4.16)$$

Then, by $\frac{\sigma}{\beta} < S^* < S^0 = \frac{B}{\delta}$, we have that $\beta B - \sigma\delta > 0$ and

$$\tilde{S} - S^* = \frac{B + \sigma qI^*}{\delta + \beta qI^*} - \frac{B + \sigma I^*}{\delta + \beta I^*} = \frac{(\beta B - \sigma\delta)(1 - q)I^*}{(\delta + \beta qI^*)(\delta + \beta I^*)} > 0. \quad (4.17)$$

Thus, we obtain that $\hat{S} \geq \tilde{S} > S^*$. Then, there exists an integer $p_1 \geq 0$ such that $s(p+1) > S^*$ for any $p \geq p_1$. Therefore, by the second part of Lemma 4.2,

$$\min_{1 \leq j \leq n+1} \tilde{y}_j(p+1) \geq \min_{1 \leq j \leq n+1} \tilde{y}_j(p), \quad \text{for any } p \geq p_1, \quad (4.18)$$

and hence, there exists a positive constant \underline{y} such that $\min_{1 \leq j \leq n+1} \tilde{y}_j(p) \geq \underline{y}$ for any $p \geq p_1$. Thus, from (4.15), we have $\lim_{p \rightarrow \infty} w(p) = +\infty$. However, by (4.14) and Lemma 2.2, it holds that there is a positive constant \bar{w} such that $w(p) \leq \bar{w}$ for any $p \geq p_1$, which leads to contradiction.

ii) Consider the case that $\{s(p)\}_{p=0}^\infty$ is not eventually monotone increasing. Then, there exists a sequence $\{p_l\}_{l=0}^\infty$ such that

$$s(p_l + 1) \leq s(p_l), \quad \text{and} \quad \lim_{l \rightarrow \infty} s(p_l + 1) = \underline{S} \leq \frac{B}{\delta}. \quad (4.19)$$

We show that $\beta\underline{S} - \sigma \geq 0$. If $\beta\underline{S} - \sigma < 0$, then there exists an integer $l_1 \geq 0$ such that $\beta s(p_l + 1) - \sigma < 0$ for any $l \geq l_1$, and by the first equation of (1.10) and (4.19), we have that

$$\begin{aligned} 0 &\geq s(p_l + 1) - s(p_l) = B - \delta s(p_l + 1) - \{\beta s(p_l + 1) - \sigma\}i(p_l + 1) \\ &> B - \delta s(p_l + 1), \end{aligned}$$

and by $l \rightarrow +\infty$, we have that $0 \geq B - \delta\underline{S}$, that is, $\underline{S} \geq \frac{B}{\delta}$, which implies that there is a limit of $\lim_{p \rightarrow \infty} s(p) = \frac{B}{\delta} > S^*$ and by the above discussion on (4.15), we conclude that $\lim_{p \rightarrow \infty} (s(p), y_1(p), y_2(p), \dots, y_n(p), i(p)) = (S^*, y_1^*, y_2^*, \dots, y_n^*, I^*)$.

This is a contradiction. Thus, we prove that $\beta\underline{S} - \sigma \geq 0$.

Then, by the first equation of (1.10) and (4.19), we have that

$$\begin{aligned} B - \delta s(p_l + 1) &\geq 0 \geq s(p_l + 1) - s(p_l) = B - \delta s(p_l + 1) - \{\beta s(p_l + 1) - \sigma\}i(p_l + 1) \\ &\geq B - \delta s(p_l + 1) - \{\beta s(p_l + 1) - \sigma\}qI^*, \end{aligned}$$

and by $l \rightarrow +\infty$, we have that $B - \delta\underline{S} \geq 0 \geq B - \delta\underline{S} - \{\beta\underline{S} - \sigma\}qI^*$, that is, $\frac{B}{\delta} > \underline{S} \geq \tilde{S} = \frac{B+\sigma qI^*}{\delta+\beta qI^*} > \frac{\sigma}{\beta}$, and hence, by $\beta B - \sigma\delta > 0$ and (4.17), we obtain

that $\underline{S} \geq \tilde{S} > S^*$, which similarly leads to contradiction by the above discussion on (4.15). Hence, the claim is proved.

Put $\tilde{y}(p) = \min_{1 \leq j \leq n+1} \{y_j(p)\}$. Then, by the claim, we are left to consider the two possibilities. First, $\tilde{y}(p) \geq q$ for all p sufficiently large. Second, we consider the case that $\tilde{y}(p)$ oscillates about q for all sufficiently large p . If the first condition that $\tilde{y}(p) \geq q$ holds for all sufficiently large p , then we get the conclusion of the proof.

For the second case that $\tilde{y}(p)$ oscillates about q for all sufficiently large p , let $p_3 < p_4$ be sufficiently large such that

$$\tilde{y}(p_3 - 1), \tilde{y}(p_4 + 1) > q, \quad \text{and} \quad \tilde{y}(p) \leq q \quad \text{for any } p_3 \leq p \leq p_4.$$

We first estimate the lower bound of $\tilde{y}_{n+1}(p)$ for $p_3 \leq p \leq p_3 + l_0(q)$. By the last equation of (4.4), we have that for $p_3 \leq p \leq p_3 + l_0(q)$,

$$\begin{aligned} \tilde{y}_{n+1}(p) &\geq \frac{1}{1+e+\sigma} \tilde{y}_{n+1}(p-1) \geq \cdots \geq \left(\frac{1}{1+e+\sigma}\right)^{p+1-p_3} \tilde{y}_{n+1}(p_3-1) \\ &> \left(\frac{1}{1+e+\sigma}\right)^{l_0(q)+1} q. \end{aligned}$$

Second, since by (1.10) and (4.13), one can obtain that for $p_3 \leq p \leq p_4$,

$$\begin{aligned} s(p+1) &\geq s(p) + B - \delta s(p+1) - \beta s(p+1) q I^* \\ &= s(p) + B - (\delta + \beta q I^*) s(p+1), \\ &= s(p) + B - k_q s(p+1), \end{aligned}$$

we obtain that

$$s(p+1) \geq \frac{s(p)}{1+k_q} + \frac{B}{1+k_q}, \quad \text{for } p_3 \leq p \leq p_4,$$

which yields

$$\begin{aligned} s(p+1) &\geq \left(\frac{1}{1+k_q}\right)^{p+1-p_3} s(p_3) + \frac{B}{1+k_q} \sum_{l=0}^{p-p_3} \left(\frac{1}{1+k_q}\right)^l \\ &\geq \frac{B}{1+k_q} \frac{1 - \left(\frac{1}{1+k_q}\right)^{p+1-p_3}}{1 - \frac{1}{1+k_q}} \\ &\geq \frac{B}{k_q} \left\{ 1 - \left(\frac{1}{1+k_q}\right)^{p+1-p_3} \right\}, \quad \text{for any } p_3 \leq p \leq p_4. \end{aligned}$$

Therefore, if $p_4 - p_3 \geq l_0(q) - 1$, then by (4.13) we have that for any $p_3 + l_0(q) \leq p + 1 \leq p_4$,

$$s(p+1) \geq s^\Delta \equiv \frac{B}{k_q} \left\{ 1 - \left(\frac{1}{1+k_q}\right)^{l_0(q)} \right\} > S^*, \quad (4.20)$$

and by the second part of Lemma 4.2, we obtain that

$$\min_{1 \leq j \leq n+1} \{y_j(p+1)\} \geq \min_{1 \leq j \leq n+1} \{y_j(p)\} \quad \text{for any } p_3 + l_0(q) \leq p \leq p_4, \quad (4.21)$$

which implies that $\min_{1 \leq j \leq n+1} \{y_j(p)\} \geq \min_{1 \leq j \leq n+1} \{y_j(p_3 + l_0(q))\}$ for any $p_3 + l_0(q) \leq p \leq p_4$. Thus, $s(p_3 + l_0(q)) \geq S^*$ and by (4.9) in Lemma 4.3, we have that

$$\begin{aligned} & \min_{1 \leq j \leq n+1} \tilde{y}_j(p_3 + l_0(q)) \\ & \geq \min \left\{ \min \left(1, \frac{d}{1+d}, \frac{d^2}{(1+d)^2}, \dots, \frac{d^{n-1}}{(1+d)^{n-1}} \right) \frac{d}{1+d} \frac{c(s(p+1))}{c(s^*)}, 1 \right\} \\ & \quad \times \tilde{y}_{n+1}(p_3 + l_0(q)), \\ & \geq \frac{d^n}{(1+d)^n} \tilde{y}_{n+1}(p_3 + l_0(q)) \\ & \geq \frac{d^n}{(1+d)^n} \left(\frac{1}{1+e+\sigma} \right)^{l_0(q)+1} q. \end{aligned}$$

Hence, we prove that

$$\underline{y}_{n+1} \geq \frac{d^n}{(1+d)^n} \left(\frac{1}{1+e+\sigma} \right)^{l_0(q)+1} q I^*.$$

Since q ($0 < q < 1$) is arbitrarily chosen, we may conclude that

$$\underline{y}_{n+1} \geq \frac{d^n}{(1+d)^n} \left(\frac{1}{1+e+\sigma} \right)^{l_0(q)+1} I^*.$$

Hence, we prove the last equation of (4.12). This completes the proof. \square

By Lemmas 2.1 and 4.4, we obtain the permanence of system (1.10).

5. Global stability of the endemic equilibrium for $R_0 > 1$

Assume $R_0 > 1$. Then, by Lemma 4.4, the system (1.10) is permanent and by Lemma 2.3, system (1.10) has a unique endemic equilibrium $E^* = (S^*, y_1^*, y_2^*, \dots, y_n^*, I^*)$. Moreover, (1.10) is equivalent to (4.4), which has a unique endemic equilibrium $\tilde{E}^* = (S^*, \tilde{y}_1^*, \tilde{y}_2^*, \dots, \tilde{y}_n^*, \tilde{y}_{n+1}^*)$ with $\tilde{y}_1^* = \tilde{y}_2^* = \dots = \tilde{y}_n^* = \tilde{y}_{n+1}^* = 1$. In the rest of this paper, we prove that the endemic equilibrium \tilde{E}^* of (4.4) is globally asymptotically stable.

By $R_0 = \frac{c(S^0)}{e+\sigma} = \frac{\beta S^0}{(1+\delta b)^n(\delta+\epsilon+\gamma+\sigma)} > 1$, we have that

$$S^0 = \frac{B}{\delta} > \frac{(1+\delta b)^n(\delta+\epsilon+\gamma+\sigma)}{\beta} = S^*$$

and

$$\beta S^* - \sigma = (1+\delta b)^n(\delta+\epsilon+\gamma+\sigma) - \sigma = (1+\delta b)^n(\delta+\epsilon+\gamma) + \{(1+\delta b)^n - 1\}\sigma > 0.$$

Define

$$\begin{cases} U_s(p) = g\left(\frac{s(p)}{S^*}\right), & U_{y_j}(p) = g\left(\frac{y_j(p)}{y_j^*}\right), \quad j = 1, 2, \dots, n \\ U_i(p) = g\left(\frac{i(p)}{I^*}\right), & \text{and } g(x) = x - 1 - \ln x \geq g(1) = 0, \quad \text{for any } x > 0. \end{cases}$$

and for simplicity, put

$$x(p+1) = \frac{s(p+1)}{S^*}, \quad z(p+1) = \frac{i(p+1)}{I^*} = \tilde{y}_{n+1}(p+1). \quad (5.1)$$

The following lemma is a key result which is a discrete version to that in Prüss *et al.* [11].

Lemma 5.1

$$\begin{aligned} & U_s(p+1) - U_s(p) \\ \leq & -\delta \frac{\{x(p+1) - 1\}^2}{x(p+1)} + \beta I^* \left(1 - \frac{1}{x(p+1)}\right) \left(1 - x(p+1) \cdot z(p+1)\right) \\ & + \frac{\sigma I^*}{S^*} \left(1 - \frac{1}{x(p+1)}\right) \{z(p+1) - 1\}, \end{aligned} \quad (5.2)$$

and

$$\begin{aligned} U_s(p+1) - U_s(p) \leq & -\{\beta I^* z(p+1) + \delta\} \frac{\{x(p+1) - 1\}^2}{x(p+1)} \\ & - \frac{(\beta S^* - \sigma) I^*}{S^*} \{z(p+1) - 1\} \left(1 - \frac{1}{x(p+1)}\right), \end{aligned} \quad (5.3)$$

and as a result, it holds that

$$\begin{aligned} & \left(1 + \frac{\sigma}{\beta S^* - \sigma}\right) S^* \{U_s(p+1) - U_s(p)\} \\ \leq & -\frac{\beta S^* \{\sigma I^* z(p+1) + \delta S^*\} \{x(p+1) - 1\}^2}{\beta S^* - \sigma} \frac{1}{x(p+1)} \\ & + \beta S^* I^* \left(1 - \frac{1}{x(p+1)}\right) \left(1 - x(p+1) \cdot z(p+1)\right). \end{aligned} \quad (5.4)$$

Proof. By (1.10), we have that

$$\begin{aligned} & U_s(p+1) - U_s(p) \\ = & \frac{s(p+1) - s(p)}{S^*} - \ln \frac{s(p+1)}{s(p)} \\ \leq & \frac{s(p+1) - s(p)}{S^*} - \frac{s(p+1) - s(p)}{s(p+1)} \\ = & \frac{s(p+1) - S^*}{S^* s(p+1)} \{s(p+1) - s(p)\} \\ = & \frac{s(p+1) - S^*}{S^* s(p+1)} \left(B - \beta s(p+1)i(p+1) - \delta s(p+1) + \sigma i(p+1)\right), \end{aligned} \quad (5.5)$$

because $\ln(1-x) \leq -x$ holds for any $x < 1$, one can obtain that

$$-\ln \frac{s(p+1)}{s(p)} = \ln \left\{1 - \left(1 - \frac{s(p)}{s(p+1)}\right)\right\} \leq -\left(1 - \frac{s(p)}{s(p+1)}\right) = -\frac{s(p+1) - s(p)}{s(p+1)}.$$

Substituting $B = \beta S^* I^* + \delta S^* - \sigma I^*$ into (5.5), we see that

$$\begin{aligned}
& U_s(p+1) - U_s(p) \\
\leq & \frac{s(p+1) - S^*}{S^* s(p+1)} \left((\beta S^* I^* + \delta S^* - \sigma I^*) - \beta s(p+1)i(p+1) - \delta s(p+1) + \sigma i(p+1) \right) \\
= & -\frac{\delta \{s(p+1) - S^*\}^2}{S^* s(p+1)} + \beta I^* \left(1 - \frac{S^*}{s(p+1)} \right) \left(1 - \frac{s(p+1)}{S^*} \cdot \frac{i(p+1)}{I^*} \right) \\
& + \frac{\sigma I^*}{S^*} \left(1 - \frac{S^*}{s(p+1)} \right) \left(\frac{i(p+1)}{I^*} - 1 \right) \\
= & -\delta \frac{\{x(p+1) - 1\}^2}{x(p+1)} + \beta I^* \left(1 - \frac{1}{x(p+1)} \right) \left(1 - x(p+1) \cdot z(p+1) \right) \\
& + \frac{\sigma I^*}{S^*} \left(1 - \frac{1}{x(p+1)} \right) \{z(p+1) - 1\}.
\end{aligned}$$

On the other hand, by $B = \beta S^* I^* + \delta S^* - \sigma I^*$, we have that

$$\begin{aligned}
& s(p+1) - s(p) \\
= & B - \beta s(p+1)i(p+1) - \delta s(p+1) + \sigma i(p+1) \\
= & (\beta S^* I^* + \delta S^* - \sigma I^*) - \beta s(p+1)i(p+1) - \delta s(p+1) + \sigma i(p+1) \\
= & -\{\beta i(p+1) + \delta\}\{s(p+1) - S^*\} - (\beta S^* - \sigma)\{i(p+1) - I^*\} \\
& + [(\beta S^* I^* + \delta S^* - \sigma I^*) \\
& - \{\beta i(p+1) + \delta\}S^* + \beta S^*\{i(p+1) - I^*\} + \sigma I^*] \\
= & -\{\beta i(p+1) + \delta\}\{s(p+1) - S^*\} - (\beta S^* - \sigma)\{i(p+1) - I^*\},
\end{aligned}$$

and hence,

$$\begin{aligned}
& U_s(p+1) - U_s(p) \\
= & \frac{s(p+1) - s(p)}{S^*} - \ln \frac{s(p+1)}{s(p)} \\
\leq & \frac{s(p+1) - s(p)}{S^*} - \frac{s(p+1) - s(p)}{s(p+1)} \\
= & \{s(p+1) - s(p)\} \left(\frac{1}{S^*} - \frac{1}{s(p+1)} \right) \\
= & -\left(\{\beta i(p+1) + \delta\}\{s(p+1) - S^*\} + (\beta S^* - \sigma)\{i(p+1) - I^*\} \right) \frac{s(p+1) - S^*}{s(p+1)S^*} \\
= & -\{\beta I^* z(p+1) + \delta\} \frac{\{x(p+1) - 1\}^2}{x(p+1)} - \frac{(\beta S^* - \sigma)I^*}{S^*} \{z(p+1) - 1\} \left(1 - \frac{1}{x(p+1)} \right).
\end{aligned}$$

Thus, we obtain (5.2) and (5.3). Moreover, from (5.2) and (5.3), we have that

$$\begin{aligned}
& \left(1 + \frac{\sigma}{\beta S^* - \sigma}\right) S^* \{U(p+1) - U(p)\} \\
\leq & \left\{ -\delta S^* \frac{\{x(p+1) - 1\}^2}{x(p+1)} + \beta S^* I^* \left(1 - \frac{1}{x(p+1)}\right) \left(1 - x(p+1) \cdot z(p+1)\right) \right. \\
& \left. + \sigma I^* \left(1 - \frac{1}{x(p+1)}\right) \{z(p+1) - 1\} \right\} \\
& - \frac{\sigma S^*}{\beta S^* - \sigma} \{\beta I^* z(p+1) + \delta\} \frac{\{x(p+1) - 1\}^2}{x(p+1)} - \sigma I^* \{z(p+1) - 1\} \left(1 - \frac{1}{x(p+1)}\right) \\
= & -S^* \left(\delta + \frac{\sigma}{\beta S^* - \sigma} \{\beta I^* z(p+1) + \delta\} \right) \frac{\{x(p+1) - 1\}^2}{x(p+1)} \\
& + \beta S^* I^* \left\{ \left(1 - \frac{1}{x(p+1)}\right) \left(1 - x(p+1) \cdot z(p+1)\right) \right\} \\
= & -\frac{\beta S^* \{\sigma I^* z(p+1) + \delta S^*\} \{x(p+1) - 1\}^2}{\beta S^* - \sigma} \frac{1}{x(p+1)} \\
& + \beta S^* I^* \left\{ \left(1 - \frac{1}{x(p+1)}\right) \left(1 - x(p+1) \cdot z(p+1)\right) \right\}.
\end{aligned}$$

Hence, the proof of this lemma is completed. \square

The following lemma plays an important role to apply techniques of equation deformation in McCluskey [9, Proof of Theorem 4.1] to the global stability analysis of endemic equilibrium for system (1.10).

Lemma 5.2 *If $R_0 > 1$, then it holds that*

$$\begin{aligned}
& \left(1 - \frac{1}{x(p+1)}\right) \left(1 - x(p+1) \cdot z(p+1)\right) \\
& + \left(1 - \frac{1}{\tilde{y}_1(p+1)}\right) \left(x(p+1) \cdot \tilde{y}_{n+1}(p+1) - \tilde{y}_1(p+1)\right) \\
& + \sum_{j=2}^{n+1} \left(1 - \frac{1}{\tilde{y}_j(p+1)}\right) \left(\tilde{y}_{j-1}(p+1) - \tilde{y}_j(p+1)\right) \\
= & -\left\{ g\left(\frac{1}{x(p+1)}\right) + g\left(\frac{x(p+1) \cdot \tilde{y}_{n+1}(p+1)}{\tilde{y}_1(p+1)}\right) + \sum_{j=2}^{n+1} g\left(\frac{\tilde{y}_{j-1}(p+1)}{\tilde{y}_j(p+1)}\right) \right\} \\
\leq & 0. \tag{5.6}
\end{aligned}$$

Proof.

$$\begin{aligned}
& \left(1 - \frac{1}{x(p+1)}\right) \left(1 - x(p+1) \cdot z(p+1)\right) \\
& + \left(1 - \frac{1}{\tilde{y}_1(p+1)}\right) \left(x(p+1) \cdot \tilde{y}_{n+1}(p+1) - \tilde{y}_1(p+1)\right) \\
& + \sum_{j=2}^{n+1} \left(1 - \frac{1}{\tilde{y}_j(p+1)}\right) \left(\tilde{y}_{j-1}(p+1) - \tilde{y}_j(p+1)\right) \\
= & \left(1 - \frac{1}{x(p+1)} - x(p+1) \cdot z(p+1) + z(p+1)\right) \\
& + x(p+1) \cdot \tilde{y}_{n+1}(p+1) - \frac{x(p+1) \cdot \tilde{y}_{n+1}(p+1)}{\tilde{y}_1(p+1)} - \tilde{y}_1(p+1) + 1 \\
& + \sum_{j=2}^{n+1} \left(\tilde{y}_{j-1}(p+1) - \frac{\tilde{y}_{j-1}(p+1)}{\tilde{y}_j(p+1)} - \tilde{y}_j(p+1) + 1\right) \\
= & \left((n+2) - \frac{1}{x(p+1)} - \frac{x(p+1) \cdot \tilde{y}_{n+1}(p+1)}{\tilde{y}_1(p+1)} - \sum_{j=2}^{n+1} \frac{\tilde{y}_{j-1}(p+1)}{\tilde{y}_j(p+1)}\right) \\
= & - \left\{ g\left(\frac{1}{x(p+1)}\right) + g\left(\frac{x(p+1) \cdot \tilde{y}_{n+1}(p+1)}{\tilde{y}_1(p+1)}\right) + \sum_{j=2}^{n+1} g\left(\frac{\tilde{y}_{j-1}(p+1)}{\tilde{y}_j(p+1)}\right) \right\} \\
\leq & 0.
\end{aligned}$$

Hence this completes the proof. \square

Proof of the second part of Theorem 1.1

Consider the following Lyapunov function (see Prüss *et al.* [11]).

$$U(p) = \frac{1}{\beta S^* I^*} \left(1 + \frac{\sigma}{\beta S^* - \sigma}\right) S^* U_s(p) + \frac{1}{d} \sum_{j=1}^n U_{y_j}(p) + \frac{1}{e + \sigma} U_i(p), \quad (5.7)$$

where

$$\begin{cases} U_s(p) = g\left(\frac{s(p)}{S^*}\right), & U_{y_j}(p) = g\left(\tilde{y}_j(p)\right), & j = 1, 2, \dots, n, \\ U_i(p) = g\left(\tilde{y}_{n+1}(p)\right), & \text{and } g(x) = x - 1 - \ln x \geq g(1) = 0, & \text{for any } x > 0. \end{cases}$$

First, we calculate $U_{y_j}(p+1) - U_{y_j}(p)$, $j = 1, 2, \dots, n+1$. By (1.4), $\frac{c(s(p+1))}{c(S^*)} =$

$\frac{s(p+1)}{S^*} = x(p+1)$. Therefore, by (4.4),

$$\begin{aligned}
& U_{y_j}(p+1) - U_{y_j}(p) \\
&= \{\tilde{y}_j(p+1) - \tilde{y}_j(p)\} - \ln \frac{\tilde{y}_j(p+1)}{\tilde{y}_j(p)} \\
&\leq \{\tilde{y}_j(p+1) - \tilde{y}_j(p)\} - \frac{\tilde{y}_j(p+1) - \tilde{y}_j(p)}{\tilde{y}_j(p+1)} \\
&= \frac{\tilde{y}_j(p+1) - 1}{\tilde{y}_j(p+1)} \{\tilde{y}_j(p+1) - \tilde{y}_j(p)\} \\
&= \begin{cases} d \left(1 - \frac{1}{\tilde{y}_1(p+1)}\right) \left(x(p+1) \cdot \tilde{y}_{n+1}(p+1) - \tilde{y}_1(p+1)\right), & \text{if } j = 1, \\ d \left(1 - \frac{1}{\tilde{y}_j(p+1)}\right) \left(\tilde{y}_{j-1}(p+1) - \tilde{y}_j(p+1)\right), & \text{if } j = 2, 3, \dots, n, \\ (e + \sigma) \left(1 - \frac{1}{\tilde{y}_{n+1}(p+1)}\right) \left(\tilde{y}_n(p+1) - \tilde{y}_{n+1}(p+1)\right), & \text{if } j = n+1. \end{cases}
\end{aligned}$$

Therefore, by Lemmas 5.1 and 5.2, we have that

$$\begin{aligned}
U(p+1) - U(p) &\leq -\frac{\{\sigma I^* z(p+1) + \mu S^*\} \{x(p+1) - 1\}^2}{I^*(\beta S^* - \sigma) x(p+1)} \\
&\quad + \left(1 - \frac{1}{x(p+1)}\right) \left(1 - x(p+1) \cdot z(p+1)\right) \\
&\quad + \left(1 - \frac{1}{\tilde{y}_1(p+1)}\right) \left(x(p+1) \cdot \tilde{y}_{n+1}(p+1) - \tilde{y}_1(p+1)\right) \\
&\quad + \sum_{j=2}^n \left(1 - \frac{1}{\tilde{y}_j(p+1)}\right) \left(\tilde{y}_{j-1}(p+1) - \tilde{y}_j(p+1)\right) \\
&\quad + \left(1 - \frac{1}{\tilde{y}_{n+1}(p+1)}\right) \left(\tilde{y}_n(p+1) - \tilde{y}_{n+1}(p+1)\right) \\
&= -\frac{\{\sigma I^* z(p+1) + \delta S^*\} \{x(p+1) - 1\}^2}{I^*(\beta S^* - \sigma) x(p+1)} \\
&\quad - \left\{ g \left(\frac{1}{x(p+1)} \right) + g \left(\frac{x(p+1) \cdot \tilde{y}_{n+1}(p+1)}{\tilde{y}_1(p+1)} \right) \right. \\
&\quad \left. + \sum_{j=2}^{n+1} g \left(\frac{\tilde{y}_{j-1}(p+1)}{\tilde{y}_j(p+1)} \right) \right\} \\
&\leq 0.
\end{aligned}$$

Hence, $U(p+1) - U(p) \leq 0$ for any $p \geq 0$. Since $U(p) \geq 0$ is monotone decreasing sequence, there is a limit $\lim_{p \rightarrow +\infty} U(p) \geq 0$. Then, $\lim_{p \rightarrow +\infty} (U(p+1) - U(p)) = 0$, from which we obtain that

$$\lim_{p \rightarrow +\infty} s(p+1) = S^*, \quad \lim_{p \rightarrow +\infty} \tilde{y}_j(p+1) = \tilde{y}_j^*, \quad j = 1, 2, \dots, n+1,$$

that is, $\lim_{p \rightarrow +\infty} (s(p), y_1(p), y_2(p), \dots, y_{n+1}(p)) = (S^*, y_1^*, y_2^*, \dots, y_{n+1}^*)$. Since $U(p) \leq U(0)$ for all $p \geq 0$ and $g(x) \geq 0$ with equality if and only if $x = 1$, E^* is uniformly stable. Hence, the proof is complete. \square

6. Conclusions

In this paper, we propose a discrete epidemic model for disease with immunity and latency spreading in a heterogeneous host population which is derived from the continuous case of model by using the well-known backward Euler method, and applying a Lyapunov functional technique which is a discrete version to that in Prüss *et al.* [11], it is shown that the global dynamics of this discrete epidemic model with latency are fully determined by a single threshold parameter.

Despite the proofs of main results in Yuan and Zou [15] make use of the theory of non-negative matrices, Lyapunov functions and a subtle grouping technique in estimating the derivatives of Lyapunov functions guided by graph theory, we apply the techniques of Lyapunov functions in McCluskey [9] and Prüss *et al.* [11] to prove the global asymptotic stability for the endemic equilibrium of system (1.10) for the case $R_0 > 1$ which is simpler and no longer needs using any of the theory of non-negative matrices and graph theory (cf. Guo *et al.* [4]). Moreover, we offer new techniques (cf. Muroya *et al.* [10]) for obtaining lower bounds for the permanence of group epidemic models which are derived by the backward Euler method from continuous group epidemic models and will be useful in applications (cf. persistence theory in dynamical systems, for example, Thieme [14], Freedman *et al.* [3] and Guo *et al.* [4]). The extension of these techniques to the other types of discrete and continuous group epidemic models will be considered in future work.

Acknowledgements

The authors wish to express their gratitude to anonymous referees for their careful readings and the insightful and constructive comments which greatly improved the quality of the paper.

References

- [1] Y. Enatsu, Y. Nakata and Y. Muroya, Global stability for a class of discrete SIR epidemic models, *Math. Biosci. Eng.* **7** (2010), 347-361.
- [2] M. Fan, M. Y. Li and K. Wang, Global stability of an SEIS epidemic model with recruitment and a varying total population size, *Math. Biosci.* **170** (2001) 199-208.
- [3] H. I. Freedman, M. X. Tang and S.G. Ruan, Uniform persistence of flows near a closed positively invariant set, *J. Dynam. Differential Equations* **6** (1994), 583-600.

- [4] H. Guo, M.Y. Li and Z. Shuai, Global stability of the endemic equilibrium of multigroup SIR epidemic models, *Can. Appl. Math. Q.* **14** (2006), 259-284.
- [5] G. Izzo and A. Vecchio, A discrete time version for models of population dynamics in the presence of an infection, *J. Comput. Appl. Math.* **210** (2007), 210-221.
- [6] G. Izzo, Y. Muroya and A. Vecchio, A general discrete time model of population dynamics in the presence of an infection, *Discrete Dyn. Nat. Soc.* 2009, Art. ID 143019, 15pp.
- [7] S. Jang and S.N. Elaydi, Difference equations from discretization of a continuous epidemic model with immigration of infectives, *MTBI Cornell University Technical Report* **92** (2004).
- [8] M. Y. Li and J. S. Muldowney, A geometric approach to global-stability problems, *SIAM J. Math. Anal.* **27** (1996) 1070-1083.
- [9] C. C. McCluskey, Complete global stability for an SIR epidemic model with delay-Distributed or discrete, *Nonl. Anal. RWA* **11** (2010) 55-59.
- [10] Y. Muroya, Y. Nakata, G. Izzo and A. Vecchio, Permanence and global stability of a class of discrete epidemic models, *Nonl. Anal. RWA* **12** (2011) 2105-2117.
- [11] J. Prüss, L. Pujo-Menjouet, G.F. Webb and R. Zacher, Analysis of a model for the dynamics of prions, *Dis. Con. Dyn. Sys. Series B*, **6** (2006), 225-235.
- [12] M. Sekiguchi, Permanence of some discrete epidemic models, *Int. J. Biomath.* **2** (2009), 443-461.
- [13] M. Sekiguchi and E. Ishiwata, Global dynamics of a discretized SIRS epidemic model with time delay, *J. Math. Anal. Appl.* **371** (2010), 195-202.
- [14] H. R. Thieme, *Mathematics in Population Biology*, Princeton University Press, Princeton, 2003.
- [15] Z. Yuan and X. Zou, Global threshold property in an epidemic model for disease with latency spreading in a heterogeneous host population, *Nonl. Anal. RWA* **11** (2010), 3479-3490.